



OPEN ACCESS

Original research

Identification of PRMT5 as a therapeutic target in cholangiocarcinoma

Jasmin Elurbide,^{1,2} Leticia Colyn,¹ Maria U Latasa,³ Iker Uriarte,^{1,2} Stefano Mariani,^{1,4} Amaya Lopez-Pascual ,^{1,5} Emiliana Valbuena,¹ Borja Castello-Urbe,¹ Robert Arnes-Benito,⁶ Elena Adan-Villaescusa,⁷ Luz A Martinez-Perez ,^{1,8} Mikel Azkargorta,⁹ Felix Elortza,¹⁰ Hanghang Wu,¹¹ Marcin Krawczyk,^{12,13} Kai Markus Schneider,¹⁴ Bruno Sangro,¹⁵ Luca Aldrighetti,¹⁶ Francesca Ratti,¹⁷ Andrea Casadei Gardini ,¹⁸ Jose J G Marin ,^{2,19} Irene Amat,^{20,21} Jesus M Urman,^{21,22} Maria Arechederra ,²³ Maria Luz Martinez-Chantar ,^{2,24} Christian Trautwein,²⁵ Meritxell Huch,⁶ Francisco Javier Cubero ,^{2,26} Carmen Berasain,²⁷ Maite G Fernandez-Barrena ,²⁸ Matias A Avila ⁷

► Additional supplemental material is published online only. To view, please visit the journal online (<https://doi.org/10.1136/gutjnl-2024-332998>).

For numbered affiliations see end of article.

Correspondence to
Professor Matias A Avila;
maavila@unav.es

JE, LC, MUL and IU are joint first authors.
MGF-B and MAA are joint senior authors.

Received 31 May 2024
Accepted 14 August 2024
Published Online First
11 September 2024



Watch Video

<https://gut.bmj.com/>


© Author(s) (or their employer(s)) 2025. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ Group.

To cite: Elurbide J, Colyn L, Latasa MU, *et al.* *Gut* 2025;**74**:116–127.

ABSTRACT

Background Cholangiocarcinoma (CCA) is a very difficult-to-treat cancer. Chemotherapies are little effective and response to immune checkpoint inhibitors is limited. Therefore, new therapeutic strategies need to be identified.

Objective We characterised the enzyme protein arginine-methyltransferase 5 (PRMT5) as a novel therapeutic target in CCA.

Design We evaluated the expression of PRMT5, its functional partner MEP50 and methylthioadenosine phosphorylase (MTAP)—an enzyme that modulates the sensitivity of PRMT5 to pharmacological inhibitors—in human CCA tissues. PRMT5-targeting drugs, currently tested in clinical trials for other malignancies, were assessed in human CCA cell lines and organoids, as well as in two immunocompetent CCA mouse models. Transcriptomic, proteomic and functional analyses were performed to explore the underlying antitumoural mechanisms.

Results PRMT5 and MEP50 proteins were correlatively overexpressed in most CCA tissues. MTAP was absent in 25% of intrahepatic CCA. PRMT5-targeting drugs markedly inhibited CCA cell proliferation, synergising with cisplatin and gemcitabine and hindered the growth of cholangiocarcinoma organoids. PRMT5 inhibition blunted the expression of oncogenic genes involved in chromatin remodelling and DNA repair, consistently inducing the formation of RNA loops and promoting DNA damage. Treatment with PRMT5-targeting drugs significantly restrained the growth of experimental CCA without adverse effects and concomitantly induced the recruitment of CD4 and CD8 T cells to shrinking tumourous lesions.

Conclusion PRMT5 and MEP50 are frequently upregulated in human CCA, and PRMT5-targeting drugs have significant antitumoural efficacy in clinically relevant CCA models. Our findings support the evaluation of PRMT5 inhibitors in clinical trials, including their combination with cytotoxic and immune therapies.

WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ Cholangiocarcinomas (CCAs) are very difficult-to-treat cancers, very resistant to chemotherapy and exhibit a very limited response to immune checkpoint inhibitors. The identification of new targets to improve the response to systemic therapies is much needed.

WHAT THIS STUDY ADDS

⇒ We demonstrate that the protein arginine-methyltransferase (PRMT5) is upregulated in a high proportion of CCA tissues. Pharmacological targeting of PRMT5 with clinically approved small molecules potentially inhibits CCA cell growth, synergising with chemotherapeutic drugs. This study shows that treatment with PRMT5 inhibitors has antitumourigenic effects in clinically relevant mouse CCA models.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ Our findings highlight the relevance of PRMT5 as a new therapeutic target in CCA and provide experimental evidence supporting the initiation of clinical trials with PRMT5 inhibitors in patients with CCA.

INTRODUCTION

Cholangiocarcinomas (CCAs) are very aggressive cancers that can be classified according to their origin within the biliary tree as intrahepatic (iCCA), perihilar and distal CCAs, with the latter two collectively known as extrahepatic CCAs (eCCA).¹ Overall, the incidence of CCA is globally increasing, mainly due to a rise in iCCA cases.^{1,2} Patients are usually diagnosed at late stages when surgical resection is not possible and chemotherapy is mostly palliative, resulting in a dramatically low 5-year overall survival between 7% and 20%.^{2–4} High-throughput sequencing analyses have elucidated




Figure 1 PRMT5 and MEP50 gene expression in human cholangiocarcinoma (CCA). (A) PRMT5 and MEP50 mRNA levels in intrahepatic CCA (iCCA) compared with normal bile ducts (NBDs) (transcriptomic dataset GSE32225), analysis of their mutual gene expression correlation and correlation with the indicated genes. (B) PRMT5 and MEP50 mRNA levels in eCCA compared with NBDs (transcriptomic dataset GSE132305), analysis of their mutual gene expression correlation and correlation with the indicated genes. (C) PRMT5 and MEP50 gene expression in an integrated dataset of iCCA samples (n=276) according to the levels of expression of the S100P gene, a marker of highly aggressive tumours. * $p < 0.05$, ** $p < 0.01$.

the molecular landscape of CCAs, identifying actionable alterations such as *FGFR2* fusions and *IDH1* mutations in iCCA, and *HER2* amplifications and *KRAS* mutations more frequently found in eCCAs, enabling the development of targeted therapies.¹ Encouraging signs of clinical activity are observed for drugs targeting *FGFR2* fusions and *IDH1* mutations in selected patients. However, the low prevalence of these mutations and the emergence of acquired drug resistance limits their effectiveness.¹⁵ On the other hand, clinical studies of immune checkpoint inhibitors (ICIs) as monotherapy have shown limited efficacy, consistent with the low frequency of mismatch repair deficiency, microsatellite instability or tumour mutational burden in biliary cancers.⁶ These outcomes have promoted the integration of cytotoxic, targeted and immune approaches in combined therapies, yielding promising results.^{5,7} It has become evident that CCAs are notoriously difficult-to-treat cancers, and therefore, while the outcomes of ongoing trials testing combination strategies are eagerly awaited new therapeutic avenues should be explored.

These novel approaches may come from mechanisms emerging as the biological makeup of CCA is unravelled. In this context, accumulating evidence indicates an important role for epigenetic changes in the induction and progression of CCA.⁸ These protumorigenic changes extend beyond mutations in genes involved in chromatin remodelling and include alterations in DNA methylation and histone post-translational modifications (PTMs).⁸ The reversible nature of epigenetic modifications makes them attractive targets for pharmacological intervention, as recently demonstrated for DNA-methyltransferase and histone lysine-methyltransferase inhibitors in experimental models of CCA.^{9–11} Arginine methylation is another type of PTM carried out by a family of nine protein arginine-methyltransferases (PRMTs) classified into three groups: type I enzymes that catalyse asymmetric arginine dimethylation (PRMT1, 2, 3, 4, 6 and 8); type II enzymes which are symmetrical dimethyltransferases (PRMT5 and 9) and a type III monomethyltransferase (PRMT7).¹² Among them, PRMT5 is highly expressed

in cancer cells and tissues, and its overexpression is directly linked to tumour progression in a broad range of solid malignancies.^{12–14} Together with methylome protein 50 (MEP50), also known as WDR77, PRMT5 forms a hetero-octameric complex that catalyses the methylation of arginine residues in histone tails, leading to the repression of tumour suppressor genes and the activation of genes involved in cell proliferation and invasion.^{12,13,15} Importantly, PRMT5 targets also comprise many non-histone proteins involved in key pathways critical for the growth and survival of cancer cells, including growth factor signalling, pre-mRNA splicing, protein translation and DNA damage repair.^{13,15–17} These observations, together with robust preclinical studies demonstrating the efficacy of PRMT5 inhibition in therapy-refractory tumours such as pancreatic cancer and hepatocellular carcinoma,^{18–21} led to the development of PRMT5 inhibitors that are showing promising preliminary results in clinical trials.^{22–24} In this study, we show that PRMT5 and MEP50 are frequently overexpressed in CCAs, and we provide experimental in vitro and in vivo evidence supporting the efficacy of PRMT5 inhibition for the treatment of this malignancy.

MATERIALS AND METHODS

Human samples

CCA tissues were obtained from patients with iCCA (n=73; 43 surgical samples, 30 needle biopsies) or eCCA (n=23) that underwent biopsies or surgical resection for diagnostic or therapeutic purposes. Tumour grade was established on anatomopathological examination of tissue sections according to the WHO Classification of Tumours, Digestive System Tumours, fifth edition. Patients or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Mouse models of CCA

The *Jnk^{Δhepa}* mice were previously described.²⁵ For CCA induction, 2-week-old male mice received 25 mg/kg (intraperitoneally)




Figure 2 Immunohistochemical analysis of PRMT5, MEP50 and MTAP proteins in human cholangiocarcinoma (CCA) tissues. (A) Representative images of PRMT5 and MEP50 immunohistochemical analyses in human iCCA and eCCA tissue samples, including iCCA samples obtained by needle biopsies. Graphs show the immunohistochemical scores for each protein in intrahepatic CCA (iCCA) and extrahepatic CCA (eCCA) tissue samples. (B) Analysis of the correlation between PRMT5 and MEP50 protein levels in iCCA tissue samples. (C) Graphs showing the distribution of PRMT5 and MEP50 scores according to tumour grade (G1–G3). (D) Representative images of MTAP immunohistochemical analyses in human iCCA and eCCA tissue samples. Graphs show the immunohistochemical scores for MTAP in iCCA and eCCA tissue samples.

of diethylnitrosamine (DEN) and from week 8 to week 25 were treated with CCl_4 (0.5 mL/kg, intraperitoneally) twice per week.^{9,26} From weeks 22 to 25, one group of mice ($n=6$) was treated with GSK3326595 (MedChemExpress, Monmouth, New Jersey, USA), at 50 mg/kg orally, daily. Control mice ($n=6$) received the same volume of vehicle, as described.²⁰ All mice were littermates.

CCA was also induced in 5-week-old C57BL/6J mice by hydrodynamic tail vein injection (HTVI) of plasmids coding for mutant TAZ (TAZS89A, 10 $\mu\text{g}/\text{mouse}$), myr-AKT (10 $\mu\text{g}/\text{mouse}$) and the sleeping beauty transposase (SB, 0.8 $\mu\text{g}/\text{mouse}$) (GenScript, Piscataway, New Jersey, USA) as described.²⁷ One group of mice ($n=12$) was treated with JNJ-64619178 (MedChemExpress) (10 mg/kg orally daily) and control mice ($n=12$) received the same volume of vehicle, as described.²¹ As reported for other HTVI oncogene-induced CCA models treatments started 2 days after HTVI,¹¹ and mice were treated for 2 or 3 weeks. C57BL/6J mice ($n=4$) without HTVI injection nor treatment were used as healthy controls and were sacrificed at same ages as the treated groups. Animals received humane care and protocols were according to the Animal Care Committee of the University of Navarra (approval #R-CP001-15GN) and the Autonomous Region of Madrid (approval #PROEX210/18 and 125.1/20) guidelines.

Additional information is provided in online supplemental materials.

RESULTS

PRMT5 and MEP50 expression in human CCA

First, we examined the expression of PRMT5 and MEP50 in well-characterised transcriptomic datasets from human CCA tissues. In

the cohort of Sia *et al.*, including iCCAs,²⁸ we observed a significant upregulation of PRMT5 and MEP50 gene expression compared with normal bile ducts (NBDs), and a direct positive correlation between the expression levels of both genes in tumour tissues (figure 1A). Here, we also found a positive correlation in PRMT5 and MEP50 expression and that of the CCA marker KRT7, and an inverse correlation with the tumour suppressor gene SOX17³ (figure 1A). In another transcriptomic dataset including eCCA samples,²⁹ we did not observe a statistically significant upregulation of PRMT5 and MEP50 in tumourous tissues versus NBDs; however, we did find a positive correlation between the expression of both genes, and with that of KRT7, MUC1 and EPCAM (for MEP50), while a negative correlation with SOX17 was observed (figure 1B). S100P expression has been robustly associated with iCCA aggressiveness.³⁰ When we analysed the integrated data from eight RNAseq studies performed in iCCA (see online supplemental materials and methods), we found that PRMT5 and MEP50 expression was significantly higher in tumours showing higher levels of S100P expression (figure 1C).

PRMT5 and MEP50 were also examined by immunohistochemistry. Both proteins were readily detected in malignant tissues, including needle iCCA biopsies (patients' and tumours' characteristics are described in online supplemental table 1), and quantitative evaluation encompassing signal intensity and the proportion of positive cells stained indicated that both iCCAs and eCCAs frequently overexpress PRMT5 and MEP50 (figure 2A) (representative images are shown in online supplemental figure 1). A positive correlation was also observed in the expression levels of both proteins in iCCAs tissues (figure 2B). Nuclear localisation of PRMT5 has been associated with worse




Figure 3 Impact of PRMT5 targeting on the growth of cholangiocarcinoma (CCA) cells. (A) CCA cell lines viability (fitness score) on CRISPR/Cas9 drop-out screen for PRMT5 and MEP50. Negative values indicate reduced survival on gene knockout. Data were retrieved from <https://score.depmap.sanger.uk/>. (B) GI₅₀ values for the PRMT5 inhibitors GSK3326595 and JNJ64619178 in the indicated CCA cell lines. (C) Immunoblot analysis of PRMT5-dependent symmetric dimethylarginine (SDMA) protein marks in control and GSK3326595 or JNJ64619178 treated HuCCCT-1 and TFK-1 cells at the indicated doses for 5 days. (D) Upper panel: GI₅₀ values for MRTX1719 in HuCCCT-1 and TFK-1 cells and analysis of MTAP protein levels by immunoblot. Lower panel: GI₅₀ values for MRTX1719 in TFK-1 control cells (TFK-1 WT) and TFK-1 cells expressing MTAP (TFK-1 MTAP), and immunoblot analyses for MTAP and SDMA protein marks in these cells. (E) Colony formation assays in the indicated cell lines treated with GSK3326595 or JNJ64619178. (F) Combination studies of the growth inhibitory effects of GSK3326595 or JNJ64619178 with cisplatin or gemcitabine in HuCCCT-1 and TFK-1 cells. Dark grey bars denote the existence of synergism at the indicated doses (combination index, CI, <1). CI was calculated as described in online supplemental materials and methods. **p*<0.05, ***p*<0.01 and ****p*<0.001 vs controls.

clinical outcomes in other solid tumours.³¹ We observed that in 63% of iCCA samples PRMT5 was found in the cells' cytoplasm and also in 100% of cells' nuclei while this was the case for 78% of eCCA samples. Only in about 4% of samples, both iCCA and eCCA, PRMT5 was localised exclusively in the cytoplasm. Noteworthy, iCCAs with higher PRMT5 and MEP50 scores were more represented among higher-grade tumours (figure 2C).

It is known that PRMT5 methyltransferase activity is highly sensitive to the intracellular levels of methylthioadenosine (MTA),³² a metabolite in the methionine salvage pathway and a S-adenosylmethionine competitive inhibitor of PRMT5.^{16 33 34} MTA is catabolised by the ubiquitous enzyme MTA-phosphorylase (MTAP), and the MTAP gene is frequently deleted in solid tumours.³⁵ MTAP loss leads to MTA accumulation and a potential improvement in the therapeutic efficacy of PRMT5-targeting drugs.^{24 33–35} Therefore, we evaluated MTAP expression by immunohistochemistry in CCA tissues. We observed that MTAP protein was undetectable in 25% of iCCA samples (9 out of 36 samples analysed) (figure 2D and online

supplemental figure 1). Together, these observations highlight the frequent overexpression of PRMT5 and MEP50 in human CCA in association with a more advanced disease, and that homozygous MTAP loss is a relatively frequent event in iCCA.

Effect of PRMT5 targeting on the growth of human CCA cells

A first indication of PRMT5 as a relevant therapeutic target in CCA was obtained by accessing data of a CRISPR/Cas9 drop-out screen in human CCA cell lines (*n*=24) (<https://score.depmap.sanger.uk/>).³⁶ We found that 83% of CCA cell lines had significantly impaired growth on genetic inactivation of PRMT5, and a similar observation was made for MEP50 (figure 3A). Next, we tested the effect of two clinically approved PRMT5 pharmacological inhibitors, GSK3326595 and JNJ64619178,²² on the growth of four well-characterised CCA cell lines.³⁷ In all cases, GI₅₀ values were in the low nanomolar range (figure 3B), and as expected both drugs significantly reduced the global levels of symmetric dimethyl arginine (SDMA) in protein extracts from CCA cells




Figure 4 Pharmacological targeting of PRMT5 inhibits the growth of human CCA tumouroids. Effect of GSK3326595 on the growth of human CCA tumouroids. Representative images of control and treated tumouroids at the end of treatments are shown along with the quantification of tumouroids growth in the indicated conditions. ** $p < 0.01$ and *** $p < 0.001$ vs controls (DMSO, vehicle-treated cultures). CCA, cholangiocarcinoma; DMSO, dimethyl sulfoxide.

(figure 3C). MRTX1719 is a small molecule that selectively binds and inhibits PRMT5 complexed with MTA. It was recently developed with the aim of increasing the efficacy of PRMT5 targeting in *MTAP*-deleted cancers in which MTA levels build up, enabling a personalised therapy and potentially reducing toxicities.²⁴ Interestingly, when we tested MRTX1719 we found a higher GI_{50} value in the *MTAP*-positive HuCCT-1 cells compared with *MTAP*-negative TFK-1 cells (figure 3D), contrary to what we observed for GSK3326595 and JNJ64619178 (figure 3B). To test if this enhanced response to MRTX1719 could indeed be attributed to the *MTAP* status, we generated a TFK-1 cell line stably expressing *MTAP*. Consistent with the inhibitory effect of MTA on PRMT5 activity,^{24 33 34} TFK-1-*MTAP* cells had higher basal SDMA levels than TFK-1-WT (figure 3D). Importantly, we found that TFK-1-WT cells were more sensitive to MRTX1719 than TFK-1-*MTAP* (figure 3D), suggesting that patients harbouring *MTAP*-deleted CCAs could benefit from this personalised therapy. In agreement with their antiproliferative activity PRMT5 inhibitors also markedly impaired the clonogenic potential of CCA cells (figure 3E). Combination strategies are actively being explored to improve the efficacy of cisplatin and gemcitabine-based CCA chemotherapy.^{5 7} We tested the effects of GSK3326595 and JNJ64619178 combined with cisplatin or gemcitabine and found a clear synergistic effect in HuCCT-1 and TFK-1 cells at most dose combinations (figure 3F). To further examine the antitumoural effects of PRMT5 inhibition, we used human CCA-derived organoids (tumouroids) that retain the histological characteristics, expression profile, genomic landscape and neoplastic behaviour of the original tumours from which they were derived.³⁸ We observed that treatment with GSK3326595 potentially reduced tumouroid growth, with the presence of smaller organoids filled with dying (figure 4). In sum, these findings indicate that PRMT5 is required for CCA cell growth, and that its inhibition markedly enhances the sensitivity of CCA cells to chemotherapeutic agents.

Mechanisms involved in the anti-CCA effect of PRMT5 inhibition

To explore the antitumoural mechanisms of PRMT5 targeting in an unbiased manner, we performed a transcriptomic analysis in the well-characterised HuCCT-1 iCCA cells^{9 10 37} on JNJ64619178 treatment. We detected 602 upregulated and 193 downregulated genes compared with controls. As expected, gene ontology biological process (GO-BP) functional classification of differentially expressed genes identified general categories related to tumourous behaviour, such as cell cycle progression and cell migration (figure 5A). Consistent with the biological

roles of PRMT5, we observed changes in genes related to chromatin regulation, mRNA splicing, DNA repair and DNA damage response.^{13 15 39} Interestingly, we also found a significant enrichment of genes involved in processes like inflammation, immune response and lipid metabolism (figure 5A). PRMT5 is critical for spliceosome assembly, and tumour cells are quite dependent on a hyperactive core splicing machinery.^{15 39} To broadly assess whether splicing impairment occurred on JNJ64619178 treatment, we analysed global alternative splicing (AS) changes in our RNAseq dataset. We identified 1374 AS events in protein-coding genes, with the majority being skipped exons and retained introns (figure 5B), two events that have shown PRMT5 dependency in other tumour types.^{17 40} GO analysis of the altered splicing events revealed significant enrichment in genes involved in cell division, the regulation of RNA splicing, inflammatory and metabolic pathways and several categories related to DNA damage and repair (figure 5C,D). Among the aberrantly spliced genes we identified the chromatin modifiers *KAT5* and *KMT5C*, critically involved in the DNA-damage response and that undergo altered splicing in *PRMT5* knockout cells¹⁷; and *POLD1*, the major catalytic DNA polymerase subunit essential for DNA replication fidelity,⁴¹ which splicing also depends on PRMT5⁴⁰ (figure 5D).

To further characterise the antineoplastic effects of PRMT5 inhibition in CCA cells, we performed complementary proteomic analyses. We identified 795 proteins that were differentially expressed (602 upregulated and 193 downregulated) in JNJ64619178-treated HuCCT-1 cells. Consistent with the RNAseq analysis, GO-BP functional classification identified enriched categories encompassing genes involved in cell cycle, chromatin regulation, RNA metabolism and splicing, DNA repair, and again immune response and lipid metabolism (figure 6A,B). We validated the effects of PRMT5 inhibition on the expression of relevant cancer-associated genes involved in chromatin remodelling, epigenetic regulation and DNA synthesis (*HELLS*, *UHRF1*, *POLD1*),^{9 41 42} antigen presentation (*NLRC5*, *B2M*, *PSMB9*, *TAP1*)⁴³ and DNA repair (*BRCA1*, *BRCA2*, *ATR*, *ATM*, *RAD51*, *RAD51AP1*)⁴⁴ in HuCCT-1, TFK1 and KMCH cells (figure 6C and online supplemental figure 2). Noteworthy, the remarkable effects of PRMT5 inhibition on *HELLS* expression may also take place at the level of protein stability, as JNJ64619178-induced *HELLS* protein downregulation was significantly attenuated by the proteasome inhibitor MG-132 (online supplemental figure 4). Moreover, *POLD1* and *HELLS*, which are important genes in DNA replication are also highly relevant in DNA repair and genome stability.^{41 42} Collectively, these observations suggest that PRMT5 activity would be critical to support mRNA splicing dynamics and DNA integrity in rapidly proliferating CCA cells, along with a potential role in tumour immune evasion.

PRMT5 inhibition promotes DNA damage in CCA cells

In view of the synergism of PRMT5 inhibitors when combined with cisplatin and gemcitabine (figure 3E), and their prominent effects on the expression of genes involved in the DNA damage response (DDR), we directly measured the levels of DNA strand breaks in a comet assay.¹⁸ As shown in figure 7A, this analysis indicated an increased frequency of DNA breaks in JNJ64619178-treated CCA cells. This response was also accompanied by the induction of apoptosis (online supplemental figure 4). DNA damage can be triggered by the formation of RNA/DNA hybrid structures known as R-loops, which lead to the exposure of single-stranded DNA and the impairment of replication fork




Figure 5 PRMT5 inhibition markedly alters the expression of genes involved in chromatin regulation, DNA damage repair, lipid metabolism and immune response and induces aberrant mRNA splicing in CCA cells. (A) Most relevant categories of differentially expressed genes identified by Gene Ontology Biological Process (GO-BP) functional classification in HuCCT-1 cells treated with JNJ64619178 (3 nM, 3 days treatment). (B) Number of alternative splicing events in protein-coding genes differentially affected by PRMT5 inhibition in HuCCT-1 cells. (C) GO-BP functional classification of the aberrant splicing events in HuCCT-1 cells treated with JNJ64619178. (D) Volcano plot representing all the significant ($p < 0.05$) splicing events. Selected genes from the indicated GO-BP functional categories are highlighted. The chromatin modifiers and DNA replication and repair genes *KAT5*, *KMT5C* and *POLD1* are identified.

progression causing replication stress.⁴⁵ By immunofluorescence analysis with the R-loop specific S9.6 antibody we found that PRMT5 inhibition leads to R-loop formation in both HuCCT-1 and TFK-1 cells (figure 7B). On DNA damage, histone 2AX (H2AX) is phosphorylated at DNA lesion sites and critically contributes to the recruitment of repair factors.⁴⁶ Interestingly, we found that the levels of H2AX protein were significantly reduced on PRMT5 inhibition (figure 7C). These findings are consistent with the role of PRMT5 in the preservation of H2AX protein stability recently described in glioblastoma cells⁴⁷ and emphasise the role of PRMT5 in maintaining genomic stability in CCA cells.

Pharmacological targeting of PRMT5 inhibits CCA development in vivo

Next, we evaluated the antitumoural efficacy of PRMT5 inhibition in two complementary mouse CCA models using two different PRMT5 pharmacological inhibitors. In the first model, mice with hepatocellular c-Jun N-terminal kinase 1/2 (*Jnk1/2*) deletion (*Jnk^{Δhepa}*) treated with CCl₄ and DEN (*Jnk^{Δhepa}*+CCl₄+DEN) develop transplantable tumours with histological and molecular features consistent with human iCCA in a context of inflammation and fibrosis.^{9 25 26} Therefore, after a single dose of DEN, 8 weeks old *Jnk^{Δhepa}* mice started to receive CCl₄ twice per week until 25 weeks of age (*Jnk^{Δhepa}*+DEN+CCl₄ mice) (figure 8A). A second group of *Jnk^{Δhepa}*+DEN+CCl₄ mice were treated with GSK3326595 for 4 weeks, from week 22 when tumours were established to week 25 (figure 8A). Immunohistochemical analysis demonstrated PRMT5 and MEP50 expression in the CK19-positive

CCA lesions (figure 8B). Tumour burden, as estimated by the liver-to-body weight ratio (liver index), was significantly reduced in treated mice, and macroscopic inspection of livers showed a marked reduction in the number of nodules (figure 8C). Consistently, histological analyses demonstrated significantly reduced CCA-like structures in GSK3326595-treated mice (figure 8D), and these structures displayed reduced tumour cell proliferation (Ki67), increased apoptosis (cleaved caspase-3) and increased DNA damage (p-H2AX) on immunohistochemical analyses (online supplemental figure 5). Interestingly, we detected increased CD4⁺ and CD8⁺ T cells infiltration in tumourous areas of GSK3326595-treated animals (figure 8E). By the end of treatments, mice receiving GSK3326595 showed a reduction in body weight. However, serum levels of hepatic enzymes, which were significantly increased in *Jnk^{Δhepa}*+DEN+CCl₄ mice, markedly diminished on GSK3326595 treatment (figure 8F).

The antineoplastic effect of PRMT5 inhibition was tested in a second clinically relevant mouse iCCA model induced by the hepatic expression of activated forms of TAZ and AKT by HTVI²⁷ (figure 9A). Here, very aggressive tumours develop rapidly after the administration of the oncogenes, replacing most of the normal parenchyma by 10 weeks postinjection, when mice die.²⁷ PRMT5 and MEP50 were clearly detected in the early lesions already present 2 weeks after HTVI and were markedly overexpressed in the extensive tumours found 1 week later (figure 9A). Next, we tested the effects of PRMT5 inhibition on iCCA development. To this end, mice injected with TAZ and AKT plasmids were treated with JNJ64619178, or vehicle, up to 3 weeks post-HTVI (figure 9A). According to




Figure 6 The effects of PRMT5 inhibition on the proteome of cholangiocarcinoma (CCA) cells are consistent with the observed transcriptional alterations. (A) Most relevant categories of differentially expressed proteins identified by Gene Ontology Biological Process (GO-BP) functional classification in HuCCT-1 cells treated with JNJ64619178 (6 nM, 4 days treatment). (B) Volcano plot representing all the significant ($p < 0.05$) differentially expressed proteins. Selected proteins included in the indicated GO-BP functional categories are identified. (C) Validation of the effects of PRMT5 inhibition on the expression of the indicated genes and proteins identified in the transcriptomic and proteomic analyses. HuCCT-1 and TFK-1 cells were treated with the indicated concentrations of JNJ64619178 for 3 or 4 days for qPCR and Western blot analyses, respectively. a, $p < 0.001$; b, $p < 0.01$; c, $p < 0.05$ vs controls.

serum transaminase levels, JNJ64619178 treatment was well tolerated, moreover, it significantly attenuated liver injury (figure 9B). At the end of treatments, extensive macroscopic tumorous lesions were appreciated in control TAZ-AKT mice while these lesions were less apparent in JNJ64619178-treated animals. The liver index confirmed a significantly reduced tumour burden (figure 9C). Consistently, histological analyses revealed a marked reduction in tumorous lesions area in mice receiving the PRMT5 inhibitor (figure 9D). These lesions also displayed reduced tumour cell proliferation (Ki67), increased apoptosis (cleaved caspase-3) and increased DNA damage (p-H2AX) (online supplemental figure 6). As observed in the *Jnk*^{Δhepa} + DEN + CCl₄ model, we clearly detected increased infiltration of CD4⁺ and CD8⁺ T cells in tumorous areas of JNJ64619178 treated mice compared with TAZ-AKT control mice (figure 9E). To better understand the antitumoural mechanisms of PRMT5 inhibition in this model, we performed transcriptomic analyses in liver tissues from TAZ-AKT mice treated with JNJ64619178 or vehicle. GO-BP functional classification of differentially expressed genes

identified general categories related to cell proliferation and cell cycle progression, consistent with the observed antineoplastic effects (figure 9F). Intriguingly, the expression of genes involved in lipid metabolism was also affected, which may be in line with the recently described role for PRMT5 in liver fat metabolism.⁴⁸ In agreement with our in vitro findings, PRMT5 targeting significantly affected gene categories related to immune and inflammatory responses, mRNA splicing and the DDR (figure 9F). Interestingly, a recent report demonstrated how aberrant TAZ/YAP expression contributes to human CCA development by promoting DNA damage and chromosomal instability.⁴⁹ Consistent with this human study, and with our transcriptomic observations, we found that the expression of several key DDR genes (*BRCA1*, *BRCA2*, *RAD51*, *POLD1*, *HELLS*) was induced in the liver of TAZ-AKT mice, and this response was blunted on JNJ64619178 treatment (figure 9G). Finally, it should be noted that the expression of MTAP was preserved in both murine CCA models (online supplemental figure 7).




Figure 7 PRMT5 inhibition induces DNA damage in cholangiocarcinoma (CCA) cells. (A) Representative images of comet assays showing levels of overall DNA strand breaks in HuCCT-1 and TFK-1 cells treated with JNJ64619178 (3 nM) for 3 days. Graph shows the quantification of the comet tail length at the level of individual cells in the number of cells indicated. (B) HuCCT-1 and TFK-1 cells were probed for R-loops using the S9.6 antibody after 3 days of treatment with JNJ64619178. Graph shows the quantification of RNA–DNA hybrids (R-loops) per nucleus. (C) Effect of JNJ64619178 on H2AX protein levels in HuCCT-1 and TFK-1 cells. Cells were treated for 3 days with the indicated JNJ64619178 concentrations. *** $p < 0.001$ vs controls (DMSO, vehicle-treated cultures). DMSO, dimethyl sulfoxide.

DISCUSSION

The identification of novel therapeutic strategies for CCA patients, particularly those that can be combined with chemotherapy or immunotherapies, is much needed. In this study, we provide evidence supporting that the arginine-methyltransferase PRMT5 could be a new pharmacological target for the systemic treatment of this extremely aggressive cancer. PRMT5 gene mutations are not commonly detected in tumours; however, overexpression of this gene is frequently observed in a broad range of malignancies.¹³ Here, we found that PRMT5 expression, and that of its functional partner MEP50, was upregulated in a significant proportion of CCAs of different anatomical origins in association with advanced stages of the disease. Previous experimental studies demonstrated that high PRMT5 levels directly contribute to malignant progression, and therefore, it should not be regarded as a mere cancer-related passenger alteration.^{13,15} Most relevantly from a therapeutic point of view, both genetic and pharmacological approaches have demonstrated that cancer cells require 2–5 fold more PRMT5 activity than normal cells, which can survive with about 15%–20% residual PRMT5 activity.³⁹ Hence, a number of ongoing clinical trials are currently testing PRMT5 small molecule inhibitors in solid and haematological malignancies.^{22,23,50} Interestingly, it was previously demonstrated that cancer cells with MTAP gene deficiency are more vulnerable to PRMT5 targeting, as MTAP-deleted cells accumulate the metabolite MTA, which is an endogenous PRMT5 inhibitor catabolised by MTAP.^{13,35,39} Homozygous MTAP deletion has been observed in various solid malignancies, and a very recent genomic investigation performed in gastrointestinal cancers found that MTAP deletion occurred in about 15% iCCAs.⁵¹ Our immunohistochemical analysis confirmed the loss of MTAP protein in more than 20% of iCCAs while additional studies including more samples are needed to establish the frequency of MTAP loss in eCCAs. Nevertheless, we show that MTAP status can be readily established in CCA tissue samples by routine IHC. This could guide the selection of patients who might respond better to PRMT5 inhibitors-based therapy,

particularly to molecules like MRTX1719²⁴ as suggested by our in vitro observations.

Our findings demonstrate that the growth and viability of human CCA cells are highly dependent on PRMT5 expression and activity. Enhanced PRMT5 activity is increasingly recognised as a critical adaptation of cancer cells to sustain growth and survival programmes through the rewiring of epigenetic mechanisms, pre-mRNA splicing, DDR and even immune evasion.^{13,15,39,43,52} In our hands, PRMT5 inhibition led to profound alterations in the transcriptome and proteome of CCA cells, suggesting that the antitumorigenic mechanisms of PRMT5 targeting are likely multifarious. We found that the expression of chromatin modifiers such as UHRF1, an oncogenic protein in CCA,⁹ and the mRNA splicing of the histone modifiers KAT5 and KMT5C,¹⁷ was markedly impaired on PRMT5 inhibition. In agreement with the strong antiproliferative effects, we also detected a significant reduction in the expression of POLD1 and HELLS genes, involved in DNA replication and chromatin remodelling, respectively. Noteworthy, HELLS protein has been recently identified as a target of PRMT5 methyltransferase activity, and HELLS arginine methylation influences its tumour-promoting activities.⁵³ Now, we also demonstrate that PRMT5 activity also contributes to preserve HELLS protein stability in CCA cells. POLD1 and HELLS are frequently overexpressed in tumours, and it is important to note that both genes also play key functions in DNA repair and genome maintenance.^{41,42} Their downregulation on PRMT5 inhibition, along with that of other important DDR genes such as BRCA1, BRCA2, ATR, RAD51AP1 and H2AX protein, suggests that PRMT5 could be a master regulator of DNA repair in CCA cells as it has been recently demonstrated for other malignancies.^{17,40,47,54,55} Indeed, we could show that PRMT5 inhibition leads to apoptosis, the accumulation of DNA strand breaks and the formation of R-loops, DNA–RNA hybrids that cause DNA damage.⁴⁵ These effects may underlie the synergistic cytotoxicity observed for PRMT5 inhibitors when combined with either gemcitabine or cisplatin and suggest that such combinatorial strategies may have a strong therapeutic




Figure 8 Antitumoural effects of PRMT5 inhibition in a mouse model of cholangiocarcinoma (CCA) development in the context of liver injury. (A) Diagram showing the *Jnk1/2^{Δhepa}*+DEN+CCl₄ (*Jnk1/2^{Δhepa}*+DC) experimental CCA model and the treatments applied (n=6 mice per group). (B) Immunohistochemical detection of PRMT5, MEP50 and CK19 proteins in liver tissue sections harbouring CCA lesions developed in this model. Representative images are shown. (C) Graphs show tumour burden, estimated as the liver-to-body weight ratio (liver index), and the quantification of nodules in the surface of the livers from the different groups of mice. Representative images of the livers of control *Jnk1/2^{Δhepa}*+DEN+CCl₄ and GSK3326595-treated mice are shown. (D) Immunohistochemical analysis of CK19 and H&E stainings showing the extent of CCA lesions in control *Jnk1/2^{Δhepa}*+DEN+CCl₄ and GSK3326595-treated mice. Representative images are shown. (E) Representative images showing the immunohistochemical detection of CD4⁺ and CD8⁺ T cells (yellow arrows), and quantification of tumour-infiltrating CD8⁺ and CD4⁺ T cells in control *Jnk1/2^{Δhepa}*+DEN+CCl₄ and GSK3326595-treated mice. (F) Body weights and serum liver parameters (AST, ALT and ALP) in the different groups of mice at the end of treatments. *p<0.05, **p<0.01 and ***p<0.001.

index in CCA.⁵⁶ Moreover, the marked increase in R-loops accumulation induced by PRMT5 targeting may also improve the efficacy of therapeutic regimens including topoisomerase inhibitors,⁵ as recently shown in a molecular subclass of glioblastoma cells prone to R-loop formation.⁵⁷ An important consequence of impaired mRNA splicing and enhanced DNA damage would be the generation of immunogenic signals and neoantigens that may induce antitumour immunity.⁵⁸ However, deficiencies in antigen processing and presentation are an independent hallmark of tumour immune evasion,⁴³ and although these mechanisms have not been thoroughly characterised in CCA they are likely to contribute to the low immunogenicity of biliary cancers.⁶ In this context, epigenetic repression of the antigen-presenting machinery has been recently described in CCA, particularly in the predominant non-inflamed subtype.⁵⁹ Interestingly, we found that PRMT5 inhibition in CCA cells induced the expression of key genes involved in antigen presentation such as *NLRC5* and *B2M*, which is consistent with the recent description of PRMT5-mediated repression of major histocompatibility complex class I-related genes in melanoma cells.⁴³

We also provide in vivo evidence of the antitumoural potential of two PRMT5 inhibitors in two complementary immunocompetent mouse models that recapitulate key molecular features of human iCCA.^{9,27} We found a marked reduction in the extent of histological lesions and overall tumour burden. Importantly, no apparent signs of toxicity were observed in drug-treated animals, on the contrary, a significant attenuation in serum markers of liver damage was observed. The in vivo antitumoural mechanisms likely include interference with the overstressed DNA repair and mRNA splicing machinery characteristic of cancer cells, which to a great extent depend on PRMT5 activity.³⁹ However, we also noticed that PRMT5 inhibition increased lymphocyte infiltration in tumourous lesions, suggesting the activation of antitumour immunity. While this response may be related to an enhanced expression in tumour cells of key genes involved in antigen presentation, as found in vitro, we cannot discard other effects such as the inhibition of T regulatory cells as previously suggested.¹³ Altogether, our study provides experimental evidence supporting the clinical evaluation of PRMT5 inhibitors in CCA patients to leverage the efficacy of




Figure 9 Antitumoural effects of PRMT5 inhibition in an aggressive cholangiocarcinoma (CCA) model triggered by hydrodynamic tail vein injection (HTVI) of activated forms of TAZ and AKT. (A) Diagram showing the TAZ-AKT model and the treatments applied (n=6 mice per group). Lower panels show representative images of the immunohistochemical analysis of PRMT5, and MEP50 in the lesions developing in TAZ-AKT mice at 2 and 3 weeks after plasmids injection, along with the demonstration of TAZ and p-AKT (Ser473) expression. Representative images are shown. (B) Serum transaminases (AST and ALT) levels in the different groups of mice. (C) Graphs show tumour burden, estimated as the liver-to-body weight ratio (liver index), and representative macroscopic images of the livers from control, TAZ-AKT mice and TAZ-AKT mice treated with JNJ64619178. (D) Representative H&E-stained liver tissue sections showing the extent of CCA lesions in TAZ-AKT and JNJ64619178-treated TAZ-AKT mice. Graph shows the quantification of tumourous areas. (E) Representative images showing the immunohistochemical detection of CD4⁺ and CD8⁺ T cells and quantification of tumour-infiltrating CD8⁺ and CD4⁺ T cells (yellow arrows) in control TAZ-AKT and JNJ64619178-treated TAZ-AKT mice. (F) Most relevant categories of differentially expressed genes identified by Gene Ontology Biological Process (GO-BP) functional classification of transcriptomic data from TAZ-AKT and JNJ64619178-treated TAZ-AKT mice livers. (G) qPCR analysis of the expression of DNA damage response (DDR)-related genes in control, TAZ-AKT and JNJ64619178-treated TAZ-AKT mice livers. *p<0.05, **p<0.01 and ***p<0.001. ALT, alanine transaminase; AST, aspartate aminotransferase.

chemotherapy or ICI-based strategies. In support of this latter possibility, a recent study demonstrated increased efficacy of ICI in combination with PRMT5 inhibitors in experimental models of breast cancer,⁶⁰ while on the other hand, the efficacy of immunotherapy was significantly lower in lung adenocarcinoma patients with high tumourous PRMT5 expression.⁶¹

Author affiliations

- ¹Hepatology Laboratory, CIMA-University of Navarra, Pamplona, Spain
- ²CIBEREHD, Madrid, Spain
- ³Hepatology and Gene Therapy, Cima. University of Navarra, Pamplona, Spain
- ⁴Oncology, University Hospital of Cagliari Department of Medicine, Cagliari, Italy
- ⁵IdiSNA, Pamplona, Spain
- ⁶Max-Planck Institute for Molecular Cell Biology and Genetics, Dresden, Germany
- ⁷Hepatology, CIMA-University of Navarra, Pamplona, Spain
- ⁸Universidad de Guadalajara Centro Universitario de Ciencias de la Salud, Guadalajara, Mexico
- ⁹Proteomics Platform, Bizkaia Science and Technology Park, Derio, Spain
- ¹⁰Proteomics Platform, CIC bioGUNE, ProteoRed-ISCIII, Bizkaia Science and Technology Park, CIC bioGUNE, Bizkaia, Spain
- ¹¹Immunology, Ophthalmology and ENT, Complutense University of Madrid Faculty of Medicine, Madrid, Spain
- ¹²Department of Gastroenterology, Hepatology and Transplant Medicine, Medical Faculty, University of Duisburg-Essen, Essen, Germany
- ¹³Laboratory of Metabolic Liver Diseases, Medical University of Warsaw, Warszawa, Poland
- ¹⁴Department of Internal Medicine III, University Hospital RWTH Aachen, Aachen, Germany
- ¹⁵Liver Unit, Dept. of Internal Medicine, Clinica Universitaria de Navarra, Pamplona, Spain
- ¹⁶IRCCS Ospedale San Raffaele, Milano, Italy
- ¹⁷Hepatobiliary surgery division, San Raffaele Hospital, Milano, Italy
- ¹⁸San Raffaele del Monte Tabor Foundation, Milano, Italy
- ¹⁹HEVEFARM, Physiology and Pharmacology, IBSAL, CIBEREhd, University of Salamanca, Salamanca, Spain
- ²⁰Department of Pathology, Navarra University Hospital Complex, Pamplona, Spain
- ²¹Instituto de Investigaciones Sanitarias de Navarra IdiSNA, Pamplona, Spain
- ²²Department of Gastroenterology and Hepatology, Navarra University Hospital Complex, Pamplona, Spain
- ²³Centro de Investigación Médica Aplicada, Pamplona, Spain
- ²⁴Liver Disease Lab, BRTA CIC bioGUNE, Centro de Investigación Biomédica en Red de Enfermedades Hepáticas y Digestivas (CIBEREhd), CICbioGUNE, Derio, Spain
- ²⁵IfADO, Dortmund, Germany
- ²⁶Immunology, Ophthalmology and ENT. Health Research Institute Gregorio Marañón (IiSGM), Complutense University of Madrid Faculty of Medicine, Madrid, Spain
- ²⁷Division of Hepatology and Gene Therapy, CIMA University of Navarra, Pamplona, Spain
- ²⁸Hepatology, FIMA, Pamplona, Spain

X Maria Arechederra @Areche_M, Maria Luz Martinez-Chantar @chantar_m, Francisco Javier Cubero @CUBEROLab and Maite G Fernandez-Barrena @Maite G Fernandez-Barrena

Acknowledgements We thank Mr Roberto Barbero for his technical support, the Morphology and Image Service at CIMA Universidad de Navarra for their help with immunohistochemical analyses, and Mr Eduardo Avila for his generous sponsorship.

Contributors JE, LC, MUL, IU, SM, AL-P, EV, BC-U, RA-B, EA-V, LAM-P, MIA, FE, MaA and HW performed experiments and interpreted primary data. MK, KMS, BS, LA, FR, ACG, JJGM, MLM-C, IA, JMU, CT, MH, FJC, CB and MAA contributed to funding and human samples acquisition. FJC and MH supervised part of the experiments. MGF-B and MAA supervised the study and analysed the data. MAA drafted the manuscript, and MGF-B and MAA revised the submitted manuscript. MAA is the guarantor.

Funding This work was supported by grants from Ministerio de Ciencia, Innovación y Universidades MCINN/AEI: PID2022-136616OB-I00/AEI/10.13039/501100011033 (MAA); PID2020-117116RB-I00 CEX2021-001136-S (MLM-C); PID2020-117941RB-I00/AEI/10.13039/501100011033 (FJC), integrados en el Plan Estatal de Investigación Científica y Técnica e Innovación, cofinanciado con Fondos FEDER 'Una manera de hacer Europa'; grant ERA-NET TRANSCAN-3 TRANSCAN2022-784-024 (MAA, MK, LA, MH and CT); grant from the Instituto de Salud Carlos III (ISCIII) AC23_1/00008 (MAA); grant from Scientific Foundation of the Spanish Association Against Cancer (FAECC) LABAE20011GARC (MGF-B) and TRNSC235657AVIL (MAA and JE); grant from Fundación Eugenio Rodríguez Pascual 2022 (MAA and MGF-B); Caixa Impulse Innovation 2023 (CI23-20155) (for MLM-C); grant EXOHEP2-CM (S2022/BMD-7409) and HORIZON-HLTH-2022-STAYHLTH-02 under agreement No 101095679 (FJC); grant from Fundación M Torres (MAA and CB); grant from

Fundación Mario Losantos (MAA and CB); grant DFG TR285/10-2 (CT). This study is based upon work from COST Action Precision-BTC-Network CA22125, supported by COST. MGF-B receives a Ramón y Cajal Program contract RYC2018-024475-1 from the Spanish Ministry of Science and Innovation. AL-P receives a Sara Borrell Contract CD22/00109 from the Spanish Ministry of Health. MA receives a post-doctoral contract POSTD18014AREC from the Asociación Española Contra el Cancer (AECC). EA-V receives a predoctoral AC-FIMA fellowship. EV receives a predoctoral fellowship PREP2022-000609 from the Ministerio de Ciencia, Innovación y Universidades project PID2022-136616OB-I00, from MICIN/AEI. BC-U receives a predoctoral Juan Serra fellowship. LC received a predoctoral fellowship from Gobierno de Navarra. SM received an ERASMUS 2021-1-IT02-KA131-HED-000008150 Placedoc fellowship. LAM-P receives a post-doctoral fellowship from CONACYT, Mexico. RA-B and MH are funded by the Max Planck Gesellschafts and the BMBF (LISYM-KREBS, DEEP-HCC).

Competing interests MAA is editor of Gut. All other authors have no conflict of interest.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants and was approved by Istituto Scientifico Romagnolo per lo Studio e la Cura dei Tumori (IRST) Meldola, Italy (ethical committee approval # IRST B050) and Hospital Universitario de Navarra Pamplona, Spain (ethical committee approval # 2015/120). Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Amaya Lopez-Pascual <http://orcid.org/0000-0001-6091-8225>
Luz A Martinez-Perez <http://orcid.org/0000-0002-9215-4458>
Andrea Casadei Gardini <http://orcid.org/0000-0002-9962-2779>
Jose J G Marin <http://orcid.org/0000-0003-1186-6849>
Maria Arechederra <http://orcid.org/0000-0002-4830-1924>
Maria Luz Martinez-Chantar <http://orcid.org/0000-0002-6446-9911>
Francisco Javier Cubero <http://orcid.org/0000-0003-1499-650X>
Maite G Fernandez-Barrena <http://orcid.org/0000-0003-0375-6236>
Matias A Avila <http://orcid.org/0000-0001-6570-3557>

REFERENCES

- 1 Ilyas SI, Affo S, Goyal L, et al. Cholangiocarcinoma - novel biological insights and therapeutic strategies. *Nat Rev Clin Oncol* 2023;20:470–86.
- 2 Alvaro D, Gores GJ, Walicki J, et al. EASL-ILCA Clinical Practice Guidelines on the management of intrahepatic cholangiocarcinoma. *J Hepatol* 2023;79:181–208.
- 3 Macias RIR, Cardinale V, Kendall TJ, et al. Clinical relevance of biomarkers in cholangiocarcinoma: critical revision and future directions. *Gut* 2022;71:1669–83.
- 4 Rushbrook SM, Kendall TJ, Zen Y, et al. British Society of Gastroenterology guidelines for the diagnosis and management of cholangiocarcinoma. *Gut* 2023;73:16–46.
- 5 Harding JJ, Khalil DN, Fabris L, et al. Rational development of combination therapies for biliary tract cancers. *J Hepatol* 2023;78:217–28.
- 6 Greten TF, Schwabe R, Bardeesy N, et al. Immunology and immunotherapy of cholangiocarcinoma. *Nat Rev Gastroenterol Hepatol* 2023;20:349–65.
- 7 Merters J, Lamarca A. Integrating cytotoxic, targeted and immune therapies for cholangiocarcinoma. *J Hepatol* 2023;78:652–7.
- 8 Rogalska-Taranta M, Andersen JB. Involvement of Epigenomic Factors in Bile Duct Cancer. *Semin Liver Dis* 2022;42:202–11.
- 9 Colyn L, Bárcena-Varela M, Álvarez-Sola G, et al. Dual Targeting of G9a and DNA Methyltransferase-1 for the Treatment of Experimental Cholangiocarcinoma. *Hepatology* 2021;73:2380–96.

- 10 Colyn L, Alvarez-Sola G, Latasa MU, *et al.* New molecular mechanisms in cholangiocarcinoma: signals triggering interleukin-6 production in tumor cells and KRAS co-opted epigenetic mediators driving metabolic reprogramming. *J Exp Clin Cancer Res* 2022;41:183.
- 11 Hu S, Molina L, Tao J, *et al.* NOTCH-YAP1/TEAD-DNMT1 Axis Drives Hepatocyte Reprogramming Into Intrahepatic Cholangiocarcinoma. *Gastroenterology* 2022;163:449–65.
- 12 Wu Q, Schapira M, Arrowsmith CH, *et al.* Protein arginine methylation: from enigmatic functions to therapeutic targeting. *Nat Rev Drug Discov* 2021;20:509–30.
- 13 Kim H, Ronai ZA. PRMT5 function and targeting in cancer. *Cell Stress* 2020;4:199–215.
- 14 Liang Z, Liu L, Wen C, *et al.* Clinicopathological and Prognostic Significance of PRMT5 in Cancers: A System Review and Meta-Analysis. *Cancer Control* 2021;28:10732748211050583.
- 15 Xu J, Richard S. Cellular pathways influenced by protein arginine methylation: Implications for cancer. *Mol Cell* 2021;81:4357–68.
- 16 Andreu-Pérez P, Esteve-Puig R, de Torre-Minguela C, *et al.* Protein arginine methyltransferase 5 regulates ERK1/2 signal transduction amplitude and cell fate through CRAF. *Sci Signal* 2011;4:ra58.
- 17 Hamard P-J, Santiago GE, Liu F, *et al.* PRMT5 Regulates DNA Repair by Controlling the Alternative Splicing of Histone-Modifying Enzymes. *Cell Rep* 2018;24:2643–57.
- 18 Wei X, Yang J, Adair SJ, *et al.* Targeted CRISPR screening identifies PRMT5 as synthetic lethality combinatorial target with gemcitabine in pancreatic cancer cells. *Proc Natl Acad Sci USA* 2020;117:28068–79.
- 19 Orben F, Lankes K, Schneeweis C, *et al.* Epigenetic drug screening defines a PRMT5 inhibitor-sensitive pancreatic cancer subtype. *JCI insight* 2022;7.
- 20 Luo Y, Gao Y, Liu W, *et al.* Myelocytomatosis-Protein Arginine N-Methyltransferase 5 Axis Defines the Tumorigenesis and Immune Response in Hepatocellular Carcinoma. *Hepatology* 2021;74:1932–51.
- 21 Brehmer D, Beke L, Wu T, *et al.* Discovery and Pharmacological Characterization of JNJ-64619178, a Novel Small-Molecule Inhibitor of PRMT5 with Potent Antitumor Activity. *Mol Cancer Ther* 2021;20:2317–28.
- 22 Feustel K, Falchook GS. Protein Arginine Methyltransferase 5 (PRMT5) Inhibitors in Oncology Clinical Trials: A review. *J Immunother Precis Oncol* 2022;5:58–67.
- 23 Vietto M, Moreno V, Spreafico A, *et al.* Phase 1 Study of JNJ-64619178, a Protein Arginine Methyltransferase 5 Inhibitor, in Advanced Solid Tumors. *Clin Cancer Res* 2023;29:3592–602.
- 24 Engstrom LD, Aranda R, Waters L, *et al.* MRTX1719 Is an MTA-Cooperative PRMT5 Inhibitor That Exhibits Synthetic Lethality in Preclinical Models and Patients with MTAP-Deleted Cancer. *Cancer Discov* 2023;13:2412–31.
- 25 Cubero FJ, Mohamed MR, Woiot MM, *et al.* Loss of c-Jun N-terminal Kinase 1 and 2 Function in Liver Epithelial Cells Triggers Biliary Hyperproliferation Resembling Cholangiocarcinoma. *Hepatology Commun* 2020;4:834–51.
- 26 Chen C, Wu H, Ye H, *et al.* Activation of the Unfolded Protein Response (UPR) Is Associated with Cholangiocellular Injury, Fibrosis and Carcinogenesis in an Experimental Model of Fibropolycystic Liver Disease. *Cancers (Basel)* 2021;14:78.
- 27 Cigliano A, Zhang S, Ribback S, *et al.* The Hippo pathway effector TAZ induces intrahepatic cholangiocarcinoma in mice and is ubiquitously activated in the human disease. *J Exp Clin Cancer Res* 2022;41.
- 28 Sia D, Hoshida Y, Villanueva A, *et al.* Integrative molecular analysis of intrahepatic cholangiocarcinoma reveals 2 classes that have different outcomes. *Gastroenterology* 2013;144:829–40.
- 29 Montal R, Sia D, Montironi C, *et al.* Molecular classification and therapeutic targets in extrahepatic cholangiocarcinoma. *J Hepatol* 2020;73:315–27.
- 30 Wu Z, Boonmars T, Nagano I, *et al.* Significance of S100P as a biomarker in diagnosis, prognosis and therapy of opisthorchiasis-associated cholangiocarcinoma. *Int J Cancer* 2016;138:396–408.
- 31 Kumar B, Yadav A, Brown NV, *et al.* Nuclear PRMT5, cyclin D1 and IL-6 are associated with poor outcome in oropharyngeal squamous cell carcinoma patients and is inversely associated with p16-status. *Oncotarget* 2017;8:14847–59.
- 32 Avila MA, García-Trevijano ER, Lu SC, *et al.* Methylthioadenosine. *Int J Biochem Cell Biol* 2004;36:2125–30.
- 33 Kryukov GV, Wilson FH, Ruth JR, *et al.* MTAP deletion confers enhanced dependency on the PRMT5 arginine methyltransferase in cancer cells. *Science* 2016;351:1214–8.
- 34 Mavrakīs KJ, McDonald ER, Schlabach MR, *et al.* Disordered methionine metabolism in MTAP/CDKN2A-deleted cancers leads to dependence on PRMT5. *Science* 2016;351:1208–13.
- 35 Fan N, Zhang Y, Zou S. Methylthioadenosine phosphorylase deficiency in tumors: A compelling therapeutic target. *Front Cell Dev Biol* 2023;11.
- 36 Behan FM, Iorio F, Picco G, *et al.* Prioritization of cancer therapeutic targets using CRISPR-Cas9 screens. *Nature New Biol* 2019;568:511–6.
- 37 Zach S, Birgin E, Rückert F. Primary cholangiocellular carcinoma cell lines. *J Stem Cell Res Transplant* 2015;2:1013.
- 38 Broutier L, Mastrogianni G, Versteegen MM, *et al.* Human primary liver cancer-derived organoid cultures for disease modeling and drug screening. *Nat Med* 2017;23:1424–35.
- 39 Guccione E, Schwarz M, Di Tullio F, *et al.* Cancer synthetic vulnerabilities to protein arginine methyltransferase inhibitors. *Curr Opin Pharmacol* 2021;59:33–42.
- 40 Carter J, Hulse M, Sivakumar M, *et al.* PRMT5 Inhibitors Regulate DNA Damage Repair Pathways in Cancer Cells and Improve Response to PARP Inhibition and Chemotherapies. *Cancer Res Commun* 2023;3:2233–43.
- 41 Gola M, Stefaniak P, Godlewski J, *et al.* Prospects of POLD1 in Human Cancers: A Review. *Cancers (Basel)* 2023;15:1905.
- 42 Peixoto E, Khan A, Lewis ZA, *et al.* The Chromatin Remodeler HELLS: A New Regulator in DNA Repair, Genome Maintenance, and Cancer. *Int J Mol Sci* 2022;23:9313.
- 43 Kim H, Kim H, Feng Y, *et al.* PRMT5 control of cGAS/STING and NLR3 pathways defines melanoma response to antitumor immunity. *Sci Transl Med* 2020;12.
- 44 Mavroei D, Georganta A, Panagiotou E, *et al.* Targeting ATR Pathway in Solid Tumors: Evidence of Improving Therapeutic Outcomes. *Int J Mol Sci* 2024;25:2767.
- 45 Li F, Zafar A, Luo L, *et al.* R-Loops in Genome Instability and Cancer. *Cancers (Basel)* 2023;15:4986.
- 46 Fragkos M, Choleza M, Papadopoulos P. The Role of γH2AX in Replication Stress-induced Carcinogenesis: Possible Links and Recent Developments. *CDP* 2023;3:639–48.
- 47 Du C, Hansen LJ, Singh SX, *et al.* A PRMT5-RNF168-SMURF2 Axis Controls H2AX Proteostasis. *Cell Rep* 2019;28:3199–211.
- 48 Hou A, Xu X, Zhang Y, *et al.* Excessive fatty acids activate PRMT5/MDM2/Drosha pathway to regulate miRNA biogenesis and lipid metabolism. *Liver Int* 2024;44:1634–50.
- 49 Tóth M, Wehling L, Thiess L, *et al.* Co-expression of YAP and TAZ associates with chromosomal instability in human cholangiocarcinoma. *BMC Cancer* 2021;21:1079.
- 50 Rodon J, Rodriguez E, Maitland ML, *et al.* A phase I study to evaluate the safety, pharmacokinetics, and pharmacodynamics of PF-06939999 (PRMT5 inhibitor) in patients with selected advanced or metastatic tumors with high incidence of splicing factor gene mutations. *ESMO Open* 2024;9:102961.
- 51 Ngoi NYL, Tang T-Y, Gaspar CF, *et al.* Methylthioadenosine Phosphorylase Genomic Loss in Advanced Gastrointestinal Cancers. *Oncologist* 2024;29:493–503.
- 52 Abe Y, Sano T, Tanaka N. The Role of PRMT5 in Immuno-Oncology. *Genes (Basel)* 2023;14:678.
- 53 Liu N, Yang R, Shi Y, *et al.* The cross-talk between methylation and phosphorylation in lymphoid-specific helicase drives cancer stem-like properties. *Sig Transduct Target Ther* 2020;5.
- 54 Li Y, Dobrolecki LE, Sallas C, *et al.* PRMT blockade induces defective DNA replication stress response and synergizes with PARP inhibition. *Cell Rep Med* 2023;4:101326.
- 55 Kalev P, Hyer ML, Gross S, *et al.* MAT2A Inhibition Blocks the Growth of MTAP-Deleted Cancer Cells by Reducing PRMT5-Dependent mRNA Splicing and Inducing DNA Damage. *Cancer Cell* 2021;39:209–24.
- 56 Gönül Geyik Ö, Anichini G, Ulukaya E, *et al.* DNA Damage Response Inhibitors in Cholangiocarcinoma: Current Progress and Perspectives. *Cells* 2022;11:1463.
- 57 Struve N, Hoffer K, Weik A-S, *et al.* Increased replication stress and R-loop accumulation in EGFRVIII-expressing glioblastoma present new therapeutic opportunities. *Neurooncol Adv* 2022;4:vdab180.
- 58 Kornepati AVR, Rogers CM, Sung P, *et al.* The complementarity of DDR, nucleic acids and anti-tumour immunity. *Nature New Biol* 2023;619:475–86.
- 59 Nishida N, Aoki T, Morita M, *et al.* Non-Inflamed Tumor Microenvironment and Methylation/Downregulation of Antigen-Presenting Machinery in Cholangiocarcinoma. *Cancers (Basel)* 2023;15:2379.
- 60 Wang Z, Li R, Hou N, *et al.* PRMT5 reduces immunotherapy efficacy in triple-negative breast cancer by methylating KEAP1 and inhibiting ferroptosis. *J Immunother Cancer* 2023;11:e006890.
- 61 Wang J, Wu M, Sun J, *et al.* Pan-cancer analysis identifies protein arginine methyltransferases PRMT1 and PRMT5 and their related signatures as markers associated with prognosis, immune profile, and therapeutic response in lung adenocarcinoma. *Heliyon* 2023;9:e22088.