

Access to and quality of elective care: a prospective cohort study using hernia surgery as a tracer condition in 83 countries



NIHR Global Health Research Unit on Global Surgery*

Summary

Background Timely and safe elective health care facilitates return to normal activities for patients and prevents emergency admissions. Surgery is a cornerstone of elective care and relies on complex pathways. This study aimed to take a whole-system approach to evaluating access to and quality of elective health care globally, using inguinal hernia as a tracer condition.

Methods This was a prospective, international, cohort study conducted between Jan 30 and May 21, 2023, in which any hospital performing inguinal hernia repairs was eligible to take part. Consecutive patients of any age undergoing primary inguinal hernia repair were included. A measurement set mapped to the attributes of WHO's Health System Building Blocks was defined to evaluate access (emergency surgery rates, bowel resection rates, and waiting times) and quality (mesh use, day-case rates, and postoperative complications). These were compared across World Bank income groups (high-income, upper-middle-income, lower-middle-income, and low-income countries), adjusted for hospital and country. Factors associated with postoperative complications were explored with a three-level multilevel logistic regression model.

Findings 18 058 patients from 640 hospitals in 83 countries were included, of whom 1287 (7·1%) underwent emergency surgery. Emergency surgery rates increased from high-income to low-income countries (6·8%, 9·7%, 11·4%, 14·2%), accompanied by an increase in bowel resection rates (1·2%, 1·4%, 2·3%, 4·2%). Overall waiting times for elective surgery were similar around the world (median 8·0 months from symptoms to surgery), largely because of delays between symptom onset and diagnosis rather than waiting for treatment. In 14 768 elective operations in adults, mesh use decreased from high-income to low-income countries (97·6%, 94·3%, 80·6%, 61·0%). In patients eligible for day-case surgery ($n=12\,658$), day-case rates were low and variable (50·0%, 38·0%, 42·1%, 44·5%). Complications occurred in 2415 (13·4%) of 18 018 patients and were more common after emergency surgery (adjusted odds ratio 2·06, 95% CI 1·72–2·46) and bowel resection (1·85, 1·31–2·63), and less common after day-case surgery (0·39, 0·34–0·44).

Interpretation This study demonstrates that elective health care is essential to preventing over-reliance on emergency systems. We identified actionable targets for system strengthening: clear referral pathways and increasing mesh repair in lower-income settings, and boosting day-case surgery in all income settings. These measures might strengthen non-surgical pathways too, reducing the burden on society and health services.

Funding NIHR Global Health Research Unit on Global Surgery and Portuguese Hernia and Abdominal Wall Society (Sociedade Portuguesa de Hernia e Parede Abdominal).

Copyright © 2024 The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY 4.0 license.

Introduction

Elective health care remains the cornerstone to well functioning health systems,¹ with common examples including diagnostics, surgery, and cardiovascular disease optimisation.² Across a very wide range of health-care systems, elective care allows a fast recovery to normal activities, reduces the need for complex emergency care, and reduces health-care costs.¹ During and after the COVID-19 pandemic, elective care was deprioritised and continues to struggle to recover, particularly in terms of access and volume.^{3,4} Continual deprioritisation is leading to a downward spiral of increasing emergency admissions

and further stress on the remaining elective capacity, with crisis management of emergency conditions fast becoming the norm.^{5,6}

Elective health care is a broad umbrella that includes a wide range of diseases, diagnostics, and treatments, meaning that any research needs to be focused.¹ A tracer condition is a common condition where diagnosis and management are established, and a lack of treatment can cause harm.⁷ A tracer condition that is common to countries around the world and able to describe a whole-system pathway, would be able to assess post-COVID-19 elective care and identify targets for improvement.

Lancet Glob Health 2024

Published Online

May 23, 2024

[https://doi.org/10.1016/S2214-109X\(24\)00142-6](https://doi.org/10.1016/S2214-109X(24)00142-6)

See Online/Comment

[https://doi.org/10.1016/S2214-109X\(24\)00142-6](https://doi.org/10.1016/S2214-109X(24)00142-6)

*A full list of collaborating authors is shown in appendix 1 (pp 1–34)

Correspondence to:

Dr Maria Picciochi, NIHR Global Health Research Unit on Global Surgery, Institute of Applied Health Research, University of Birmingham, Birmingham B15 2TH, UK
m.picciochi@bham.ac.uk

or
Prof Aneel Bhanu, NIHR Global Health Research Unit on Global Surgery, Institute of Applied Health Research, University of Birmingham, Birmingham B15 2TH, UK
a.a.bhanu@bham.ac.uk

See Online for appendix 1

Research in context

Evidence before this study

Elective care is cost-effective, ensures patients return to normal activities quickly, and prevents emergency admissions. Partly through COVID-19 shutdowns in elective care and the subsequent slow recovery, health care is facing an international crisis, with emergency care becoming the norm. It is creating a downward spiral that is further reducing elective capacity. We planned to use inguinal hernia care as a tracer condition (ie, a condition for which there are accepted definitions and proven, cost-effective treatments) to assess elective health care. We searched PubMed for global studies linking assessment of health care and patient-level data. We used the search terms “elective healthcare”, “elective surgery”, and “inguinal hernia repair”, as well as related terms. We did not find any study with data collected from patients with inguinal hernias and health system evaluation. We also did not find any study assessing waiting times for elective surgery at a global scale before the pandemic. However, we identified inguinal hernia as part of the World Bank’s Essential Surgery Package and the 2023 HerniaSurge Collaborative guidelines, which set acceptable and achievable standards around the world. We identified the Surgical Preparedness Index, which helps hospitals establish resilient elective systems but showed wide global variation in preparedness. We reviewed the *Lancet* Commission on Surgery (2015) and the *Lancet* Commission on Diagnostics (2021), both of which advocate for stronger elective systems, and called for actionable research.

Added value of this study

This study used hernia repair as a tracer condition for elective health care. It showed that inguinal hernias are mostly a disease of working age patients around the world, and when limited to the inguinal region, are treatable with simple, elective, day-case surgery. If neglected, the need for more complex emergency surgery increases substantially, as do major life-threatening complications of bowel resection, leading to delayed recovery

and far higher total health-care costs. The delays shown were in diagnosis, not in waiting for treatment. Additionally, access to mesh in lower-income settings could be improved, and day-case surgery for simple hernias promoted. This allowed us to identify actionable targets: improving community worker knowledge and referral pathways is necessary in low-income settings, combined with a global quality improvement programme to boost mesh availability. The lack of mesh in lower-income settings exposes inequities in affordable medical devices and can be solved with industry collaboration and a global training programme. Day-case rates for eligible patients were still far from the targets desired in most income settings, so setting up dedicated pathways will reduce overall costs, boost capacity, and improve system resilience.

Implications of all the available evidence

Without correcting downwards spirals of reliance on emergency care, crisis management becomes the norm across a wide range of conditions, as is currently being seen globally. Strengthening elective pathways will also relieve the burden on emergency pathways by avoiding emergency presentation for conditions that can be fixed at an early stage with simple treatments. It can promote earlier presentation for emergency surgery, such as surgery for appendicitis, preventing the need for complex emergency care too. Policy makers can use this study as a proxy for other elective conditions, which are likely to suffer from the same weaknesses but are too numerous to study individually. Opportunistic detection and optimisation of non-surgical disorders (eg, raised blood pressure, raised blood sugar) would integrate surgery into the wider system of primary and preventive health care rather than leave it as a standalone entity. This system strengthening approach will also be realised in the management of multimorbidity. All these factors, underpinned by timely elective care, will reduce the health burden on society and health-care services.

Inguinal hernias are a common, global surgical condition and suitable for such a task.⁸ The World Bank Essential Surgery Package identifies inguinal hernia as a condition that leads to substantial global burden, treatment represents a substantial surgical need, and its repair is cost-effective.⁹ Inguinal hernia repairs are offered by health systems worldwide, and there are accepted guidelines for treatment.^{10,11} Not offering timely repair can impact patients’ ability to do daily activities, limits their capacity to do heavy work leading to societal cost, and can increase the rate of emergency surgery.^{12–14}

The study aimed to take a whole-system approach to evaluate access to and quality of elective health care around the world, using inguinal hernia as a tracer condition. We set out to collect data at both patient level and health system level, bridging the gap between the two. We designed the analysis to identify actionable

targets directly relevant to inguinal hernia repair, but with features that might also strengthen wider elective care pathways. We accepted from the very start of planning that a single outcome measure would not be appropriate for whole-system assessment, so we set out to design the study in a way that reported evenly across wider health systems.

Methods

Study design and participants

We conducted an international, multicentre, prospective cohort study of patients undergoing inguinal hernia surgery (HIPPO). We collected only routine, anonymised data, without making any change to existing clinical care pathways within each hospital. The study was prospectively registered in ClinicalTrials.gov (NCT05748886) and the full protocol is available online. The local principal

For the protocol see <https://www.globalsurgeryunit.org/clinical-trials-holding-page/hippo/>

investigators were responsible for seeking approvals according to regulations in place. In some participating hospitals, informed patient consent was taken, whereas in other hospitals it was waived by the local research ethics committees.

Any hospital performing inguinal hernia repair was eligible to take part. Eligible hospitals were identified by national leads and invited to take part. Participating hospitals identified and included consecutive patients of any age undergoing primary inguinal hernia repair as the main procedure during a 4-week inclusion window, between Jan 30 and May 21, 2023, with a follow-up window of 30 days after the date of surgery. Both elective and emergency indications were included. All surgical approaches were included apart from open surgeries performed via midline incision as these represent more complex operations, often associated with other procedures.

Measurement set

The Study Management Group identified that no single outcome would have face validity in accurately evaluating performance of whole surgical systems, based on literature review and discussions. To have a more comprehensive evaluation of elective surgery system performance, we needed to consider multiple different measures. Therefore, we defined a measurement set through a multi-stage consensus process within the Study Management Group (appendix 1 p 36). The measures we used were designed to align with the attributes of WHO's Health System Building Blocks: (1) access, including measures of access and coverage, and (2) quality, including measures of quality and safety.¹⁵

The measurement set was organised into six key performance measures and seven additional descriptive measures (appendix 1 p 38). We used staged inclusion and exclusion criteria to determine the sample in which each measure would be assessed. To address access we measured: (1) emergency rate: the proportion of patients undergoing emergency surgery (measurable in all included patients); (2) bowel resection rate: the proportion of patients in whom bowel resection occurred (all patients); and (3) waiting time: time between the date of surgery and start of symptoms for symptomatic patients and the date of diagnosis for asymptomatic patients (elective surgery only). To address quality we measured: (1) mesh use rate: the proportion of patients in whom a mesh was used out of all patients in whom a mesh would be indicated according to international guidelines¹¹ (adults undergoing elective surgery only); (2) day-case rate: the proportion of patients who were discharged on the same day out of all patients for whom day-case surgery would be recommended (adults younger than 90 years, American Society of Anesthesiologists Physical Status Classification System [ASA] grade I–II, undergoing elective surgery); and (3) postoperative complications: defined by Clavien-Dindo classification

and including all grades I–V (all patients). To better assess safety, we created a model for complications taking into account the features of the patient pathway. Detailed rationale for the key performance measures is described in appendix 1 (p 39).

Seven additional descriptive measures were also defined. To further describe access, we used hospital-level measures (standardised patient pathways, waiting list management, availability of day-case surgical unit, and financing methods). To describe quality, we used intraoperative measures including primary operator (a self-identified variable of senior surgeon, trainee surgeon, and non-surgeon medical practitioner), anaesthesia used (local, regional block, spinal, and general), and surgical approach (open, minimally invasive, and minimally invasive converted to open).

Data management

Data were collected and stored online through a secure server running the Research Electronic Data Capture (REDCap) web application.¹⁶ The service was managed by the Global Surgery REDCap system hosted at the University of Birmingham, UK, and its security was governed by the policies of the University of Birmingham. Collaborators were given REDCap project server login details, allowing secure data entry and storage. For more information on data collection see appendix 1 (p 37).




Figure 1: Flow of patients mapped to key performance measures

This figure shows the included and excluded patients when staged inclusion and exclusion criteria were applied, to evaluate the key performance measures in the correct sample. The rationale for this is explained in appendix 1 (p 39). ASA=American Society of Anesthesiologists Physical Status Classification System grade.

	HICs (n=9808)	UMICs (n=3366)	LMICs (n=3948)	LICs (n=936)	Total (n=18 058)
Age, years					
Median	63.0	53.0	42.0	44.0	57.0
IQR	51.0–73.0	32.0–66.0	11.5–58.5	20.0–60.0	39.0–69.0
Age groups					
Infants (<1 year)	124 (1.3%)	243 (7.2%)	389 (9.9%)	78 (8.4%)	834 (4.6%)
Children (1 year to <16 years)	245 (2.5%)	325 (9.7%)	640 (16.2%)	128 (13.7%)	1338 (7.4%)
Adults (≥16 years)	9432 (96.2%)	2796 (83.1%)	2918 (73.9%)	725 (77.9%)	15 871 (88.0%)
Missing	7	2	1	5	15
Sex					
Female	1008 (10.3%)	386 (11.5%)	361 (9.1%)	84 (9.0%)	1839 (10.2%)
Male	8792 (89.7%)	2980 (88.5%)	3586 (90.9%)	847 (91.0%)	16 205 (89.8%)
Missing	8	0	1	5	14
ASA grade					
I–II	7865 (80.3%)	3070 (91.2%)	3793 (96.1%)	895 (96.1%)	15 623 (86.6%)
III–V	1868 (19.1%)	290 (8.6%)	144 (3.6%)	29 (3.1%)	2331 (12.9%)
Not recorded	67 (0.7%)	6 (0.2%)	10 (0.3%)	7 (0.8%)	90 (0.5%)
Missing	8	0	1	5	14
Comorbidities					
None	7267 (74.1%)	2670 (79.4%)	3417 (86.6%)	802 (86.1%)	14 156 (78.5%)
One	1840 (18.8%)	535 (15.9%)	436 (11.1%)	119 (12.8%)	2930 (16.2%)
Two	515 (5.3%)	121 (3.6%)	84 (2.1%)	10 (1.1%)	730 (4.1%)
Three or more	172 (1.8%)	36 (1.1%)	7 (0.2%)	0 (0.0%)	215 (1.2%)
Missing	12	3	4	5	24
Symptoms					
Asymptomatic	1312 (13.4%)	520 (15.4%)	974 (24.7%)	117 (12.6%)	2923 (16.2%)
Symptomatic	8486 (86.6%)	2846 (84.6%)	2972 (75.3%)	815 (87.4%)	15 119 (83.8%)
Missing	10	0	2	4	16
Hernia size					
Limited to inguinal region	8209 (83.8%)	2498 (74.2%)	2351 (59.6%)	636 (68.3%)	13 694 (75.9%)
Limited to scrotum	1490 (15.2%)	826 (24.5%)	1535 (38.9%)	283 (30.4%)	4134 (22.9%)
Extending to mid-thigh or beyond	102 (1.0%)	42 (1.2%)	61 (1.5%)	12 (1.3%)	217 (1.2%)
Missing	7	0	1	5	13
Contamination					
Clean	9640 (98.4%)	3081 (91.6%)	3731 (94.6%)	898 (96.4%)	17 350 (96.2%)
Clean-contaminated	138 (1.4%)	271 (8.1%)	185 (4.7%)	30 (3.2%)	624 (3.5%)
Contaminated	15 (0.2%)	10 (0.3%)	25 (0.6%)	0 (0.0%)	50 (0.3%)
Dirty	5 (0.1%)	3 (0.1%)	5 (0.1%)	4 (0.4%)	17 (0.1%)
Missing	10	1	2	4	17
Hernia defect size					
<1.5 cm	2132 (21.8%)	703 (20.9%)	988 (25.0%)	238 (25.5%)	4061 (22.5%)
1.5 cm to 3 cm	3872 (39.5%)	1223 (36.3%)	1444 (36.6%)	378 (40.6%)	6917 (38.3%)
>3 cm	1975 (20.2%)	998 (29.6%)	1154 (29.2%)	235 (25.2%)	4362 (24.2%)
Not known	1816 (18.5%)	442 (13.1%)	360 (9.1%)	81 (8.7%)	2699 (15.0%)
Missing	13	0	2	4	19

Data are n (%) unless stated otherwise. Denominators for each group are presented in the top row; missing data are excluded from the denominators when calculating the percentages. Patient characteristics and intraoperative details from all patients included in the analysis are shown. HIC=high-income country. UMIC=upper-middle-income country. LMIC=lower-middle-income country. LIC=low-income country. ASA=American Society of Anesthesiologists Physical Status Classification System.

Table: Preoperative and intraoperative characteristics across income groups

Data validation

We have previously validated our data collection methodology in terms of case ascertainment and data accuracy.^{17,18} Each hospital lead was responsible for data

accuracy and data completeness collected and uploaded from their teams. The data were checked centrally and when there were missing data or invalid data, the hospital lead was contacted to complete and correct the data




Figure 2: Key performance measures to evaluate access and quality

This figure shows the variation of access, coverage, quality, and safety across the income groups. Categorical variables are presented with adjusted rates for hospital and country, and waiting times are presented as geometric means, also adjusted for hospital and country. Missing data are excluded from this figure and are reported in appendix 1 (p 46). HIC=high-income country. UMIC=upper-middle-income country. LMIC=lower-middle-income country. LIC=low-income country.

entered before the data lock. Following this, participating hospitals with data completeness below 95% were excluded.

Sample size

To ensure global generalisability of the results and to justify the resources put into the study, we estimated a minimum number of 300 hospitals contributing patient-level data from 70 countries, based on previous cohort studies (ie, GlobalSurg, COVIDSurg).^{17,18} Assuming an

average of 30 patients per hospital, we predicted a minimum sample size of 10000 patients. A sample of 10000 equates to margins of error between 0.2% and 0.85% depending on the binary outcome and a width of 0.39 for the continuous outcome (see appendix 1 p 40 for full details).

Statistical analysis

The data were mapped to country income groups, which were defined according to the World Bank into four




Figure 3: Waiting times components to elective surgery across income groups

This figure shows the distribution of waiting times across the patient pathway; diagnosis and decision for surgery represent key points in access to elective surgery. Data are presented as geometric mean and 95% CI. Missing data are excluded from this figure and are reported in appendix 1 (p 46). HIC=high-income country. UMIC=upper-middle-income country. LMIC=lower-middle-income country. LIC=low-income country.

categories: low-income countries (LICs), lower-middle-income countries (LMICs), upper-middle-income countries (UMICs), and high-income countries (HICs), as the importance of country income level in relation to health-care access and quality has been widely recognised.⁶

Continuous hospital, patient, and intraoperative characteristics were presented as mean and SD if normally distributed and median and IQR if not normally distributed. Categorical variables were described using frequencies and percentages. Rates of key performance measures by income group were presented with adjusted rates from three-level multilevel logistic regression models (hospital nested within country) and 95% CIs. Log-transformed waiting times between income groups were summarised using geometric means and 95% CIs from three-level multilevel linear regression models with similar structure as above. A three-level multilevel logistic regression model was used to explore the association of factors describing the patient pathway with complications. Clinically plausible factors agreed by the Study Management Group were included as covariates: income groups, age group, ASA groups, urgency of surgery, anaesthesia type, contamination, bowel resection, use of mesh, and day-case surgery. Hospitals nested within country were included as random effects. For the above analyses, appropriate model fit diagnostics were checked to confirm that validity and model assumptions hold for the data. All statistical analyses were performed using R (version 4.0.2). A p value of less than 0.05 was considered statistically significant.

Role of the funding source

The funders of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report.

Results

Data from 18465 patients were collected across 642 hospitals from 83 countries that participated in this study. From those, 407 patients were excluded and data from 18058 patients, across 640 hospitals, were analysed. Exclusion reasons, proportions within each income group, and distribution across countries are shown in appendix 1 (pp 48–49). Patient selection, including the patient group in which each key performance measure was assessed, is presented in figure 1.

Overall, most patients were male (16205 [89.8%] of 18044), with a median age of 57.0 years (IQR 39.0–69.0; table). There were more patients with symptoms related to their inguinal hernia (15119 [83.8%] of 18042) than patients without symptoms. Most hernias were limited in size to the inguinal region (13694 [75.9%] of 18045), with the remainder limited to the scrotum. Most operations were classified as clean (17350 [96.2%] of 18041). Other preoperative and intraoperative characteristics are described in appendix 1 (pp 41–43).

Most included patients underwent surgery in tertiary-level hospitals (11645 [65.1%] of 17887; appendix 1 p 44). From the included hospitals, most were publicly funded (500 [80.1%] of 624; appendix 1 p 45). In addition, most were able to provide emergency surgery over 24 h (565 [90.5%] of 624).

The overall emergency surgery rate was 7.1% (1287 of 18041 patients), which increased from HICs to LICs, as shown by figure 2. This was accompanied by an increase in bowel resection rate, from adjusted rates of 1.2% (105 of 9798 patients) in HICs to 4.2% (36 of 932s) in LICs.

The geometric mean of overall waiting time for elective surgery was 8.0 months (95% CI 7.8–8.1). Overall waiting times for elective patients were slightly longer in HICs (geometric mean 8.2 months) and shortest in LMICs (geometric mean 5.9 months). When looking in more detail at the components of the overall waiting time, important differences appeared, particularly after contact with the health system was established. After diagnosis, the waiting time to surgery was 3.6 months in HICs but only 1.2 months in LICs (figure 3)

Overall mesh use rate was 94.8% (13995 of 14768 patients), which decreased from HICs (adjusted rate 97.6%; 8842 of 8916) to LICs (adjusted rate 61.0%; 457 of 635).

Day-case rates were low and variable across income groups. The highest adoption of day-case surgery occurred in HICs and in LICs, with adjusted rates of 50.0% (4464 of 7182) and 44.5% (368 of 605), respectively.

Of the 18058 included patients, 14580 (80.7%) had a follow-up appointment within 30 days of the surgery. A total of 18018 patients had follow-up data available, as data were also available for some patients from emergency department or unplanned clinic records. The overall postoperative complications rate at 30 days was 13.4% (2415 of 18018 cases). Of these 2415 cases, 1808 (74.9%) were Clavien-Dindo grade I–II, and only 16 (0.7%) were Clavien-Dindo grade V (death; appendix 1 p 46). An ASA grade of III–V, clean-contaminated, contaminated, or dirty surgery, emergency surgery, and bowel resection were all associated with increased risk of postoperative complications (figure 4).

Standardised patient pathways, waiting list management, and availability of day-case surgical units were variable across the different income groups (figure 5). All were consistently highest in HICs (appendix 1 p 47).

Insurance provided by the government was the payment type used by most patients in all income groups (14297 [79.9%] of 17887). From the other payment methods, out-of-pocket payments were highest in LMICs (1081 [27.4%] of 3948) and LICs (189 [20.8%] of 908).

Senior surgeons were present in 12752 (70.7%) of 18042 cases overall, which was similar in the different income groups (figure 5). Anaesthesia type showed wide variation. General anaesthesia was more used in HICs and UMICs (5867 [59.9%] of 9798 and 1920 [57.0%] of 3366, respectively) as opposed to spinal anaesthesia, which was more commonly used in LMICs and LICs (1890 [47.9%] of 3944 and 548 [58.8%] of 932, respectively). More details on other anaesthetic practices are described in figure 5 and appendix 1 p 47.





Figure 4: Multilevel logistic regression model to test factors associated with complications at 30 days

This model shows the factors associated with complications, reflecting a multilevel model with hospital and country as the two levels that were adjusted for. All patients were included in the model, except those with missing data for any of the variables ($n=40$). OR=odds ratio. HIC=high-income country. UMIC=upper-middle-income country. LMIC=lower-middle-income country. LIC=low-income country. ASA=American Society of Anesthesiologists Physical Status Classification System. *Includes infants (<1 year).

Overall, open surgery was used in most patients (14117 [78.2%] of 18041), with a decrease in minimally invasive surgery from higher to lower incomes (HICs 2723 [27.8%] of 9798, UMICs 759 [22.6%] of 3365, LMICs 433 [11.0%] of 3946, LICs 9 [1.0%] of 932). Laparoscopic surgery represented 3769 (96.0%) of all 3924 cases of minimally invasive surgery, with robotic approach being used in the remaining 155 (4.0%).

Discussion

This study shows that inguinal hernias are mostly a disease of working-age patients around the world, and when small, are treatable with simple, day-case surgery. If neglected, the more complex emergency surgery, which might require bowel resection, leads to slow recovery and far higher total health-care costs.¹⁹ We also



found that most of the waiting time was before the diagnosis was made, especially in low-income settings, rather than waiting for treatment after seeing a surgeon. Using inguinal hernia as a tracer condition for elective care, this study showed multiple weaknesses in access and quality in current health-care systems, with a particular disadvantage in lower-income settings. As a result, there was higher emergency demand, which further reduced elective capacity and might create downward spirals. Without addressing these spirals, emergency care and crisis management become the norm with potentially worse outcomes for patients.^{6,12} Policy makers can use this study as a proxy for other elective conditions, which are likely to suffer from the same weaknesses.² If weak access and quality persist over several electively treatable conditions, both surgical and non-surgical, multimorbidity can also become established, which makes future elective care harder and emergency care even more complicated.²⁰

We showed that inguinal hernia is a low-risk procedure that can be provided by a range of practitioners and with a range of anaesthetic types, while maintaining safety. Most complications were minor. However, emergency surgery was associated with more complications. We did not capture the most complex emergency surgery performed though larger midline incisions, as these operations require different data capture.¹³ It is thus likely that this study underestimates the true burden of emergency surgery.¹² This study also shows a clear global imbalance in access to mesh repair, which probably reflects poor access to simple medical devices in lower-income settings.²¹ Mesh is well proven to reduce long-term hernia recurrence,^{10,11} is simple to place, and is low-cost and should be scaled.

The following actionable targets supported by our findings are relevant to front-line teams and policy makers. In some lower-income settings, improving referral systems and increasing the use of mesh repair will improve access and quality. Educating communities and community health workers around the symptoms of hernias and establishing referral pathways might facilitate timely evaluation by a surgeon. A global quality improvement programme in mesh placement for hernias is justified. This would involve strengthening industrial supply chains, making mesh affordable to patients, and increasing training to ensure safe and effective placement. In all income settings, boosting day-case surgery will improve capacity for simple, cost-effective surgery while also making systems resilient against

Figure 5: Additional descriptive measures panel
 This figure shows the variation of additional descriptive data across income groups. The error bars represent the 95% CIs for the rates. Missing data are excluded from this figure and are reported in appendix 1 (p 47). HIC=high-income country. UMIC=upper-middle-income country. LMIC=lower-middle-income country. LIC=low-income country. MIS=minimally invasive surgery.

external threats.²² Mesh placement and use of available anaesthetic resources would fit with the latest guidelines from the HerniaSurge Collaborative and for future global burden of disease studies with a focus on inguinal hernias, mesh and anaesthetic use could be taken into account during the modelling exercises.¹⁰

These targets might also strengthen other areas of simple, common elective health care, as identified through the *Lancet* Commission on Surgery and the *Lancet* Commission on Diagnostics.^{2,21} The assessment and referral pathway could be combined with a set of simple assessments to detect other common diseases that would benefit from timely diagnosis and management to reduce multimorbidity. At the point of first appointment, blood pressure checks, blood sugar checks, and HIV checks are all deliverable and effective, making surgery part of an embedded pathway rather than a standalone speciality.²³

The strengths of this study include its prospective, dedicated nature, which creates a unique dataset, allowing us to report a measurement set that comprehensively evaluates the whole system, rather than focusing on a single outcome measure. By doing so, we provide more holistic evidence relevant to all stakeholders, from front-line teams to policy makers. The design also allows us to identify several actionable targets and thus solutions to global problems around elective health care.

This study has important limitations. Using inguinal hernia as a tracer condition is a limitation in itself, considering that the randomised controlled trials conducted so far have shown that the individual risk for emergency surgery seems to be low.²⁴ We did not capture patients who did not undergo surgery, meaning the societal burden of untreated hernia could be far higher than we can estimate.²⁵ As discussed, we did not capture hernia repair through a midline incision, which is often used for the most complex emergency situations, underestimating the true burden of emergency surgery. There might be under-representation from some countries where minimally invasive surgery rates are currently very high (eg, Denmark, Norway, and Sweden), skewing results for those parts of the world.²⁶ Interpretation of the findings in the context of secondary-level and primary-level hospitals needs to be carefully made as most participating hospitals were tertiary level. We did not explore variation of the approval processes conducted nor the detailed differences in primary operator across countries, and we recognise there might be hidden bias related to these differences. Some data captured at hospital level (eg, payment mechanism) might not fully describe each individual patient's journey. We did not contact patients routinely at 30 days after surgery, which can be impossible in some settings, but patients with major events would be likely to re-present in their hospitals within 30 days and be captured by this study. There is a risk of over-representation from some countries, which is inherent to a global cohort study such as HIPPO. There should be some caution

around conclusions on the whole system as the results might not be completely representative of every country included in each income group.

This study has recognised topics for future research. Inguinal hernia is one of many potential tracer conditions, for both elective care and surgery. Future researchers can select their own tracer conditions and use these key performance measures to provide a whole-system assessment. Further research will be needed to evaluate waiting times in emergency surgery as it can highlight hidden problems. Qualitative studies would be helpful to understand the variation in asymptomatic patients having an operation. Finally, from a methodology perspective, it would be useful to clarify the differences in study approval processes in different countries to see if, in limited settings, they are subject to any bias.

This study has identified areas for intervention by policy makers. Financing these pathways will be a challenge, especially if elective care needs to compete with emergency care. The additional descriptive measures in this study show the importance of affordable health insurance (whether government-funded or individually funded), which should be increased for common conditions around the world. Relying on out-of-pocket payments is likely to limit elective care to those with savings, or plunge families into catastrophic expenditure.^{27,28}

Contributors

MP, AOA, AA, AEA, JAC, DC, RC, ACD, ME, AG, DG, JG, PH, EH, AI, IJ, SKK, OK, IL, SL, VL, EL, JM, AMB, DM, DN, FN, OO, SZYO, RO, FP, ARM, MSA, JFFS, MS, ST, and AB were involved in study design, coordination of the study, manuscript concept, and editing. MP, RA, DB, AC, JGGN, RRG, BK, SRK, SL, OO, KSYO, RO, and CV were involved in data chasing and accuracy of the data. BK, OO, MP, and AB conducted the data analysis. OO, MP, and AB accessed and verified the data. Dissemination Committee co-authors were involved in the dissemination of the study, coordination at a national level, and data collection. Hospital-lead co-authors were involved in coordination of the study at a hospital level and data collection. Collaborator co-authors were involved in data collection. All authors had full access to all the data in the study and had final responsibility for the decision to submit for publication.

Equitable partnership declaration

The authors of this paper have submitted an equitable partnership declaration (appendix 2). This statement allows researchers to describe how their work engages with researchers, communities, and environments in the countries of study. This statement is part of *The Lancet Global Health's* broader goal to decolonise global health.

See Online for appendix 2

Declaration of interests

MP reports a project research grant from the Portuguese Hernia and Abdominal Wall Society (Sociedade Portuguesa de Hernia e Parede Abdominal). AB reports a Global Health Research Unit grant from the National Institute for Health and Care Research (NIHR). All other authors declare no competing interests.

Data sharing

Anonymised data are available upon request from the corresponding authors and successful completion of a data sharing agreement through an Application Programming Interface linked to the REDCap data server hosted at the University of Birmingham, Birmingham, UK.

Acknowledgments

This study was supported by a grant from the NIHR Global Health Research Unit on Global Surgery (NIHR133364) and a project research

grant from the Portuguese Hernia and Abdominal Wall Society (Sociedade Portuguesa de Hernia e Parede Abdominal). The views expressed are those of the authors and not necessarily those of the National Health Service, the NIHR, or the UK Department of Health and Social Care.

References

- 1 Kruk ME, Gage AD, Arsenault C, et al. High-quality health systems in the Sustainable Development Goals era: time for a revolution. *Lancet Glob Health* 2018; **6**: e1196–252.
- 2 Fleming KA, Horton S, Wilson ML, et al. The *Lancet* Commission on diagnostics: transforming access to diagnostics. *Lancet* 2021; **398**: 1997–2050.
- 3 COVIDSurg Collaborative. Elective surgery cancellations due to the COVID-19 pandemic: global predictive modelling to inform surgical recovery plans. *Br J Surg* 2020; **107**: 1440–49.
- 4 Khan Y, Verhaeghe N, Devleeschauwer B, et al. Impact of the COVID-19 pandemic on delayed care of cardiovascular diseases in Europe: a systematic review. *Lancet* 2023; **402** (suppl 1): S61.
- 5 Columbus AB, Morris MA, Lilley EJ, et al. Critical differences between elective and emergency surgery: identifying domains for quality improvement in emergency general surgery. *Surgery* 2018; **163**: 832–38.
- 6 GlobalSurg Collaborative. Mortality of emergency abdominal surgery in high-, middle- and low-income countries. *Br J Surg* 2016; **103**: 971–88.
- 7 Wolfe I, Forman J, Cecil E, et al. Effect of the Children and Young People's Health Partnership model of paediatric integrated care on health service use and child health outcomes: a pragmatic two-arm cluster randomised controlled trial. *Lancet Child Adolesc Health* 2023; **7**: 830–43.
- 8 Abebe MS, Tareke AA, Alem A, Debebe W, Beyene A. Worldwide magnitude of inguinal hernia: systematic review and meta-analysis of population-based studies. *SAGE Open Med* 2022; **10**: 20503121221139150.
- 9 Mock CN, Donkor P, Gawande A, Jamison DT, Kruk ME, Debas HT. Essential surgery: key messages of this volume. In: Debas HT, Donkor P, Gawande A, Jamison DT, Kruk ME, Mock CN, eds. *Essential surgery: disease control priorities*, 3rd edn (vol 1). Washington, DC: The World Bank, 2015.
- 10 Stabilini C, van Veenendaal N, Aasvang E, et al. Update of the international HerniaSurge guidelines for groin hernia management. *BJS Open* 2023; **7**: zrad080.
- 11 HerniaSurge Group. International guidelines for groin hernia management. *Hernia* 2018; **22**: 1–165.
- 12 Samuel JC, Tyson AF, Mabedi C, et al. Development of a ratio of emergent to total hernia repairs as a surgical capacity metric. *Int J Surg* 2014; **12**: 906–11.
- 13 Köckerling F, Heine T, Adolf D, et al. Trends in emergent groin hernia repair—an analysis from the Herniated Registry. *Front Surg* 2021; **8**: 655755.
- 14 Hwang MJ, Bhangu A, Webster CE, Bowley DM, Gannon MX, Karandikar SS. Unintended consequences of policy change to watchful waiting for asymptomatic inguinal hernias. *Ann R Coll Surg Engl* 2014; **96**: 343–47.
- 15 Borghi J, Brown GW. Taking Systems Thinking to the global level: using the WHO building blocks to describe and appraise the global health system in relation to COVID-19. *Glob Policy* 2022; **13**: 193–207.
- 16 Harris PA, Taylor R, Minor BL, et al. The REDCap consortium: building an international community of software platform partners. *J Biomed Inform* 2019; **95**: 103208.
- 17 GlobalSurg Collaborative and National Institute for Health Research Global Health Research Unit on Global Surgery. Global variation in postoperative mortality and complications after cancer surgery: a multicentre, prospective cohort study in 82 countries. *Lancet* 2021; **397**: 387–97.
- 18 COVIDSurg Collaborative. Timing of surgery following SARS-CoV-2 infection: an international prospective cohort study. *Anaesthesia* 2021; **76**: 748–58.
- 19 Margenthaler JA, Longo WE, Virgo KS, et al. Risk factors for adverse outcomes following surgery for small bowel obstruction. *Ann Surg* 2006; **243**: 456–64.
- 20 van Ginneken E, Reed S, Siciliani L, et al. Addressing backlogs and managing waiting lists during and beyond the COVID-19 pandemic. 2022. <https://eurohealthobservatory.who.int/publications/i/addressing-backlogs-and-managing-waiting-lists-during-and-beyond-the-covid-19-pandemic> (accessed Nov 6, 2023).
- 21 Meara JG, Leather AJ, Hagander L, et al. Global Surgery 2030: evidence and solutions for achieving health, welfare, and economic development. *Lancet* 2015; **386**: 569–624.
- 22 NIHR Global Health Unit on Global Surgery. Elective surgery system strengthening: development, measurement, and validation of the surgical preparedness index across 1632 hospitals in 119 countries. *Lancet* 2022; **400**: 1607–17.
- 23 Kivuyo S, Birungi J, Okebe J, et al. Integrated management of HIV, diabetes, and hypertension in sub-Saharan Africa (INTE-AFRICA): a pragmatic cluster-randomised, controlled trial. *Lancet* 2023; **402**: 1241–50.
- 24 Van den Dop LM, Van Egmond S, Heijnen J, et al. Twelve-year outcomes of watchful waiting versus surgery of mildly symptomatic or asymptomatic inguinal hernia in men aged 50 years and older: a randomised controlled trial. *EClinicalMedicine* 2023; **64**: 102207.
- 25 Ma Q, Jing W, Liu X, Liu J, Liu M, Chen J. The global, regional, and national burden and its trends of inguinal, femoral, and abdominal hernia from 1990 to 2019: findings from the 2019 Global Burden of Disease Study—a cross-sectional study. *Int J Surg* 2023; **109**: 333–42.
- 26 OECD. Surgical operations and procedures statistics. 2023. https://ec.europa.eu/eurostat/statistics-explained/index.php?title=Surgical_operations_and_procedures_statistics#Number_of_surgical_operations_and_procedures (accessed Nov 20, 2023).
- 27 Shrimme MG, Dare AJ, Alkire BC, O'Neill K, Meara JG. Catastrophic expenditure to pay for surgery worldwide: a modelling study. *Lancet Glob Health* 2015; **3** (suppl 2): S38–44.
- 28 CROCODILE study group. Catastrophic expenditure and treatment attrition in patients seeking comprehensive colorectal cancer treatment in India: a prospective multicentre study. *Lancet Reg Health Southeast Asia* 2022; **6**: 100058.

THE LANCET

Global Health

Supplementary appendix 1

This appendix formed part of the original submission and has been peer reviewed. We post it as supplied by the authors.

Supplement to: NIHR Global Health Research Unit on Global Surgery. Access to and quality of elective care: a prospective cohort study using hernia surgery as a tracer condition in 83 countries. *Lancet Glob Health* 2024; published online May 23. [https://doi.org/10.1016/S2214-109X\(24\)00142-6](https://doi.org/10.1016/S2214-109X(24)00142-6).

Supplement

Access to and quality of elective care: a prospective cohort study using hernia surgery as a tracer condition in 83 countries

NIHR Global Health Research Unit on Global Surgery

A full list of collaborating authors is shown in Appendix 1

Corresponding authors:

Dr Maria Picciochi, NIHR Global Health Research Unit on Global Surgery, Institute of Applied Health Research, University of Birmingham, Birmingham B15 2TH, UK

m.picciochi@bham.ac.uk

and

Professor Aneel Bhangu, NIHR Global Health Research Unit on Global Surgery, Institute of Applied Health Research, University of Birmingham, Birmingham B15 2TH, UK

a.a.bhangu@bham.ac.uk

Acknowledgments: This study was supported by NIHR Global Health Research Unit Grant (NIHR133364) and a project research grant from Portuguese Hernia and Abdominal Wall Society (Sociedade Portuguesa de Hernia e Parede Abdominal). The funders had no role in study design or writing of this report. The views expressed are those of the authors and not necessarily those of the National Health Service, the NIHR or the UK Department of Health and Social Care.

Appendices

Section 1 – List of authors

Pages 1-34

Section 2 – Supplementary tables and materials

Pages 35-45

Section 1: List of authors

Dissemination Committee (listed by country):

Albania: I Dajti; Algeria: Z Djama; Argentina: M Lucchini, RM Palacios Huatuco; Australia: K Atherton, AC Dawson, E Lun; Austria: F Aigner; Belgium: F Berrevoet; Benin: I Lawani, S Lawani, C Bokossa; Bosnia and Herzegovina: S Delibegovic; Bulgaria: M Slavchev; Burkina Faso: AF Sanon, A Sanou; Burundi: JB Gusa, JC Mbonicura; Cameroon: A Bang, O Gabom, C Nwegbu; Canada: A Brar, J Martin; Chile: MM Modolo, M Olivos; Colombia: JA Calvache; Croatia: J Mihanovic; Cyprus: N Gouvas, A Yiallourou; Czech Republic: B East; Dominican Republic: S Batista, R Rivas; Ecuador: EP Lincango; Egypt: S Emile; Ethiopia: AB Aregawi; France: AP Arnaud; Gabon: N Boumas; Georgia: Z Demetrashvili; Germany: H Lederhuber, MW Löffler; Ghana: AE Agbeko, NB Sam, S Tabiri, F Agyei, FE Gyamfi, S Mohammed; Greece: I Katsaros, G Tsoulfas; India: L Bains, J Dhiman, D Ghosh, PD Haque, A Suroy; Ireland: S Ramjit; Israel: G Marom; Italy: F Pata, G Gallo; Jordan: F Ayasra; Kazakhstan: I R Fakhradiyev; Kenya: IHS Hamdun; Kyrgyzstan: A Iqbal; Liberia: E Mbanzabugabo; Libya: M Elhadi; Lithuania: A Gulla; Madagascar: L Samison; Malawi: M Nyirenda, R Nyirenda; Malaysia: AC Roslani; Mali: B Bengaly; Malta: J Psaila; Mexico: L Martinez, A Ramos-De la Medina; Namibia: PR Nashidengo; New Zealand: M McGuinness, D Wright; Niger: A Ousseini; Nigeria: A Adisa, AO Ademuyiwa; North Macedonia: T Risteski; Oman: Z Al Balushi, B Dawud, A AlSharqi, F Ali; Pakistan: AU Qureshi; Palestine: H Abu-Arish; Paraguay: H Gomez-Fernandez; Philippines: JM Faylona, MD Sacdalan; Poland: W Krawczyk; Portugal: JG Goncalves-Nobre, M Sampaio-Alves, I Santos; Romania: I Negoii; Russian Federation: A Butyrskii; Rwanda: JC Allen, F Ntirenganya; Sierra Leone: I Fortune; Slovenia: J Kosir; South Africa: N Parker, K Chu; Spain: A Minaya Bravo; Sri Lanka: D Wickramasinghe, U Jayarajah; Sudan: M Elmujtaba; Sweden: M Nikberg; Switzerland: E Gialamas; Syria: M Alshaar; Tanzania: M Nkoronko; Turkey: A Isik; Uganda: I Mubesi; United States: J Ng-Kamstra; Venezuela: O Bahsas-Zaky.

Writing group (Study Management Group): M Picciochi, AO Ademuyiwa, A Adisa, AE Agbeko, JA Calvache, D Chaudhry, R Crawford, AC Dawson, M Elhadi, A Ghaffar, D Ghosh, J Glasbey, PD Haque, E Harrison, A Isik, I Jakaityte, SK Kamarajah, O Kouli, I Lawani, S Lawani, V Ledda, E Li, J Martin, A Minaya Bravo, D Morton, D Nepogodiev, F Ntirenganya, O Omar, SZY Ooi, R Oppong, F Pata, A Ramos-De la Medina, M Sampaio-Alves, JFF Simoes, M Steinruecke, S Tabiri, A Bhangu.

Data handling and management: M Picciochi, R Acharya, D Badran, A Chaudhry, JG Goncalves-Nobre, RR Gujjuri, B Kadir (statistician), SR Knight, S Lawday, O Omar (senior statistician), KSY Ooi, R Ooi, C Varghese.

Hospital leads (listed by country and city):

Albania: E Agastra (Korca; Regional Hospital of Korca); I Dajti (Tirana; University hospital Koco Gliozheni).

Algeria: R Belouz (Algiers; CHU Isaad Hassani); Z Djama (Constantine; University Hospital Abdelhamid Ben Badis); A Mouffokes (Oran; EHU-1st November 1954).

Argentina: ME Muriel (Allende, Cordoba; Sanatorio Allende - Sede Cerro); RM Palacios Huatucu (Buenos Aires; Hospital Italiano de Buenos Aires); M Santillan (Buenos Aires; Hospital Universitario CEMIC); A Duro (Buenos Aires; Hospital municipal de vicente López); JI Valenzuela (City of Buenos Aires; Hospital Velez Sarsfield); DA Pantoja Pachajoa (Cordoba; Clinica Universitaria Reina Fabiola); G Romero reyna (Cordoba; Sanatorio Allende - Sede Nueva Cordoba); CM Florián Villa (San Francisco; Clinica Regional del Este).

Aruba: M Gosselink (Oranjestad; Dr. Horacio E Oduber Hospital).

Australia: YH Lam, A Nguyen (Adelaide; Northern Adelaide Local Health Network); JA Duffield (Adelaide; Royal Adelaide Hospital); A Frankel (Brisbane; Princess Alexandra Hospital); S Bowman (Brisbane; Queen Elizabeth 2 Jubilee Hospital); D Mitchell (Brisbane; Surgical Treatment and Rehabilitation Service); H Iswariah (Brisbane; The Prince Charles Hospital (TPCH)); S Abeykoon (Campbelltown; Campbelltown Hospital); S Gananadha (Canberra; Calvary Hospital); S Gananadha (Canberra; Canberra Hospital); S Salindera (Coffs Harbour NSW; Coffs Harbour Health Campus); S Stevens (Colac; Colac Area Health); AC Dawson (Gosford; Gosford Hospital); M Issa (Hamilton; Hamilton Base Hospital); E Wong (Melbourne; Angliss Hospital); L Bromley (Melbourne; Austin Hospital); E Wong (Melbourne; Box Hill Hospital); K Jaffry (Melbourne; Casey Hospital); M Bickford (Melbourne; Knox Private Hospital); E Wong (Melbourne; Maroondah Hospital); R Nataraja (Melbourne; Monash Childrens Hospital); R Hodgson (Melbourne; Northern Hospital); A Fox (Melbourne; St Vincent's Hospital); M Wichmann (Mount Gambier; Mount Gambier and Districts Health Service); A Davis (Mount Nasura; Armadale Health Service); S Zhang (Newcastle; Calvary Mater Newcastle); M Ishak (Newcastle; John Hunter Hospital); R Liang (Robina; Robina Hospital); P Tang (Sale; Sale Hospital / Central Gippsland Health Service); R Liang (Southport; Gold Coast University Hospital); M Park (Sydney; Royal North Shore Hospital); C Cornwell (Sydney; The Royal Prince Alfred Hospital); E Page-Taylor (Toowoomba; Toowoomba Hospital); A Hameed (Westmead; Westmead Hospital); R McGee (Wyong; Wyong Public Hospital).

Austria: I Königsrainer (Feldkirch; Landeskrankenhaus Feldkirch); F Aigner (Graz; Barmherzige Brüder Krankenhaus, Graz); S Mikalauskas (Graz; Medical University of Graz); L Havranek (Linz; Ordensklinikum Linz Elisabethinen); A Binder (Tulln; Universitätsklinikum Tulln).

Bangladesh: S Islam (Dhaka; Dhaka Medical College Hospital); MP Singh (Dinajpur; Lamb Hospital).

Benin: G Gbessi (Cotonou; Centre National Hospitalier et Universitaire Hubert Koutoukou Maga); H Aouagbe Behanzin (Cotonou; Hopital de Menontin); M Agbadebo (Dassa-Zoumè; Hôpital de Zone de Dassa-Zoumè); E Bara (Kandi; Hôpital de zone de Kandi); AB Yevide (Klouékanme; Hopital de Zone de Klouékanme); TK Hessou (Natitingou; Centre Hospitalier Départemental de l'Atacora); AM Hodonou (Parakou; Centre Hospitalier Universitaire Borgou Alibori); I Lawani (Porto Novo; Centre Hospitalier Universitaire et Departemental Oueme Plateau).

Bosnia and Herzegovina: Z Matkovic (Doboj; Genera Hospital 'Sveti aposto Luka' Doboj); N Lalovic (Foča; University Hospital Foča); J Miskovic (Mostar; SKB University Clinical Hospital Mostar); M Salibašić (Sarajevo; Clinical Center University of Sarajevo); A Cerovac (Tešanj; General Hospital Tešanj); A Tursunovic (Tuzla; University Clinical Center Tuzla).

Brazil: C Panis (Francisco Beltrão; Universidade Estadual do Oeste do Paraná).

Bulgaria: T Ivanov (Pleven; Heart and Brain - Pleven Hospital); M Karamanliev (Pleven; University Hospital Dr Georgi Stranski, Medical University - Pleven); R Donchev (Plovdiv; MHAT St. Karidad); D Hadzhiev (Plovdiv; UMHAT Sveti Georgi); T Yotsov (Ruse; University Hospital Medika); E Hristova (Sofia; Fifth City Hospital Sofia - 5th MBAL).

Burkina Faso: AF Sanon (Ouagadougou; Tengandogo University Hospital).

Burundi: JC Mbonicura (Bujumbura; Centre Hospitalo-Universitaire de Kamenge); N Diomede (Bujumbura; Kamenge Military Hospital); J Gusa (Bujumbura; Prince Regent Charles Hospital).

Cambodia: S Stock (Battambang; Handa Medical Centre).

Cameroon: O Ndizeye (Bamenda; Nkwen Baptist Hospital).

Canada: R Spence (Halifax; Queen Elizabeth II); S Lee (Port Moody; Eagle Ridge Hospital); S Lee (Vancouver; Royal Columbian Hospital).

Chile: MM Modolo (Santiago; Hospital Barros Luco Trudeau).

China: W Yang (Guangzhou; The First Affiliated Hospital of Jinan University).

Colombia: CJ Perez Rivera (Bogota; Fundacion Cardioinfantil-IC); S Sierra (Medellin; Clínica CES); DS Garcés Palacios (Popayan; Hospital Susana Lopez de Valencia); JA Calvache (Popayán; Hospital Universitario San José).

Croatia: J Mihanovic (Zadar; Zadar General Hospital); G Augustin (Zagreb; University Hospital Centre Zagreb).

Cyprus: A Yiallourou (Nicosia; Nicosia General Hospital).

Czech Republic: J Moravik (Decin; Krajská zdravotní as - Hospital Decin); A Al Kaddah (Hradec Kralove; Charles University Hospital); Z Musilová (Ivančice; Hospital Ivančice); M Schön (Novy Jicin; Hospital & Oncological Centre Novy Jicin); J Roman (Ostrava; University Hospital Ostrava); B East (Prague; Motol University Hospital).

Dominican Republic: R Rivas (Santo Domingo; CEDIMAT - Centro de Diagnóstico, Medicina Avanzada, Laboratorio y Telemedicina).

Egypt: G Abouelnagah (Alexandria; Alexandria Main University Hospital); D Attia (Alexandria; Alexandria Medical Research Institute); G Abouelnagah (Alexandria; Smouha University Hospital); A Maher (Assiut; Assiut University Children Hospital); A Kedwany (Assiut; Assiut University Hospital); S Abdelmohsen (Aswan; Aswan University Hospital, Aswan University); R Adel Diab (Cairo; Al Zahraa University Hospital); A Al-Mallah (Cairo; Al-Azhar University Hospitals); H Taher (Cairo; Cairo University Children's Hospitals (CUSPH & CUCH)); H Abozied (Cairo; EL-Hussein University Hospital, Al-Azhar University, Faculty Of Medicine); ASM Abdelrahman (Cairo; Giza International Hospital); M EIFiky, N Shehata (Cairo; Kasr Al Ainy Faculty of Medicine, Cairo University); B Fahmy (Giza; The Memorial Soaad Kafafi University Hospital); H Elghadban (Mansoura; Mansoura University Hospital); E Adel Mahmud Sultan (Menofia; Menofia University Hospital); M Omar (Qena; Qena University Hospital); EA Ahmed (Sohag; Sohag University Hospital); AM Elkhoully (Tanta; Tanta University Hospital); A Asla (Zagazig; Al Ahrar Zagazig Teaching Hospital).

Ethiopia: F Terefe (Addis Ababa; Yekatit 12 hospital medical college); A Yeshitila (Deberbirhan; Hakim Gizaw Hospital); M Taeme (Dessie; Dessie Referral Hospital); A Demessie (Gondar; Gondar University Comprehensive specialized hospital); AB Aregawi (Hawassa; Hawassa University

Comprehensive Specialized Hospital); N S.Bayleyegn (Jimma; Jimma University Medical Center); B Sime (Yirgalem; Yirgalem Hospital Medical College).

France: A Police (Eaubonne; Hôpital Simone Veil); A Castaldi (Nimes; Hôpital Carèmeau); E Reitano (Strasbourg; Nouvel Hopital Civil de Strasbourg).

Gabon: N Boumas (Libreville; Centre Hospitalier universitaire mère enfant Fondation Jeanne Ebori).

Georgia: Z Demetrashvili (Tbilisi; N.Kipshidze Central University Clinic).

Germany: C Kamphues (Berlin; Park-Klinik Weissensee); D Hackner (Erlangen; Universitätsklinikum Erlangen); U Ronellenfitsch (Halle; University Hospital Halle); J Rolinger (Moenchengladbach; Kliniken Maria Hilf); D Reim (Munich; Klinikum Rechts der Isar TUM School of Medicine); N Börner (Munich; Ludwig Maximilian University of Munich - Großhadern); J Goedeke (Munich; Ludwig Maximilian University of Munich - Innenstadt); AE Gut (München; Isarklinikum); M Janda (Rostock; University Hospital Rostock); J De Deken (Saarbruecken; Klinikum Saarbruecken); MW Löffler (Tuebingen; University Hospital Tuebingen).

Ghana: R Armah (Accra; Greater Accra Regional Hospital); A Bediako Bowan (Accra; Korle-Bu Teaching Hospital); E Kafui Ayodeji (Accra; Pentecost Hospital); EA Arkoh (Ankaful; Ankaful Leprosy General Hospital); FE Gyamfi (Berekum; Berekum Holy Family Hospital); A Davor (Bolgatanga; Upper East Regional Hospital); MT Morna (Cape-Coast; Cape Coast Teaching Hospital); N Agboadoh (Damongo; St. Anne's Hospital); EA Nacheleh (Ho; Ho Teaching Hospital); AE Agbeko (Kumasi; Komfo-Anokye Teaching Hospital); S Mensah (Kumasi; University Hospital, KNUST); P Taah-Amoako (Nsawkaw; Tain District Hospital); K Collins (Sunyani; Brong-Ahafo Regional Hospital); AS Seidu (Tamale; Tamale Teaching Hospital); BK Seshie (Tema; Tema General Hospital); G Ansong (Walewale; Walewale Government Hospital).

Greece: K Kambouri (Alexandroupolis; Alexandroupolis University General Hospital); A Kyriakidis (Amfissa; General Hospital of Amfissa); D Korkolis (Athens; Agios Savvas Anticancer Hospital); N Memos (Athens; Aretaieion Hospital); D Kelgiorgi (Athens; Athens Euroclinic); C Chouliaras (Athens; Athens Medical Center); E Fradelos (Athens; Athens Naval and Veterans Hospital); N Michalopoulos (Athens; Attikon University General Hospital); N Dimitrokallis (Athens; Evaggelismos General Hospital); A Paspala (Athens; Evgenideio Hospital); P Christodoulou (Athens; General Hospital Asklepieio Voulas); EC Tampaki (Athens; KAT Athens General Hospital); D Schizas (Athens; Laiko University Hospital); K Kontzoglou (Athens; National and Kapodistrian University of Athens); M Spartalis (Athens; Sotiria General Hospital of Thoracic Diseases); M Billis (Filiates; General Hospital of Filiates); N Tsakiridis (Florina; Florina General Hospital 'Eleni Th. Dimitriou'); A Karakosta (Ioannina; University Hospital of Ioannina); D Panagopoulos (Kyparissia; General Hospital of Messinia, Hospital Unit of Kyparissia); G Koukoulis (Larissa; General Hospital of Larissa 'Koutlimpaneio and Triantafylleio'); G Christodoulidis (Larissa; General University Hospital of Larissa); K Bouchagier (Patras; General University Hospital of Patras); V Mousafeiris (Patras; Karamandaneio Prefecture Children Hospital of Patras); A Papadopoulos (Piraeus; General Hospital of Nikaia); L Katsiaras (Piraeus; Metaxa Cancer Hospital); N Zampitis (Piraeus; Tzaneio General Hospital); O Ioannidis (Thessaloniki; George Papanikolaou General Hospital of Thessaloniki); M Drogouti

(Thessaloniki; O Agios Dimitrios General Hospital); C Kaselas (Thessaloniki; Papageorgiou General Hospital); D Lytras (Volos; Achillopoyleio General Hospital of Volos).

Guatemala: M Aguilera-Arevalo (Guatemala City; Hospital General San Juan De Dios); L Talé-Rosales (Guatemala City; Hospital Juan Jose Arevalo Bermejo); ST Torres Rodríguez (Guatemala City; Hospital de Referencia Nacional de Enfermedades Respiratorias).

India: N Krishnappa (Bangalore, Karnataka; BGS Global Institute of Medical Sciences); S Kumar Venkatappa (Bangalore; Victoria Hospital); TS Mishra (Bhubaneswar; All India Institute Of Medical Sciences - Bhubaneswar); Y Sakaray (Chandigarh; Postgraduate Institute of Medical Education & Research, Chandigarh, India); R Kottayasamy Seenivasagam (Coimbatore; PSG Institute of Medical Sciences and Research); R Sharma (DELHI; St Stephen's Hospital); R Gupta (Dehradun; Synergy Institute of Medical Sciences); M Luthra (Delhi; Holy Family Hospital); T Longkumer (Dimapur; Christian Institute of Health Sciences and Research); A Chhabra (Faridkot; Guru Gobind Singh Medical College & Hospital (Baba Farid University of Health Sciences)); T Doma Bhutia (Gangtok; Sir Thutob Namgyal Memorial Hospital Sochakgang); M Pathak (Jodhpur; All India Institute of Medical Sciences (AIIMS), Jodhpur); J Rathod (Karamsad; Shree Krishna Hospital); MK Agrawal (Lucknow; King George's Medical University); D Jain (Ludhiana; Christian Medical College & Hospital); NK Chaudhry (Ludhiana; Satguru Partap Singh Hospital); A Mathew (Madhepura; Madhepura Christian Hospital); P Alexander (Manali; Lady Willingdon Hospital); V Kumar (Manipal; Kasturba Medical College Hospital, Manipal); RD Sharma (Mumbai; Lilavati Hospital & Research Centre); B Sarang (Mumbai; Terna Medical College and Hospital); D Singh (Nandurbar; Chinchpada Christian Hospital); N Gupta (New Delhi; ABVIMS Dr RML Hospital); S Kulkarni (New Delhi; Army Hospital Research & Referral New Delhi); T Rashid (New Delhi; Hamdard Institute of Medical Sciences & Research); L Bains (New Delhi; Maulana Azad Medical College); T Iahmo (Padhar; Padhar Hospital); A Kumar (Patiala; Government Medical College Patiala); M Kumar (Patna; All India Institute of Medical Sciences, Patna); R Abhinaya (Pondicherry; Jawaharlal Institute of Postgraduate Medical Education and Research); VS Jha (Pune; Command Hospital, Southern Command); D Dugar (Raipur; All India Institute of Medical Sciences Raipur); S Basu (Rishikesh; All India Institute Of Medical Sciences); K Singh (SAS Nagar (Mohali) ; BR Ambedkar State Institute of Medical Sciences Mohali); C Mahakalkar (Sawangi (Meghe), Wardha; Acharya Vinoba Bhave Rural Hospital); FQ Parray (Srinagar; Sher-i-Kashmir Institute of Medical Sciences); JA Kalyanapu (Tezpur; Baptist Christian Hospital); M Chisthi (Thiruvananthapuram; Government Medical College Thiruvananthapuram); A Kavalakat (Thrissur; Jubilee Mission Medical College & Research Institute); B Roopavathana. S (Vellore; Christian Medical College & Hospital).

Iran, Islamic Rep.: N Yousefzadeh Kandevari (Bastak; Farabi hospital); M Pourfridoni (Jiroft; Imam Khomeini Hospital).

Iraq: R Raheem Attallah Al_obaidy (Anbar; Heet General hospital); Z Alkhuzaiie (Najaf; Al Batool private hospital); MA Al-Juaifari (Najaf; Al-Najaf Al-Ashraf Teaching Hospital).

Ireland: S Ramjit (Dublin; St James's Hospital); R Tummon (Kerry; University Hospital Kerry).

Israel: S Abu Salem (Jerusalem; Hadassah Medical Center).

Italy: R Sulce (Arezzo; Ospedale San Donato USL Toscana Sud Est); PM Cicerchia (Ariccia; Ospedale dei Castelli (N.O.C.)); E Marra (Aversa; San Giuseppe Moscati); M Rottoli (Bologna; IRCCS Azienda Ospedaliero-Universitaria di Bologna); J Andreuccetti (Brescia; ASST Spedali Civili, Ospedale di Brescia); E Locci (Cagliari; Cagliari University Hospital); F Cappellacci (Cagliari; Chirurgia Generale e Polispecialistica, Cagliari University Ospital 'Duilio Casula'); N Cillara (Cagliari; Santissima Trinità - ATS Sardegna); E Abate (Carate Brianza (MB); Ospedale Vittorio Emanuele III - Carate Brianza); F Ascari (Carpi; Ramazzini); S Romano (Casarano; Francesco Ferrari Hospital); M Veroux (Catania; Azienda Ospedaliero- Universitaria Policlinico San Marco); B Nardo (Cosenza; Azienda Ospedaliera di Cosenza); D Sasia (Cuneo; Santa Croce e Carle Hospital, Cuneo); G Baronio (Esine; ASST Valcamonica Ospedale di Esine); N Fabbri (Ferrara; Azienda Unità Sanitaria Locale di Ferrara); J Martellucci (Firenze; Azienda Ospedaliera Universitaria Careggi); G Canonico (Firenze; Ospedale San Giovanni di Dio); V Lizzi (Foggia; Ospedali Riuniti Azienda Ospedaliera Universitaria); F D'acapo (Forlì; Morgagni-Pierantoni); D Merlini (Garbagnate Milanese; ASST Rhodense - Ospedale di Garbagnate Milanese); A Barberis (Genoa; E.O. Ospedali Galliera); MF Amisano (Genoa; IRCCS Ospedale Policlinico San Martino); A Luzzi (Genoa; Ospedale Villa Scassi); F Palmieri (Gravedona ed Uniti; Ospedale Moriggia Pelascini); CL Bertoglio (Magenta; Ospedale 'G.Fornaroli', ASST-OVEST Milanese); E Baldini (Melzo; Ospedale Santa Maria delle Stelle, ASST Melegnano Martesana); M Ceolin (Milan; Humanitas Research Hospital); P De Nardi (Milan; IRCCS San Raffaele Scientific Institute, Milan); MG Piacentini (Milan; Ospedale Fatebenefratelli e Oftalmico); F Ferrara (Milan; San Carlo Borromeo); F Brucchi (Milan; Sesto San Giovanni Hospital); F Di Marco (Modica; Ospedale Maggiore); N Tamini (Monza; Fondazione IRCCS San Gerardo dei Tintori Monza, Scuola di Medicina e Chirurgia, Università Milano Bicocca); P Anoldo (Naples; Federico II University of Naples); R Patrone (Naples; Istituto Nazionale Tumori Fondazione, Pascale IRCCS); F Selvaggi (Naples; Primo Policlinico di Napoli); G Bellio (Padova; Piove di Sacco Hospital); P Venturelli (Palermo; Policlinico Universitario Paolo Giaccone); L Conti (Piacenza; G. Da Saliceto); L Morelli (Pisa; Azienda Ospedaliero Universitaria Pisana); SMM Basso (Pordenone; Azienda Sanitaria Friuli Occidentale (AS FO)); F Biolchini (Reggio Emilia; Azienda Unità Sanitaria Locale - IRCCS di Reggio Emilia); C Marafante (Rivoli; Ospedale degli Infermi di Rivoli); A Antinori (Rome; Fondazione Policlinico Universitario Agostino Gemelli); M Campanelli (Rome; Policlinico Tor Vergata Hospital, Rome); P Lapolla (Rome; Policlinico Umberto I Sapienza University of Rome); G Palomba (Salerno; San Giovanni di Dio e Ruggi d'Aragona); L Cardinali (San Benedetto del Tronto; Madonna del Soccorso Hospital); E Andolfi (Sansepolcro; Valtiberina); L Verre (Siena; Azienda Ospedaliero Universitaria Senese); G Poillucci (Spoleto (PG); San Matteo degli Infermi); E Pontecorvi (Sulmona; SS Annunziata); S Novello (Treviso; Ospedale Ca' Foncello - Università di Padova (DISCOG)); M Santarelli (Turin; Città della Salute e della Scienza); L Cobellis (Vallo Della Lucania; Casa di Cura Prof. Dott. Luigi Cobellis); G letto (Varese Lombardy; University of Insubria, Ospedale di Circolo e Fondazione Macchi (Varese)); F Pederiva (Varese; Filippo Del Ponte Hospital, University of Insubria); A Iacomino (Venezia; Ospedale Civile - Santi Giovanni e Paolo); D Verdi (Venice; Mirano Hospital); A Broglia (Voghera; Ospedale Civile di Voghera); P Cianci (andria; Lorenzo Bonomo); M Angelucci

(rome; policlinico universitario campus bio medico of rome); G Calini (udine; santa maria della misericordia).

Japan: H Yonekura (Aichi; Fujita Health University Bantane Hospital).

Jordan: S Alananzeh (Ajloun; Al Iman Hospital); A Qasem (Amman; Al-Basheer Hospital); Y Alawneh (Amman; Ibn Al Haitham Hospital); S Al-Tahayneh (Amman; Islamic Hospital); A Khamees (Amman; Jordan University Hospital); MEH Albanna (Amman; Marka Specialty Hospital); RKZ Almahadin (Amman; Prince Hamza hospital); M Mahafdah (Ar Ramtha; King Abdullah University Hospital/ Jordan University of Science and Technology); O Mansour (As-Salt; Al Hussain New Salt Hospital); M Mubarak (Irbid; Ar Ramtha Governmental Hospital); A Alrababah (Irbid; Princess Basma Hospital).

Kazakhstan: M Kulimbet (Almaty; City Clinical Hospital No.7, Asfendiyarov Kazakh National Medical University); I Fakhradiyev (Almaty; JSC 'Central Clinical Hospital', Asfendiyarov Kazakh National Medical University).

Kenya: R Parker (Bomet; Tenwek Hospital).

Lebanon: D Rahme (Beirut; Hopital Libanais Geitaoui); H Hamdar (Jbail ; Maritime Hospital).

Libya: W Ebrahim (Albayda; Albayda Medical Center); M Saleh (Benghazi; Al-jalaa Teaching/Trauma Hospital); S Alsaeti (Benghazi; Benghazi Children's Hospital); R Michael (Benghazi; Benghazi Medical Center); A Bojazyah (Darna; Al-Wahda Hospital); H Bileid Bakeer (Gharyan; Gharyan Central Hospital); M Abudabbous, N Albahloul (Misurata; Misurata Central Hospital); A Egdeer (Nalut; Nalut Central Hospital); H Embarek (Sebha; Al-Majd Clinic); M Abdelkabir (Sebha; Aseel Alghad Clinic); H Idheiraj (Sebha; Sabha Medical Center); R Salim (Tobruk; Tobruk Medical Center); K Ayad (Tripoli; Alkhalil hospital); A Alragheai (Tripoli; Metiga Hospital); S Egreaa (Tripoli; Sabratha teaching hospital); S Timmalah (Tripoli; Tripoli Medical Center/ Tripoli University Hospital); N Lahmer (Zawia; Zawia Teaching Hospital); N Ben Hasan (Zliten; Zliten Teaching Hospital).

Lithuania: D Venskutonis (Kaunas; LUHS Kaunas Hospital); A Dauksa (Kaunas; Lithuanian University of Health Sciences Kaunas Clinics); A Gulla (Vilnius; Vilnius University Hospital).

Madagascar: F Rasoaherinomenjanahary (Antananarivo; Joseph Ravoahangy Andrianavalona Hospital).

Malaysia: MS Mohd Shah (Kelantan; Hospital Universiti Sains Malaysia); R Noor (Kota Bharu; Hospital Raja Perempuan Zainab II); SN Loke (Kuching, Sarawak; Sarawak General Hospital); MA Yunus (Malacca; Hospital Jasin); H Amin-Tai (Serdang; Hospital Pengajar Universiti Putra Malaysia (HPUPM)).

Mali: KS Dembele (Ségou; District Hospital of Tominian).

Malta: J Psaila (Msida; Mater Dei Hospital).

Mexico: CM Nuño-Guzmán (Guadalajara; Antiguo Hospital Civil de Guadalajara); LA Flores Chávez (Guadalajara; Clínica de Especialidades más Centro de Cirugía Simplificada); CM Nuño-Guzmán, G Yanowsky-Reyes (Guadalajara; Hospital Civil Fray Antonio Alcalde); A Gonzalez Ojeda (Guadalajara; Hospital de Especialidades, CMNO-IMSS); G Ambriz González (Guadalajara; UMAE Hospital de Pediatría Centro Medico Nacional de Occidentales IMSS); EE Lozada Hernandez (León; Hospital Regional e Alta Especialidad del Bajío); M Trejo-Avila (Mexico City; Hospital General Dr. Manuel Gea González); C Moreno-Licea (Mexico City; Instituto Nacional de Ciencias Médicas y Nutrición

'Salvador Zubirán'); A Navarrete-Peón (Pachuca; Sociedad Española de Beneficencia); M Noguez Castillo (Querétaro; Hospital de especialidades del niño y la mujer); A Ramos-De la Medina (Veracruz; Hospital Español Veracruz).

Morocco: I Gouazar (Marrakech; Centre Hospitalier Universitaire Mohammed VI, Marrakech); A Ouachou (Rabat; Centre Hospitalier Universitaire Ibn Sina Rabat).

Namibia: PR Nashidengo (Windhoek; Windhoek Central Academic Hospital).

New Zealand: S Rennie (Masterton; Wairarapa Hospital); M Haimona (Wellington; Wellington Regional Hospital); F Fadzlullah (Whanganui; Whanganui Hospital); L Paterson (Whangarei; Whangarei Hospital).

Nigeria: A Adeyeye (Ado Ekiti; Afe Babalola University Multi-System Hospital); J Olaogun (Ado-Ekiti; Ekiti State University Teaching Hospital); N Oloko (Bauchi; Abubakar Tafawa Balewa University Teaching Hospital Bauchi); P Agbonrofo (Benin City; University of Benin Teaching Hospital); A Abiodun (Bida; Federal Medical Centre Bida); U Ezomike (Enugu; University of Nigeria Teaching Hospital); SA Sani (Gwagwalada; University of Abuja Teaching Hospital); TA Lawal (Ibadan; University College Hospital); AO Lawal (Idi Araba; Lagos University Teaching Hospital); AI Okunlola (Ido Ekiti; Federal Teaching Hospital, Ido Ekiti); OM Williams (Ikeja; Lagos State University Teaching Hospital); A Adisa (Ile-Ife; Obafemi Awolowo University Teaching Hospitals Complex); T Mohammed (Ilesa; Obafemi Awolowo University Teaching Hospitals Complex Wesley Guild Hospital Unit); P Elemile (Ilishan-Remo; Babcock University Teaching Hospital); II Aremu (Ilorin; General Hospital); L Abdur-Rahman (Ilorin; University of Ilorin Teaching Hospital); JG Makama (Kaduna; Barau Dikko Teaching Hospital); IU Garzali (Kano; Aminu Kano Teaching Hospital); TT Ibiyeye (Lokoja; Federal Teaching Hospital Lokoja); OH Ekwunife (Nnewi; Nnamdi Azikiwe University Teaching Hospital); OH Ekwunife (Onitsha; Holy Rosary Specialist Hospital); O Ojewuyi (Osogbo; UNIOSUN Teaching Hospital); I Ogundele (Sagamu; Olabisi Onabanjo University Teaching Hospital); M Daniyan (Zaria; Ahmadu Bello University Teaching Hospital).

North Macedonia: T Risteski (Skopje; University Clinic for Pediatric Surgery).

Oman: H Al-Aamri (Ibra; Ibra Hospital); AH ALSharqi (Muscat; Sultan Qaboos University Hospital); M Al Hinai (Nizwa; Nizwa Hospital).

Pakistan: S Ahmed (Islamabad; Dr Akbar Niazi Teaching Hospital); SH Waqar (Islamabad; The Pakistan Institute of Medical Sciences); M Shahid (Karachi; PAF Faisal Hospital); F Ashraf (Karachi; Patel Hospital); AN Syed (Karachi; The Indus Hospital); AS Ammar (Lahore; Bahria International Hospital, Bahria Orchard); K Hayat (Lahore; Services Hospital Lahore); N Talat (Lahore; The Children's Hospital & The Institute of Child Health Lahore); W Mabood (Peshawar; Mercy Teaching Hospital); HW Bhatti (Rawalpindi; Benazir Bhutto Hospital); M Usman Malik (Sargodha; District Headquarter & Teaching Hospital - Sargodha).

Palestine: B Mohamad (Bethlehem, West Bank; Beit Jala Governmental Hospital (Al Hussein)); A Alwali (Gaza; Al-Shifa Hospital); A AbuNemer (Gaza; Nasser Hospital); S Alijla (Gaza; Palestine Red Crescent Society - Al-amal Hospital); H Ayesh (Hebron, West Bank; Al-Ahli Hospital); H Abu-Arish (Hebron, West Bank; Governmental Hebron Hospital-Alia); D Rabaia, A Jaber (Jenin, West Bank; The

Martyr Dr. Khalil Sulaiman Hospital (Jenin Governmental Hospital)); M MohammedAli (Nablus, West Bank; Rafidia Hospital); A Attili (Tulkarm, West Bank; Martyr Thabet Thabet Governmental Hospital).

Paraguay: OM Cuenca Torres (Asuncion; Hospital de Clínicas, II Cátedra de Clínica Quirúrgica, Universidad Nacional de Asunción).

Peru: V Panduro-Correa (Huánuco; Hospital Regional Hermilio Valdizán Medrano); L Fuentes Rivera Lau (Lima; British American Hospital); CF Huaroto Landeo (Lima; Clinica Internacional); G Mendiola (Lima; Hospital Santa Rosa de Lima); Y Carpio Colmenares (Lima; SANNA - Clínica El Golf); C Shiraishi Zapata (Paíta; Hospital I Miguel Cruzado Vera EsSalud); R Díaz-Ruiz (Piura; Jose Cayetano Heredia III Regional Hospital).

Poland: Ł Nawacki (Kielce; Wojewódzki Szpital Zespólny w Kielcach); M Kisielewski (Krakow; 5th Military Clinical Hospital); Z Orzeszko (Krakow; Brothers Hospitalers Hospital); M Matyja (Krakow; Jagiellonian University Medical College); W Krawczyk (Sosnowiec; Wojewódzki Szpital Specjalistyczny nr 5 im. Św Barbary); M Walędziak (Warsaw; Military Institute Of Medicine); JK Zajac (Wroclaw; Regional Specialist Hospital in Wroclaw); F Brzeszczyński (Łódź; Copernicus Memorial Hospital).

Portugal: S Henriques (Almada; Hospital Garcia de Orta); J Frazão (Amadora; Hospital Prof. Doutor Fernando Fonseca, E.P.E.); S Gaspar Reis (Barreiro; Centro Hospitalar Barreiro Montijo, EPE); AR Mateus Loureiro (Caldas da Rainha; Hospital das Caldas da Rainha - Centro Hospitalar do Oeste, E.P.E.); M Reia (Elvas; Hospital Santa Luzia Elvas); J Pinho (Figueira da Foz; Hospital Distrital da Figueira da Foz); DG Alves (Funchal; Hospital Dr. Nélio Mendonça); R Silva Borges (Horta; Hospital da Horta, E.P.E.); E Borges (Lisbon; Centro Hospitalar Lisboa Norte); M Nunes (Matosinhos; Unidade Local de Saude de Matosinhos - Hospital Pedro Hispano); A Faustino (Ponta Delgada; Hospital do Divino Espírito Santo); G Fialho (Portalegre; Hospital Doutor José Maria Grande); J Dias-Ferreira (Porto; Centro Hospitalar e Universitário de São João); M Santos (Porto; Hospital da Prelada); J Marques Antunes (Santa Maria da Feira; Centro Hospitalar Entre o Douro e Vouga); H Devesa (Santarem; Hospital de Santarem); J Ricardo (Santiago do Cacém; Hospital do Litoral Alentejano); R Branquinho (Tomar; Centro Hospitalar Médio Tejo); J Fernandes (Vila Franca de Xira; Hospital Vila Franca de Xira); J Pereira-Macedo (Vila Nova de Famalicao; Centro Hospitalar do Medio Ave); B Vieira (Vila Real; Centro Hospitalar de Trás-os-Montes e Alto Douro, E.P.E.).

Qatar: CE Guldogan (Doha; Turkish Hospital).

Romania: ST Makkai-Popa (Brasov; Regina Maria); F Grama (Bucharest; Coltea Clinical Hospital); EA Toma (Bucharest; Elias Emergency Hospital); I Negoii (Bucharest; Emergency Clinical Hospital Bucharest); M Muresan (Cluj-Napoca; Medicover Hospital Cluj).

Russian Federation: V Kakotkin (Kaliningrad; Immanuel Kant Baltic Federal University, Regional Clinical Hospital); A Bedzhanyan (Moscow; Petrovsky National Research Centre of Surgery); S Katorkin (Samara; Hospital Surgery Clinic of Samara State Medical University); A Butyrskii (Simferopol; Municipal Emergency Hospital No.6); V Ten (Yuzhno-Sakhalinsk; Private healthcare institution 'RZD-Medicine').

Rwanda: N Christian (Huye, Gisagara; Butare university teaching hospital (CHUB)); C Mpirimbanyi (Kigali; Kibagabaga Hospital); A Costas-Chavarri (Kigali; Rwanda Military Hospital).

Saudi Arabia: N Alzerwi (Al-Majmaah; King Khalid General Hospital); A Shabkah (Jeddah; International Medical Center); A Nawawi (Jeddah; King Abdulaziz University Hospital); N Alzerwi (Riyadh; King Salman Hospital in Riyadh); S Chowdhury (Riyadh; King Saud Medical City); DY Alalawi (Tabuk; King Salman Armed Forces Hospital); S Awad (Taif City; King Faisal Medical Complex).

Serbia: A Karamarkovic (Belgrade; Zvezdara University Medical Center).

Slovenia: JA Košir (Ljubljana; University Medical Centre).

South Africa: M Flint (Cape Town; Groote Schuur Hospital); A Victor (Cape Town; Karl Bremer Hospital); SS Verhage (Cape Town; Khayelitsha District Hospital); F Gool (Cape Town; Mitchell's Plain District Hospital); T Mabogoane (Cape Town; Victoria Hospital Wynberg); B Phakathi (Durban; King Edward VIII Hospital); V Pillay (Durban; Stanger Hospital); R Jayakrishnan (East London; Cecilia Makiwane Hospital); Y Manickchund (East London; Frere Hospital); O Jolayemi (Empangeni; Ngwelezana Hospital); H Stark (George; George Hospital); C Molewa (Johannesburg; Edenvale); H Wain (Pietermaritzburg; Edendale Hospital); D Montwedi (Pretoria; Kalafong Academic Hospital); G De Wee (Upington; Dr Harry Surtie Hospital); C Dempers (Worcester; Worcester Provincial Hospital).

Spain: H Aguado López (Albacete; Hellín Hospital); MDM Martí-Ejarque (BARCELONA; Hospital Universitari Sagrat Cor); A Torroella (Barcelona; HM Nou Delfos); O Martin Sole (Barcelona; Hospital Sant Joan de Deu); A Landaluce-Olavarria (Bizkaia; Hospital Urduliz); V Alonso (Burgos; Hospital Universitario de Burgos); Á Fernández Camuñas (Ciudad Real; Hospital General Universitario de Ciudad Real); M Estaire Gómez (Leganés; Severo Ochoa University Hospital); AG Barranquero (Lleida; Hospital Universitari Arnau de Vilanova); L Marquez (Madrid; Hospital Central de la Cruz Roja San Jose y Santa Adela); P Serrano Méndez (Madrid; Hospital Clinico San Carlos); A Vilar (Madrid; Hospital Universitario Principe de Asturias); J Guevara (Madrid; Hospital Universitario la Paz); AM Minaya Bravo (Madrid; Hospital del Henares); JL Rodicio Miravalles (Oviedo; Hospital Universitario Central de Asturias (HUCA)); R Díaz Pedrero (Rivas-Vaciamadrid, Madrid; Hospital Universitario HM Rivas); L Tallon-Aguilar (Sevilla; Hospital Universitario Virgen del Rocío); A Curado Soriano (Seville; Hospital Universitario Virgen Macarena); Z Balciscueta (Valencia; Hospital Arnau de Vilanova); D Moro-Valdezate (Valencia; Hospital Clínico Universitario de Valencia); B De Andrés-Asenjo (Valladolid; Hospital Clínico Universitario de Valladolid); A Vazquez Melero (Vitoria-Gasteiz; Hospital Universitario Araba); J Escartin (Zaragoza; Hospital Royo Villanova); C Gracia-Roche (Zaragoza; Hospital Universitario Miguel Servet).

Sri Lanka: S Srishankar (Anuradhapura; Teaching Hospital Anuradhapura); D Wickramasinghe (Colombo; National Hospital of Sri Lanka); U Jayarajah (Dehiwala; Colombo South Teaching Hospital); S Gobishangar (Jaffna; Teaching Hospital, Jaffna); W Wijenayake (Werahera; University Hospital, Kotelawala Defence University).

Sudan: I Abdalla (Ed Dueim; Ed Dueim Teaching Hospital); S Ibrahim Tour Harakan (El Geneina; El Geneina teaching Hospital); MM Yassin (Gadarif city; Gadarif teaching hospital); Z Aljalabi (Kassala; Police hospital); I Adel (Khartoum; Bashair Teaching Hospital); IMG Ahmed (Khartoum; Ibrahim Malik Teaching Hospital); M Hajalamin (Khartoum; Omdurman Teaching Hospital); EE Abuobaida Banaga

Hag El Tayeb (Khartoum; Ribat university hospital); HA Fadlalmola (Khartoum; Soba University Hospital); E Alkhalifa (Wad Madani; University of Gezira Hospital).

Sweden: H Zaigham (Malmö; Skåne University Hospital); M Nikberg (Vasteras; Västmanlands Hospital Västerås).

Switzerland: P Probst (Frauenfeld; Spital Thurgau AG); E Gialamas (Geneva; Geneva University Hospitals); J Gass (Luzern; Luzerner Kantonsspital); A Tampakis (Olten; Kantonsspital Olten); G Peros (Winterthur; Kantonsspital Winterthur); MA Schneider (Zurich; University Hospital of Zurich).

Syrian Arab Republic: MA Farho (Aleppo; Abd Al Wahab Agha Hospital); AA Kayali (Aleppo; Al-Shahbaa Private Hospital); M Aloulou (Aleppo; Aleppo Private Hospital); A Ghazal (Aleppo; Aleppo University Hospital); B Alsaïd (Damascus; Al Assad university hospital); M Klib (Damascus; Al-Mouwasat University Hospital); H Dalati (Damascus; Children's University Hospital); S Jomaa (Damascus; Damascus Hospital); Y Alhammoud (Homs; Al-Basel Specialized Hospital in Karm El-Louz); S Abbas (Homs; The Military Hospital); G Hneino (Latakia; Al Saydeh Surgical Hospital); I Ali (Latakia; National Hospital); G Bashour (Latakia; Othaman Hospital); A Hammed (Latakia; Tishreen University Hospital).

Thailand: S Techapongsatorn (Bangkok; Vajira hospital).

Togo: F Alassani (Lomé; CHU Sylvanus Olympio).

Tunisia: A Sebai (Tunis; La Rabta Hospital).

Turkey: GC Bulbuloglu (Adana; Adana Seyhan State Hospital); MA Koç (Ankara; Ankara University Medical School); MY Uzunoglu (Bursa; Bursa City Hospital); B Yigit (Elazig; Elazig Fethi Sekin City Hospital); AN Sanli (Gaziantep; Abdulkadir Yuksel State Hospital); GK Kurtoglu (Istanbul; Acibadem Altunizade Hospital); E Tuzuner (Istanbul; Acibadem Maslak Hospital); Y Altinel (Istanbul; Bagcilar Research And Training Hospital); ÖP Zambak Mutlu (Istanbul; Bahçelievler State Hospital); RE Sönmez (Istanbul; Istanbul Medeniyet University, School of Medicine); S Bektas (Istanbul; Istanbul Medipol University Hospital); E Erginöz (Istanbul; Istanbul universty - Cerrahpaşa Medical faculty); A Özcan (Istanbul; Kanuni Sultan Suleyman Training and Research Hospital); Y Tosun (Istanbul; Kartal Dr. Lutfi Kirdar Training and Research Hospital); İH Özata (Istanbul; Koç University Medical School); TK Uprak (Istanbul; Marmara University, School of Medicine); E Unal (Istanbul; Sehit Prof.Dr. İlhan Varank Training and Research Hospital); N Kiziltoprak (Istanbul; Sultan 2. Abdulhamid Han Training and Research Hospital, University of Health Sciences); M Ergenç (Istanbul; Sultanbeyli State Hospital); MT Demirpolat (Istanbul; University of Health Science Umraniye Education and Research Hospital); B Citgez (Istanbul; Uskudar University Faculty of Medicine, Memorial Hospital); H Ulman (Izmir; Bakircay University Cigli Training and Research Hospital); YK Şen (Izmir; University of Health Sciences Izmir Bozyaka Training and Research Hospital); E Colak (Samsun; Samsun University Samsun Training and Research Hospital); N Kavak (Zonguldak; Zonguldak Bulent Ecevit University School of Medicine Research and Training Hospital); E Kamer (İzmir; University of Health Sciences Tepecik Training and Research Hospital).

Uganda: I Mubezi (Iganga; Iganga district hospital); H Lule (Kigumba; Kiryandongo Hospital); S Stonelake (Luwero; Kiwoko Hospital).

United Kingdom: CS Ong (Bangor, North Wales; Ysbyty Gwynedd); P Patel (Barrow in Furness; Furness General Hospital); S Dindyal (Basildon; Basildon University Hospital); F Georgiades (Bedford; Bedford Hospital); J Abbasy (Birmingham; Heartlands Hospital); M Kaur (Brighton; Royal Sussex County Hospital); B Martin (Bristol; Bristol Royal Hospital for Children); M Chauhan (Camberley; Frimley Health NHS FT - Frimley Park); S Ahmed (Chester; Countess of Chester Hospital); M Tutton (Colchester; Colchester Hospital University); N Chidumije (Coventry; University Hospitals Coventry and Warwickshire NHS Trust); W Al-Khyatt (Derby; Royal Derby Hospital); A Sukumar (Dudley; Russell's Hall Hospital); H Kamal (Dundee; Ninewells Hospital); A Nada (Durham; University Hospital North Durham); A Chaudhary (Exeter; Royal Devon and Exeter Hospital); M Bogdan (Great Yarmouth; James Paget Univeristy NHS Foundation Trust Hospital); M Peter (Huddersfield; Huddersfield Royal Infirmary); J Walshaw (Hull; Hull University Teaching Hospitals NHS Trust); M Ewedah (Ilford; King George Hospital); L Rampersad (Larbert; Forth Valley Royal Hospital); A Peckham-Cooper, NS Blencowe (Leeds; Leeds Teaching Hospitals); R Lunevicius (Liverpool; Aintree University Hospital); P Panahi (London; Ealing Hospital); E Baili (London; Guy's and St Thomas' Hospitals); K Theodoropoulou (London; Homerton University Hospital); P Kapsampelis (London; Kingston); MMH Mohammed (London; Queen Elizabeth Hospital, Woolwich); C Parmar (London; The Whittington Hospital); MMH Mohammed (London; University Hospital Lewisham); C Smart (Macclesfield; Macclesfield District General Hospital); P Wilson (Manchester; Wythenshawe Hospital); F Gareb (Margate; Queen Elizabeth the Queen Mother Hospital Margate); G Sundaram Venkatesan (Middlesbrough; James Cook University Hospital); C Hidalgo Salinas (Morecambe; Royal Lancaster Infirmary); S Tingle (North Shields; Northumbria NHS Hospital Trust); N Marzouqa (Nottingham; Queens Medical Centre); T Theivendrampillai (Oxford; John Radcliffe Hospital); M Zahed Abdalla (Portsmouth; Queen Alexandra Hospital); A Rahman (Redhill; East Surrey Hospital); M Abdelkarim (Rhyl; Glan Clwyd Hospital); O Whitehurst (Salford; Salford Royal Hospital); E Tokidis (Sheffield; Sheffield Teaching Hospital NHS Foundation Trust); S Bandyopadhyay (Southampton; Southampton General Hospital); B Al-Sarireh (Swansea; Morriston Hospital Swansea); A Salam (Walsall; Walsall Manor Hospital).

United States: M Sulciner (Boston; Brigham and Women's Hospital); G Chang (Chicago, IL; Mount Sinai Hospital); A Alecci (Chicago, IL; Rush University Medical Centre); HE Rice (Durham, NC; Duke University Medical Center); D Ridder (Honolulu, Hawaii; The Queen's Medical Center); K McKenzie (Jamaica; Jamaica Hospital); A Choudhry (Syracuse; SUNY Upstate University Hospital).

Yemen, Rep.: MY Abdualqader (Hajjah; Kowaydina hospital); B Alshaikh (Sana'a; Al-Thawra Modern General Hospital).

Collaborators (listed by country and city):

Albania: B Ibi, S Faber (Korca; Regional Hospital of Korca); I Dajti (Tirana; University hospital Koco Gliozheni).

Algeria: K Bensmain, ZR Benamrouche , IE Boumakhlouf, IE Boudis, H Abdoun , M Benamrouche , M Saidani (Algiers; CHU Isaad Hassani); Z Djama, A Chied, HA Mimouni (Constantine; university

hospital abdelhamid ben badis); AK Awad, B Radja, B Abdennour , MN Bouhafs, MEA Meghaizerou (Oran; EHU-1st November 1954).

Argentina: P Carmignani, J Mondino, R Figueroa, J Morales, FR Pascual (Allende, Cordoba; Sanatorio Allende - Sede Cerro); MA Bequis, F Suldrup, C Korzin, J Napoli , N Feijoo, F Mahnic, ME Duran, L Chantada, C Brandi, JF Viñas, F Lucero, C Samojeden, LJ Caram, S Bertone, F Corvatta (Buenos Aires; Hospital Italiano de Buenos Aires); L Garciandia (Buenos Aires; Hospital Universitario CEMIC); M Rius, S Matthiess, J Paredes, MC Kalaydjian , A Veira (Buenos Aires; Hospital municipal de vicente López); MA Fernández Zurita, JI Valenzuela (City of Buenos Aires; Hospital Velez Sarsfield); S Gomez, GR Viscido, MA Doniquian (Cordoba; Clinica Universitaria Reina Fabiola); R Badra, JS García, CI Ferrero, M Garcia, L Granero (Cordoba; Sanatorio Allende - Sede Nueva Cordoba); M Pagani (San Francisco; Clinica Regional del Este).

Aruba: M Gosselink, J Ringers (Oranjestad; Dr. Horacio E Oduber Hospital).

Australia: M Lie, B Mao, C Stennard, EMA Murphy, JE Do, M Harris, B Fosh (Adelaide; Northern Adelaide Local Health Network); M Watson, J Petric, M Maclean, XY Po, M Pham, D Patterson, V Gunasaegaram, E Hopping, P Holt, JA Duffield (Adelaide; Royal Adelaide Hospital); E Schmidt, R Colbran (Brisbane; Princess Alexandra Hospital); V Liu, E Tan, JST Tefay, R Shen, S Bowman (Brisbane; Queen Elizabeth 2 Jubilee Hospital); D Mitchell, M Kelly, A Edmundson (Brisbane; Surgical Treatment and Rehabilitation Service); H Iswariah, R Franz, M Chandrasegaram, P Yuide, SS Hlaing (Brisbane; The Prince Charles Hospital (TPCH)); S Abeykoon, D Kaushal (Campbelltown; Campbelltown Hospital); C Leung (Canberra; Calvary Hospital); S Davis, NF Franco (Canberra; Canberra Hospital); T Rawther, R McClen, W Petrushnko, E Roussos, K Das (Coffs Harbour NSW; Coffs Harbour Health Campus); S Stevens, F Alimri, S McClintock, J Maritz (Colac; Colac Area Health); S Hariharan, S Laura, B Wang, V Ng, J Linker, A Li, I Dong, R Bhatia, S Cai, WKH Lai, AC Dawson, SYD Chia, M Binks, N Tran, SHM Ng, D Shen, EWY Lun, E Reid, J Cui, M Roussos (Gosford; Gosford Hospital); M Issa, M Anandan, P Devlin, U Naidoo (Hamilton; Hamilton Base Hospital); B Balaravi Pillai, D Abeyirigunawardana , T Parker, T Valizadeh Elizeh (Melbourne; Angliss Hospital); D Liu, S Ng, J Jones, O Ladlow, JV Maida, D Proud, A Vu, N Shulman, L Bromley, V Muralidharan, K Hall, C Cheong, C Jamieson-Grigg (Melbourne; Austin Hospital); A Hilder, T Manickam, L Barnard (Melbourne; Box Hill Hospital); K Jaffry, A Gray, A Lim, R Kattini (Melbourne; Casey Hospital); M Bickford, S Kenworthy (Melbourne; Knox Private Hospital); A Crowe, J Zhu (Melbourne; Maroondah Hospital); M Pacilli, A Comella, K Taghavi, R Nataraja, SJA Robinson (Melbourne; Monash Childrens Hospital); D Lowen, A Khan, S Samadi, S Tan, AL Surkitt, S Condron, E Haege, E Francis, A Boynes, S Gill, B D'Souza, H Xiao, E Fraser, J Wong, S Fennelly, K Mori, M Muir, Y Huang, R Pajtak (Melbourne; Northern Hospital); J You, C Banal, T Abelman, N Chen, L Chong, H Jalilehvand, W Santucci, B McKay (Melbourne; St Vincent's Hospital); I Murshed, D Makary, L Green, M Wichmann, A Lim, M Kang, EA Dontoh (Mount Gambier; Mount Gambier and Districts Health Service); P Walker, A Fani, S Sundararajan, E Downes, A Davis, M Bajwa, R Geow, KK Sim (Mount Nasura; Armadale Health Service); S Smith, L Peters, S Zhang, ACK Cheung, I Caitens (Newcastle; Calvary Mater Newcastle); M Ishak, E Zhang, KKA Yu, L Beukes, L Kang, , F Amico (Newcastle; John Hunter Hospital); M Bowles, E Downing, B Williams, G Cox, V Bakshi

(Robina; Robina Hospital); N Ensor (Sale; Sale Hospital / Central Gippsland Health Service); J Ng, CT Petcu, J Hwang (Southport; Gold Coast University Hospital); TJ Hugh, S Quoy, C Knee, D Siriwardena, C Tung, G Smith (Sydney; Royal North Shore Hospital); O Camilleri, J Vu, J Hong, C Cornwell (Sydney; The Royal Prince Alfred Hospital); E Chan, C Killoran, J Mackenzie, D Fry (Toowoomba; Toowoomba Hospital); V Mahendrarvarman, A Shanmugalingam, C Li, L Allan, P Shivashankar, H Pleass (Westmead; Westmead Hospital); V Lin, R McGee, B Bereket Araya, CW Un, Y De Silva, D Maan, J Wang, M Kwon, R Kibuuka, J Chew, J Siu, J Barklmore (Wyong; Wyong Public Hospital).

Austria: N Klammer, R Schmidt-Branden, P Tschann, P Horvath (Feldkirch; Landeskrankenhaus Feldkirch); N Koter, G Moitzi, E Wallner, C Allmer, F Aigner (Graz; Barmherzige Brüder Krankenhaus, Graz); J Kahn, A Belarmino, R Sucher, V Wolfschluckner, G Singer (Graz; Medical University of Graz); R Függer, M Biebl, A Punzengruber, H Fehrer (Linz; Ordensklinikum Linz Elisabethinen); A Binder, E Haiden, P Riedl, M Enßlin (Tulln; Universitätsklinikum Tulln).

Bangladesh: K Nahar, T Akter (Dhaka; Dhaka Medical College Hospital); A Oosterkamp (Dinajpur; Lamb Hospital).

Benin: G Gbessi, J Avakoudjo, M Fiogbe, P Assouto, SP Chigblo (Cotonou; Centre National Hospitalier et Universitaire Hubert Koutoukou Maga); H Aouagbe Behanzin, M Seto, G Mevognon (Cotonou; Hopital de Menontin); M Agbadebo, A Hada, SFA Houndji (Dassa-Zoumè; Hôpital de Zone de Dassa-Zoumè); E Bara (Kandi; Hôpital de zone de Kandi); AB Yevide, ZF Tamou, E Hatangimana, B Cakpo, R Soglonou (Klouékanme; Hopital de Zone de Klouékanme); TK Hessou, SR Tobome, M Zounon (Natitingou; Centre Hospitalier Départemental de l'Atacora); AM Hodonou, C Bokossa, F Hounde, R Alinde (Parakou; Centre Hospitalier Universitaire Borgou Alibori); F Dossou, R Goudou, ACS Toi, G Natchagande (Porto Novo; Centre Hospitalier Universitaire et Departemental Oueme Plateau).

Bosnia and Herzegovina: M Stjepanovic (Doboj; Genera Hospital 'Sveti aposto Luka' Doboj); O Čančar, M Pejović (Foča; University Hospital Foča); J Miskovic, M Boras, M Kajic, V Dragisic, Z Brekalo, I Mikulic, N Soldo, M Bevanda, M Faletar (Mostar; SKB University Clinical Hospital Mostar); M Salibašić, E Hodžić, E Halilović, M Kruščica, E Bičakčić (Sarajevo; Clinical Center University of Sarajevo); A Cerovac, H Škiljo, E Hodžić, O Bedak, M Kalabić, E Begunić (Tešanj; General Hospital Tešanj); A Huremovic, E Alić (Tuzla; University Clinical Center Tuzla).

Brazil: RA Tenfen Carneiro (Francisco Beltrão; Universidade Estadual do Oeste do Paraná).

Bulgaria: D Georgiev, I Fidoshev, V Neykov, E Daleva, I Ilieva (Pleven; Heart and Brain - Pleven Hospital); M Karamanliev, D Dimitrov, A Shanker, P Vladova, MD Shoshkova, A Mehta, M Abdullahi, V Ratheesh, V Kamalathevan, C Wiesner, S Shittu, M Galasyuk, S Shanker (Pleven; University Hospital Dr Georgi Stranski, Medical University - Pleven); M Imirski, A Soumpasis (Plovdiv; MHAT St. Karidad); E Hadzhieva, D Chakarov (Plovdiv; UMHAT Sveti Georgi); T Yotsov, P Kamenova, A Vricheva, I Yotsov (Ruse; University Hospital Medika); E Hristova, K Spassov (Sofia; Fifth City Hospital Sofia - 5th MBAL).

Burkina Faso: A Sanou, M Windsouri, R Doamba, IW Bahikoro, AST Sanon (Ouagadougou; Tengandogo University Hospital).

Burundi: N Ildephonse, N Steve , Y Fulgence , GD Nibogora, C Nimbona , B Paul (Bujumbura; Centre Hospitalo-Universitaire de Kamenge); G Kazobinka, C Rukundo, N Renovat, E Ndizeye , M Dauphin (Bujumbura; Kamenge Military Hospital); FF Irakiza, L Niyidukunda, G Nkunguzi, N Oscar, N Theophile, B Révérien (Bujumbura; Prince Regent Charles Hospital).

Cambodia: S Oum, S Eam (Battambang; Handa Medical Centre).

Cameroon: NS Bibila, NN Cabrel, J Dongmo (Bamenda; Nkwen Baptist Hospital).

Canada: R Spence, G Berger, D Hannedige, C Hoogerboord, A Abidali (Halifax; Queen Elizabeth II); M Mozel, L Monteiro, R Lertnamvongwan, D Konkin, R Kaur (Port Moody; Eagle Ridge Hospital); M Mozel, R Lertnamvongwan, L Monteiro, S MacKenzie, D Konkin (Vancouver; Royal Columbian Hospital).

Chile: MM Modolo, E Sepúlveda, Á Molero, M Perez, LV Torres Bavestrello, ÓA Soublett Rivas , C Carrillo Sarango, L Paredes, P Sornoza, A Pinto, C González (Santiago; Hospital Barros Luco Trudeau).

China: J Wang (Guangzhou; The First Affiliated Hospital of Jinan University).

Colombia: L García-Zambrano, PA Cabrera Rivera, N Paez, SV Agudelo Mendoza, MS Mosquera Paz, A Kadamani Abiyomaa, CF Roman Ortega, F Casas J, B Guerra (Bogota; Fundacion Cardioinfantil-IC); JD Molina Marin, C Maya, C Vasquez Maya, B Dieck, F Zapata (Medellin; Clínica CES); VA Ruiz López, MA Ñañez (Popayan; Hospital Susana Lopez de Valencia); DC Cardona Gomez, A Rojas, DC Patiño García , LI Bolaños, C Pastás, DA Pérez Muñoz (Popayán; Hospital Universitario San José).

Croatia: J Mihanovic, I Bacic, D Vukosav , V Žufić, O Jurić, E Dijan, N Jović, I Čoza, I Rakvin, Z Katusic, T Soric, I Vidić, D Rukavina (Zadar; Zadar General Hospital); I Separovic, R Radojković, J Mavrek (Zagreb; University Hospital Centre Zagreb).

Cyprus: M Theodoridou, A Pilavas, R Moukarzel, R Andreou, N Gouvas, K Lambri, S Charitonos , IC Mylona, G Kokkinos, N Dimitriou, MM El Ghouli Miliotou, V Ioannou, NA Ververidis, N Kalampokis, A Yiallourou, R Sokratous, D Tsiardas , P Evangelou, D Evripidou, C Thrasyvoulou (Nicosia; Nicosia General Hospital).

Czech Republic: T Reichelt, P Hudáč (Decin; Krajská zdravotní as - Hospital Decin); K Akter, L Moolla, F Rudisch, A Ibrahim Hassan, O Ahmad, E ELShennawy, M Shalaby, M Khaled, A Akiba, F Philips, E Bankart, R Elshennawy, A Al Kaddah, H Al Atassi, S Ashry, N Salgadoe (Hradec Kralove; Charles University Hospital); L Majerčák , P Levíček, A Lukáč, L Pánči (Ivančice; Hospital Ivančice); J Roman, L Tulinsky, I Mrazkova, P Ostruszka, A Varga, L Martinek (Ostrava; University Hospital Ostrava); H Novák, J Woleský, P Francúz (Prague; Motol University Hospital).

Dominican Republic: R Rivas, B Calcaño, J Michel, Y Perez, R Ubiñas, P Garcia-Dubus, S Batista, S Strachan (Santo Domingo; CEDIMAT - Centro de Diagnóstico, Medicina Avanzada, Laboratorio y Telemedicina).

Egypt: Y Tanas, Y Kerolous, Y El Okazy, M Mokhtar, M Lotfy, M AL Sayed, H Altabbaa, AGMM Abouelnagah, O Al Shaqran, D M. Awad, A Sabry, G Nagy, E Amer, M Khalil, A El Shamarka, B Sharaf eldin, AAA Aboshosha, A Farrag, H Sherif Farouk Ahmed Hassan, Y Badr (Alexandria; Alexandria Main University Hospital); Y Orabi, M Kamal matter, A Alrifaae (Alexandria; Alexandria

Medical Research Institute); M Elnour , M Zahran, A Aladl, M Bahnacy, Y Seada, M Kotb, A Ragab, Y Farag, L Khalifa, M Elmiesiry , D Abdalaziz, I Maharem, O AbouHiekal, O Hany, S Hanna, Y Dean, A Faisal (Alexandria; Smouha University Hospital); M Mostafa, I Ali, T Sabra, H Ibrahim , A K. Ali, M Osman , A Eltayeb (Assiut; Assiut University Children Hospital); A Morad, MO Herdan , A Abdelshafi, M M. Nathan, M Shalkamy, M Hamada Takrouney, R Sayad, FA Monib, A A Elhars, MM Saad, A Rashad Temerik, AM Abbas, O Mohamed Mokbel, E AbdElBaset, A Barakat, Z Bady, S Arafa, Z Osama, A Elzanaty , S Salama (Assiut; Assiut University Hospital); MEM Madany (Aswan; Aswan University Hospital, Aswan University); A Khaity , R Adel Diab, A Ghazal, A Ehab, A Abd Elsattar (Cairo; Al Zahraa University Hospital); A El-bastwesy (Cairo; Al-Azhar University Hospitals); A Eisa, M Ellesseily, R Radwan, Y Asar, D Waleed, S Tawfik , AF Nixon Fulli , MJI Albert, A Autiak Ayii Chol, AR AbdelHalim, B Azhar, H Al-derume , M Alqadasi, N Alasbahi (Cairo; Cairo University Children's Hospitals (CUSPH & CUCH)); H Abozied, Y Ashour, Y Mohamed , M Abdelmaboud, H Abdelazim, AES El kady, M Omar, A Haty, M Abd Al-Fattah, I Tagreda , IM Kereet, AG Montaser, M Faisal, M Masoud , M ElSayed Metwally (Cairo; EL-Hussein University Hospital, Al-Azhar University, Faculty Of Medicine); ASM Abdelrahman, S Mansour (Cairo; Giza International Hospital); A Nabil, MMA Marei, A Elmosalamy, L ElGebaly, AM Allam, T Awad, H Taher, K Fayed, M Abdelfattah, DH Khattab, N Ali, A Saleh, K Nassim, NK Aly, I Abo Elhagag, M Doss, M Elzayat, Y Samer Morsy, M EIFiky (Cairo; Kasr Al Ainy Faculty of Medicine, Cairo University); M Erfan, M Zaazou , M Reda, M Kouta, M Mohamad Amin, I Guirguis, H Amir , S Aboseif, M Abdelhafez, O Agha, A Khairy , A Dawoud, M Hamoud Almahly, S Mostafa Yassin , A El-Sherbiney, A Adel, H Foda, D Ahmed, AS Elkhodary , AA Mansour (Giza; The Memorial Soaad Kafafi University Hospital); A Elghrieb, M Natey, A Elshazli Mahmoud , A Khalleefah, H Elfeki, M Shalaby, M Sadek, M Abdelmaksoud, M Mostafa, M Waseem, A Adel, A Azam, A Sakr, A Sanad, م عبدالفتاح (Mansoura; Mansoura University Hospital); H Foad, S Elnoamany, S Selim, DS Alrokh, A Hassanin (Menofia; Menofia University Hospital); M Alansary (Qena; Qena University Hospital); A Ragheb, M Fahmy , M Mehanny (Sohag; Sohag University Hospital); AGE Aboelnasr, KMG Mohammed, M Eissa, S Allam, M Kamar (Tanta; Tanta University Hospital); A Asla (Zagazig; Al Ahrar Zagazig Teaching Hospital).

Ethiopia: F Terefe, WA Zerefa, E Gallo , A Yingess, T Kebede, M Mesfin, G Alemayehu , SM Djote, T Girma, DA Muhie (Addis Ababa; Yekatit 12 hospital medical college); E Yeshialem, G Seyfu, F Tsige, A Yeshitila, N Solomon (Deberibirhan; Hakim Gizaw Hospital); M Worku, S Lakew, Y Melkamu, ST Workineh, M Beletachew (Dessie; Dessie Referral Hospital); B Mengesha, G Getachew , I Tesfahun, M Teressa , BG Chimam, AB Aregawi (Hawassa; Hawassa University Comprehensive Specialized Hospital); N S.Bayleyegn, YY Metaferia, TG Moges, D Mengiste, A Teshome Sahilemariam (Jimma; Jimma University Medical Center); B Sime, T Jemal (Yirgalem; Yirgalem Hospital Medical College).

France: E Volpin, H Braham, C Lionel, R Arena, Y Malki (Eaubonne; Hôpital Simone Veil); M Bertrand, A Castaldi, L Theuil, A L'Hostis, M Prudhomme (Nimes; Hôpital Carèmeau); P Riva, A Lapergola, D Mutter, S Perretta (Strasbourg; Nouvel Hopital Civil de Strasbourg).

Gabon: PC Nze Obiang (Libreville; Centre Hospitalier universitaire mère enfant Fondation Jeanne Ebori).

Georgia: Z Demetrashvili, G Devidze, G Pisarevi, L Petashvili, E Ekaladze, N Lekiasvili, I Pipia, A Tvaladze, G Kenchadze, K Khutsishvili (Tbilisi; N.Kipshidze Central University Clinic).

Germany: LD Lee (Berlin; Park-Klinik Weissensee); J Binder, A Denz, C Krautz, M Brunner, M Maak, GF Weber, A Stollberg, D Hackner, S Engel, F Krämer, R Grützmann (Erlangen; Universitätsklinikum Erlangen); M Schüler, J Kleeff, R Rüdric (Halle; University Hospital Halle); A Kirschniak, J Rolinger, S Göller, L Van den Hil, J Miller, H Pehlivan (Moenchengladbach; Kliniken Maria Hilf); M Kießler, N Hüser, D Schippers, M Berlet, M Steffani, M Weber (Munich; Klinikum Rechts der Isar TUM School of Medicine); N Börner, M Albertsmeier, H Arbogast, M Mattis, S Jarmusch, P Zimmermann, U Wirth (Munich; Ludwig Maximilian University of Munich - Großhadern); F Anzinger (Munich; Ludwig Maximilian University of Munich - Innenstadt); FG Bader, M Sohn, ML Koschke, M Busch, N Hielscher (München; Isarklinikum); A Brosin, J Lindert, M Philipp, M Gumsheimer, F Wiese (Rostock; University Hospital Rostock); L Sahan, GA Stavrou, J De Deken, M Jabal (Saarbruecken; Klinikum Saarbruecken); MW Löffler, A Königsrainer, M Quante, C Yurttas (Tuebingen; University Hospital Tuebingen).

Ghana: R Armah, NA Christian, D Daary, S Akuffo, A Twumasi, AD Andani, J Oppong, E Agbowada, J Daleku, J Ampadu, W Afedo, Z Robertson, A Obheng, DN Lee, D Ofosuhene (Accra; Greater Accra Regional Hospital); GD Brown, F Osman, FJ Eshun, C Banka, I Amankwaa, E Ametefe, G Owusu, J Nyamekye - Baidoo, O Okrah, G Birikorang, P Kumassah, J Dei-Asamoah, J Annan, C Akli-Nartey, D Alifoe, S Tsatsu, C Ansah Larbi, U Una, K Yalley, A Bediako Bowan (Accra; Korle-Bu Teaching Hospital); KB Oduro-Boateng, S Anim, H Adjei, S Dognia, A Oppong, C Markin, JL Ahale, NKA Obuobi (Accra; Pentecost Hospital); S Agana, EB Akakpo, F Galley (Ankafu; Ankafu Leprosy General Hospital); FE Gyamfi, S Segnitome, S Agordjor, D Adjei, D Kyeremeh, Y Sarpong, F Opoku Twene, CK Ntow-Boahen, SAA Atupra, V Siepaal, IN Bakaweri, AJ Tabiim, R Agyei Boakye, A Asare Twumasi (Berekum; Berekum Holy Family Hospital); R Akankoauesi Apatewen, U Kanyan Kassim, B Owusu Ansah, F Amoako (Bolgatanga; Upper East Regional Hospital); V Kudoh, K Boakye-Acheampong, B Boakye, R Kpangkpatri, MT Morna, GA Rahman, EO Ofori, L Adagrah Aniakwo, M Amoako-Boateng, D T. Enti, S Debrah, M Nortey, P Koggoh, P Mensah, MM Agyapong, T Agyen, V Etwire, Y Adofo-Asamoah, S Yussif, M Yigah (Cape-Coast; Cape Coast Teaching Hospital); B Maanikuu, F Kuubetersob B. N, D Powell, A Gbeadese, F Tierenye (Damongo; St. Anne's Hospital); EA Nacheleh, DYD Agleby, N Jiagge, R Akpaka, D Labadah, N Naabo, R Guzmán Lambert, FJ Eshun, P Ntem, E Setsoafia, N Affram, BY Hernandez Cervantes, DB Osei, K Ewool, F Nyarko, MA Ali, MA Oyortey, I Hagbevor, ME Ashong, JN Anyorigiya (Ho; Ho Teaching Hospital); J Yorke, A Lovi, EO Osei, PA Boateng, R Oppong-Amoah, K Agbedinu, C Dally, SG Brenu, F Galley, FM Agbemafoh, I Kyei, C Aboah, AY Appiah-Kubi, B Nimako, M Aikins, M Adinku, A Opoku-Agyapong, J Adjei, R Sagoe (Kumasi; Komfo-Anokye Teaching Hospital); A Gyedu, PK Boateng, S Mensah, E Frimpong-manso (Kumasi; University Hospital, KNUST); P Taah-Amoako, S Tabiri (Nsawkaw; Tain District Hospital); F Owusu, P Yeboah Owusu (Sunyani; Brong-Ahafo Regional Hospital); EMT Yenli, AS Seidu, M Dason, M Amadu, GA Adoro, M Kyereh, I Osman, J Quansah, C Doku, A Darkwa Boateng, AM Muntaka, MA Dokurugu, R Nesco, M Yahaya, EDF Konlan, A Issaka, RA Ramirez Calas, M Sheriff, M Dery, V Dassah (Tamale; Tamale Teaching Hospital); BK Seshie, F

Caiquo, M Dum, L Ackam, K Yalley, FDA Agbodo, A Baiden Amisah, D Ashitey , LD Bray, J Ofori, M Ishak (Tema; Tema General Hospital); G Ansong (Walewale; Walewale Government Hospital).

Greece: I Gogoulis, K Bekiaridou, A Mitsala, S Botaitis, C Tsalikidis, M Asimakidou, C Nikolaou, E Efremidou, C Limas, P Chloropoulou, M Aggelidou, M Pitiakoudis, P Kostoglou, G Pappas Gogos, M Karanikas (Alexandroupolis; Alexandroupolis University General Hospital); E Kapasakis, M Karakeke, A Skarpas, C Floros, K Athanassiou, E Karakeke (Amfissa; General Hospital of Amfissa); N Tasis, A Sarafi, A Plastiras, G Kavalieratos, T Tsirlis (Athens; Agios Savvas Anticancer Hospital); L Chardalias, N Memos, V Themelidi, I Papaconstantinou, T Theodosopoulos, D Politis, K Iliakopoulos, K Bramis, P Antonakis, A Skreka, D Kotsaris, N Dafnios, A Vezakis, I Contis, T Petropoulou, KC Kordeni, T Kozonis, G Fragulidis, D Massaras (Athens; Aretaieion Hospital); E Apostolopoulos, I Karatsolis, A Mourtzouni, K Avgerinos, D Kelgiorgi, K Polychronopoulos, G Kostoulas, A Tsechpenakis, A Saridaki (Athens; Athens Euroclinic); A Ioannidis, C Chouliaras, I Tierris, MK Konstantinidis (Athens; Athens Medical Center); DK Manatakis, D Balalis, N Stamos, N Tasis, V Kalles (Athens; Athens Naval and Veterans Hospital); T Sidiropoulos, M Papadoliopoulou, N Arkadopoulos, D Sampanis, P Vassiliu, E Dylja, AI Nikolaou, I Margaritis, P Kokoropoulos, V Tsaousis, S Christodoulou, E Poulis, A Chamzin (Athens; Attikon University General Hospital); S Kapiris, E Mavrodimitraki, M Sotiropoulou, N Dimitrokallis, N Papadogianni, V Vougas, M Papamichail, K Rekouna, K Pavlopoulos, N Roukounakis, A Thanasa, P Trakosari, M Christou, E Saitoglou (Athens; Evaggelismos General Hospital); C Nastos, D Dellaportas, P Lykoudis, N Garmpis, G Kouraklis (Athens; Evgenideio Hospital); P Christodoulou, G Kapogiannatos, A Nikitaras, SM Tsoiti, J Katogiritis (Athens; General Hospital Asklepieio Voulas); EC Tampaki, C Papazacharias, O Bellou (Athens; KAT Athens General Hospital); N Machairas, P Dorovinis, C Doudakmanis, D Schizas, A Syllaios, MD Keramida, P Stamopoulos, S Davakis, M Despotidis, A Panagakis, S Kykalos, F Stavratidis, M Vailas, L Karydakidis, N Kydonakis, A Loizou, KS Giannakopoulos, P Sakarellos, A Kozadinis, I Katsaros (Athens; Laiko University Hospital); C Damaskos, E Antoniou, M Mavri, I Psilopatis, S Vernadakis , I Bokos, P Paraskeva, D Vardakostas, I Gomas, A Barlas, A Smyrnis, D Prevezanos, NN Mathioudakis, I Kozadinis, A Kozadinis, P Kanavidis, N Garmpis (Athens; National and Kapodistrian University of Athens); E Spartalis, M Spartalis (Athens; Sotiria General Hospital of Thoracic Diseases); C Stefanou, S Gkogkos, L Fountoulis, I Sougkas, M Billis, A Kontokostopoulos, A Balta, T Padioti , MA Sotiriou, A Theochari (Filiates; General Hospital of Filiates); N Tsakiridis, I Tsakiridis, E Synekidou (Florina; Florina General Hospital 'Eleni Th. Dimitriou'); E Athanasopoulou, C Ntagkas, E Samara, A Katsiou (Ioannina; University Hospital of Ioannina); D Panagopoulos, A Panagopoulos, L Katsiaras (Kyparissia; General Hospital of Messinia, Hospital Unit of Kyparissia); C Kolla, L Mansour, G Koukoulis, S Zourntou, D Papageorgouli, K Bouliaris, LI Fountarlis, M Bei, AA Kalidis, X Vagena, A Gkouniaroudi, A Migdanis, A Bakalis, E Gavriil (Larissa; General Hospital of Larissa 'Koutlimpaneio and Triantafylleio'); K Koumarelas, MN Kouliou (Larissa; General University Hospital of Larissa); K Bouchagier, F Mulita, G Verras, G Skroubis, I Maroulis (Patras; General University Hospital of Patras); V Mousafeiris, A Panagidis (Patras; Karamandaneio Prefecture Children Hospital of Patras); A Papadopoulos, P Grivas, F Spanos, A Kalogeropoulou , G Zeringa, C Kourouniotis, G Rados, E Barkolias , K Zakkas, I Demiris, V Nikolaou, K Tata, G Karakaidos (Piraeus; General Hospital of

Nikaia); E Kontis, E Papamattheou, A Efstathiou, E Efstathiou, N Kopanakis, K Ntatsis, I Katsaros, P Manikis, V Tselepidis, M Kyriazi (Piraeus; Metaxa Cancer Hospital); I Siannis, N Kouzacos, V Georgilaki, N Vlachakos, S Vederaki, A Tsiaka, A Zarafidou, N Zampitis, S Tsatsos, F Stefou, F Kyramargios, M Merrakos, G Bekakos, A Marinis (Piraeus; Tzaneio General Hospital); O Ioannidis, E Anestiadou, K Zapsalis, S Simeonidis, S Bitsianis (Thessaloniki; George Papanikolaou General Hospital of Thessaloniki); M Drogouti, A Gkoutoula, A Sarakatsanos, E Efthymiou, I Chatzis (Thessaloniki; O Agios Dimitrios General Hospital); I Spyridakis, C Kaselas, M Tsopozidi, M Florou, C Demiri, V Papadopoulos, D Giakoustidis, A Giakoustidis, D Alexandrou, P Chatzikomnitsa (Thessaloniki; Papageorgiou General Hospital); SC Liapis, K Perivoliotis, C Chatzinikolaou, N Tsantikos, ZR Karampotaki, D Lytras (Volos; Achillopoyleio General Hospital of Volos).

Guatemala: M Aguilera-Arevalo, M Rodríguez-Ordoñez, DA Sosa Méndez, M Sebastián-Mendoza, TA Salazar-Lorenzana, R Herrera, DE Reyes Rodríguez, P Vásquez, S Morales, J Gomez, C García-Salas, JR Asturias Luna, J Tabora-Zepeda, S Vasquez, JR Hernández, JOM Herrera Batres, D Muñoz, E Ayala, O Coyoy-Gaitán (Guatemala City; Hospital General San Juan De Dios); JB Pellecer Cano, DA Palma Portillo, M Blanco (Guatemala City; Hospital Juan Jose Arevalo Bermejo); D Herrera, SA Villeda, O Lima Azurdia (Guatemala City; Hospital de Referencia Nacional de Enfermedades Respiratorias).

India: S Jp, M Bhat, A Raheja, I Shariff, H Anand, B R Budihal, S Kashyap, B Arya, Y M S, N Krishnappa (Bangalore, Karnataka; BGS Global Institute of Medical Sciences); M Manangi, P Anandan, S Shivashankar chikkanayakanahalli, S Kumar Venkatappa (Bangalore; Victoria Hospital); TS Mishra, P Kumar, M Gureh, MK Sethi, AA Asharaf (Bhubaneswar; All India Institute Of Medical Sciences - Bhubaneswar); Y Sakaray, S Irrinki, S Subbiah Nagaraj, S Khare, C Tandup (Chandigarh; Postgraduate Institute of Medical Education & Research, Chandigarh, India); M K B, AB Muthunayagam, A Prasath S V, P Arunachalam, P John (Coimbatore; PSG Institute of Medical Sciences and Research); S Raul, R Vakil, R Sinha, E Dvivedi, A Thomas, S Joseph, A Sharma, B Khan, D Chatterjee (DELHI; St Stephen's Hospital); R Gupta, A Khanduri, S Singh, DH Tyagi, U Daspal, N Rawal, R Varshney (Dehradun; Synergy Institute of Medical Sciences); M Luthra, R Handa, S Basu, P Chadha, R Sethi (Delhi; Holy Family Hospital); T Longkumer, KK Mishra, S Sundaramurthy, S Kumar, D Phom, J Kaippally, D Omni, L Imchen, T Alinger (Dimapur; Christian Institute of Health Sciences and Research); A Chhabra, A Kumar (Faridkot; Guru Gobind Singh Medical College & Hospital (Baba Farid University of Health Sciences)); B Kharga, M Sarda, K Bhutia (Gangtok; Sir Thutob Namgyal Memorial Hospital Sochakgang); N Sharma, MS Rodha, N Banerjee, A Baksi, S Kaur, R Chaudhary, M Lodha, SP Meena, M Badkur, I Singh, A Sinha, KJ Rathod, R Saxena, J Tk, A Vig, M Pathak, A Sukhdev Jadhav, S Nayak, T Motiwala, K Shreyas (Jodhpur; All India Institute of Medical Sciences (AIIMS), Jodhpur); SR Pathan, J Rathod, C Agarwal, K Sharma, S Pandya (Karamsad; Shree Krishna Hospital); A Anand, A Kumar, HS Pahwa, AA Sonkar, MK Agrawal, AK Pal (Lucknow; King George's Medical University); D Jain, PD Haque, V Michael, W Bhatti, J Dhiman, DRS Thind, A Bhatt, P Gupta, A Luther, S Khurana, RR Ranadive, A Suroy, H Kaur, S D A, P Shukla (Ludhiana; Christian Medical College & Hospital); NK Chaudhry, DA Hajela, P Patel, PK Arya, DD Dhawan, DR Tripathi, KK Luthra, A Kumar, DH Gupta (Ludhiana; Satguru Partap Singh

Hospital); A Mathew, C Pun, P Dummala, M Gurung (Madhepura; Madhepura Christian Hospital); P Alexander, N Aruldas (Manali; Lady Willingdon Hospital); PS Prabhu, S Payyanur Thotan, B L, B Sv (Manipal; Kasturba Medical College Hospital, Manipal); RD Sharma, R Redkar, R Nathani, S Karmarkar, A Sharma, S Singh, S Achugatla, A Bangar, DD Kulkarni, K Raghuwanshi, DN Nikam, SN Mahendra (Mumbai; Lilavati Hospital & Research Centre); B Sarang, D Belekar, K Gaikwad (Mumbai; Terna Medical College and Hospital); A John, PA Thomas, L Pramod, D Gavit, D Singh (Nandurbar; Chinchpada Christian Hospital); N Kansakar, N Gupta, N Kapur, N Narain (New Delhi; ABVIMS Dr RML Hospital); S Kulkarni (New Delhi; Army Hospital Research & Referral New Delhi); T Rashid, M Husain, F Tauheed, SV Manzoor, S Ohri (New Delhi; Hamdard Institute of Medical Sciences & Research); L Bains, P Lal, S Neogi, A Mishra, S Ahuja (New Delhi; Maulana Azad Medical College); T lahmo, M George, M Singh, P Waghchoure, A Choudhrie (Padhar; Padhar Hospital); A Kumar, M Aggarwal, V Kanna D, S Vembar, TP Singh, DS Walia, V Singh, G Kaur, A Jindal, P Dhamija (Patiala; Government Medical College Patiala); M Kumar, A Sinha, AK Jha, M Sharma, A Bhadani (Patna; All India Institute of Medical Sciences, Patna); R Abhinaya, U Kumbhar, A Jain, S Chilaka, S P (Pondicherry; Jawaharlal Institute of Postgraduate Medical Education and Research); VS Jha, A Jayapalan, Y Vashishth, G Jalal, VV Nair, C Raphael, HK Prabhakar, Z Khan (Pune; Command Hospital, Southern Command); D Dugar, D Mohanty, TDB Tridip, DR Ramchandani (Raipur; All India Institute of Medical Sciences Raipur); S Basu, N Kumar, AG Goswami, H Panga, D Mallik, A Gupta, D Rajput, R Anjum T Siddeek, P Manjunath, S Edem, F Huda, SK Singh, S Karuppusamy Krishnasamy, S Katragadda, KMR Reddy, I Ahmed, E Yhoshu, L Manoj Joshua, A Das, P Kothari (Rishikesh; All India Institute Of Medical Sciences); K Singh, SS Malhi, R Kaur, HK Cheema, M Singh, N Saini, M Gupta, A Bhatti, A Gupta, P Kaur, N Pahuja, H Kaur, S Chopra, B Sehgal, G Singh (SAS Nagar (Mohali); BR Ambedkar State Institute of Medical Sciences Mohali); DS Kshirsagar, M Kaple, G Saxena, S Dhole, A Bhargava, C Mahakalkar, S Deshpande (Sawangi (Meghe), Wardha; Acharya Vinoba Bhave Rural Hospital); R Wani, RA Dar, AA Malik, N Bhat, ZA Shah, GA Bhat (Srinagar; Sher-i-Kashmir Institute of Medical Sciences); S Kondpan, A Kutma, JA Kalyanapu (Tezpur; Baptist Christian Hospital); A Sundaram, K J B, R Krishna raj, G George, M Chisthi, H Jafarkhan, I P s, U Govindan, MKL K S D, C Narayan, G Pillai, VV Kollengode, D Jabbar (Thiruvananthapuram; Government Medical College Thiruvananthapuram); T Tony V, A Nair, A Kavalakat, A Moncy, A Johnson, A Joseph, A Appukuttan, A J, AN Oommen, S Francis, S M, N Srinivas, M Narayanan (Thrissur; Jubilee Mission Medical College & Research Institute); T Devabalan Koil, MR Jesudason, Y Myla, ASP Dhinakar, A Tirkey, B Roopavathana. S, R Mittal, S Surendran, N Paul Ambrose, D Joshiba, P Trinity, R Raghunath, NP Paul Sigamony, PY George, R Philip Sridhar, S Chase, SJ Arthur (Vellore; Christian Medical College & Hospital).

Iran, Islamic Rep.: N Yousefzadeh Kandevari (Bastak; Farabi hospital); M Pourfridoni, H Askarpour, H Mohammadi sardoo, AA Kheirkhah Vakilabad, M Ali-Hassanzadeh (Jiroft; Imam Khomeini Hospital).

Iraq: R Raheem Attallah Al_obaidy (Anbar; Heet General hospital); Z Alkhuzai, S Salim, FRH Hassooni (Najaf; Al Batool private hospital); Y Zwain, HMA Oneizah, S Razaq, M Razaq, HHZ .zaini (Najaf; Al-Najaf Al-Ashraf Teaching Hospital).

Ireland: E Colton, KE Oderoha, A Gill, S Ramjit, F Ghazali, K Griffin, R Ahmed, C Weadick, S Nair, V Sharma, C Donohoe (Dublin; St James's Hospital); R Tummon, B Maguire, S Wrenn, R Habib, P Owens (Kerry; University Hospital Kerry).

Israel: A Shweiki, G Szydlo Shein, G Almogy, Y Mintz, A Pikarsky, R Elazary, O Cohen-Arazi, JA Demma, H Mahajna, B Helou, Y Fishman, G Marom (Jerusalem; Hadassah Medical Center).

Italy: M De Prizio, K Kröning, R Sulce, LM Fatucchi, F Tofani, V Mariottini, R Malatesti, M Scricciolo, GA Pellicano', A D'Ignazio, A Mazzoni, A Biancafarina, M Angelini, V Borgogni (Arezzo; Ospedale San Donato USL Toscana Sud Est); L Rossi, G Munzi, G Tarantino, M Castrovillari, A Serao, JR Casella Mariolo, G Del Corpo, A Natili, A Iodice (Ariccia; Ospedale dei Castelli (N.O.C.)); S Gargiulo, B Esposito, M Pannullo, L Bracciano, E Marra, A Alberico (Aversa; San Giuseppe Moscati); A Gori, S Cardelli, G Dajti, C Larotonda, IS Russo (Bologna; IRCCS Azienda Ospedaliero-Universitaria di Bologna); J Andreuccetti, S Molfino, D Alberti, G Pignata, G Emiliani, G Boroni, M Ruffoli, M Manfredini, L Sequi, G Zanni (Brescia; ASST Spedali Civili, Ospedale di Brescia); E Locci, M Podda, V Murzi, C Piras, A Carta, A Pisanu, T Pilia, P Marongiu, A Saba, M Pisano, F Campus, E Gessa, E Silanos, A Lai, F Frongia, F Corronca, S Montisci (Cagliari; Cagliari University Hospital); F Cappellacci, C Soddu, GL Canu, F Medas, PG Calò, S Puddu, M Biancu, M Abbas, F Casti (Cagliari; Chirurgia Generale e Polispecialistica, Cagliari University Ospital 'Duilio Casula'); B Demurtas, A Deserra, F D'Agostino, C Margiani (Cagliari; Santissima Trinità - ATS Sardegna); L Laface, M Casati, M Mariani, S Guarriello, A Balconi (Carate Brianza (MB); Ospedale Vittorio Emanuele III - Carate Brianza); B Scotto, N Laquatra, R Ruccella, G De Angeli (Carpi; Ramazzini); G Gravante, A Chiappini, R Lopatriello, G Mammolo (Casarano; Francesco Ferrari Hospital); C Distefano, G Riccioli, R Granata, M Veroux, DC Centonze, S Costa, R Gioco, D Zerbo, A Licciardello, L Stella (Catania; Azienda Ospedaliero-Universitaria Policlinico San Marco); L Rende, M Osso, D Paglione, F Pata (Cosenza; Azienda Ospedaliera di Cosenza); D Sasia, G Giraudo, D Ribero, S Alberti, V Schirinzi (Cuneo; Santa Croce e Carle Hospital, Cuneo); P Belotti, L Taglietti (Esine; ASST Valcamonica Ospedale di Esine); V Giordano, A Pesce, CV Feo, MC Pignanelli, S Severi (Ferrara; Azienda Unità Sanitaria Locale di Ferrara); F Cammelli, F Natali, M Scheiterle, G Maltinti, L Fortuna, F Coratti, A Manetti, J Martellucci, E Monati (Firenze; Azienda Ospedaliera Universitaria Careggi); L Gabellini, R Fratarcangeli, A Damigella, E Adinolfi, A Anastasi (Firenze; Ospedale San Giovanni di Dio); M Montagna, A Giuliani, G Procaccini, N Tartaglia, F Vovola, D Merlicco, S Schirone, ST Massa, G Pavone, M Pacilli, F Maffei, A Gerundo (Foggia; Ospedali Riuniti Azienda Ospedaliera Universitaria); D Di Pietrantonio, S Quartarone, L Solaini, G Ercolani, L Ragazzini, V Zucchini (Forlì; Morgagni-Pierantoni); M Cammelli, D Scotto Di Carlo, FAN Marin (Garbagnate Milanese; ASST Rhodense - Ospedale di Garbagnate Milanese); A Azzinnaro, A Razzore, A Petrunaro, E Mina, B Sperotto (Genoa; E.O. Ospedali Galliera); G Carganico, D Pertile, D Soriero (Genoa; IRCCS Ospedale Policlinico San Martino); R Diaz, A Luzzi, S Carrabetta, C Meola, D Caruso, F Floris, P Grondona, E Romairone, S Marzorati, F Ré, C Righetti, A Viacava, L Epis (Genoa; Ospedale Villa Scassi); R Sampietro, D Gobatti, C Zandonella (Gravedona ed Uniti; Ospedale Moriggia Pelascini); VP Dinuzzi, U Rivolta, S Luciano, GMF Marini, L Scaravilli (Magenta; Ospedale 'G.Fornaroli', ASST-OVEST Milanese); R Magarini, G Saletta, AC Sironi, G Grava, M Mercurio (Melzo; Ospedale Santa Maria

delle Stelle, ASST Melegnano Martesana); D Zulian, A Izzo, M Gritti, S Giudici, E Desiato (Milan; Humanitas Research Hospital); M Molteni, L Ottaviani (Milan; IRCCS San Raffaele Scientific Institute, Milan); G Grande, E Mazzotta (Milan; Ospedale Fatebenefratelli e Oftalmico); S Grimaldi (Milan; San Carlo Borromeo); F Brucchi, F Ferraina, S Lauricella (Milan; Sesto San Giovanni Hospital); E Di Marco (Modica; Ospedale Maggiore); LC Nespoli, G De Carlo, EA Baccalini, D Palmisano, A Scacchi, N Tamini, L Ripamonti, M Rennis, C Vitiello, A Davolio, L Degrade, P Masseria, V Brocco, P Chiacchio, M Ceresoli, E Signaroli, A Finocchio, C Fumagalli, M Binda (Monza; Fondazione IRCCS San Gerardo dei Tintori Monza, Scuola di Medicina e Chirurgia, Università Milano Bicocca); M Milone, M Manigrasso, S Vertaldi, A D'Amore, GD De Palma, A Mareello, L Fedele, C Sorrentino, D Pignatelli, G Luglio, FP Tropeano, M Cricri, A Miele, G Aprea, G Palomba, M Capuano, R Basile, G Sorrentino (Naples; Federico II University of Naples); D Rega, A Ottaiano, V Granata, A Belli, F Izzo (Naples; Istituto Nazionale Tumori Fondazione, Pascale IRCCS); G Pellino, D Massaro, V Mosca, F Selvaggi (Naples; Primo Policlinico di Napoli); G Bellio, L Rubin, N De Santis, N Schiavon, A Zerbinati, S Corso, C Cecconi (Padova; Piove di Sacco Hospital); P Venturelli, G Cocorullo, G Carollo, R Tutino, A Bonelli, G Salamone, MP Proclamà, R Guercio, L Licari, N Finocchiaro, G Graziano, G Orlando, G Guercio, M Marcianò, G Galatioto, F Vassallo (Palermo; Policlinico Universitario Paolo Giaccone); G Palmieri, F Banchini (Piacenza; G. Da Saliceto); G Di Franco, F Porcelli, N Furbetta, A Comandatore, S Guadagni, M Palmeri (Pisa; Azienda Ospedaliero Universitaria Pisana); P Ubiali, F Maffeis, J Velkoski (Pordenone; Azienda Sanitaria Friuli Occidentale (AS FO)); M Giuffrida, GE Nita (Reggio Emilia; Azienda Unità Sanitaria Locale - IRCCS di Reggio Emilia); C Marafante, M Garino, SL Birolo, M Dugo, M Pisano, A Borello, R Barone, LD Bonomo, MV Facchino, M Caccetta, E Moggia, MR D'Anna, C Mosca, S Mungo, A Masciandaro (Rivoli; Ospedale degli Infermi di Rivoli); F Tirelli, I Neri, M Aulicino, C Vacca (Rome; Fondazione Policlinico Universitario Agostino Gemelli); M Campanelli, M Grande, L Siragusa, G Sica (Rome; Policlinico Tor Vergata Hospital, Rome); A Mingoli, G Brachini, B Cirillo, S Meneghini, S Giovampietro, G Duranti, F Ciccarone, L Simonelli, I Clementi, B Binda, GB Fonsi, E Spalice, E Cianci, MI Bellini, G Sgarzini, S Sorrenti, E Lori, P Palumbo, D Pironi (Rome; Policlinico Umberto I Sapienza University of Rome); A Amendola, C De Martino, E Bisogno (Salerno; San Giovanni di Dio e Ruggi d'Aragona); S Di Saverio, A Morello, L Lely, I Merlini, G Travaglini, S Sabbatini, M Zambon (San Benedetto del Tronto; Madonna del Soccorso Hospital); D Giullitti, L Barni, GE Poto (Sansepolcro; Valtiberina); D Fusario, L Resca, L Carbone, A Francia, GE Poto, AL Pesce, O Carpineto Samorani, F Roviello, A Ongaro, SA Piccioni, M Gambelli, L Catozzi, M Gjoka, A Bartalini Cinughi de Pazzi, NN Leonelle Lore, G Grassi, F Manasci, V Ricchiuti (Siena; Azienda Ospedaliero Universitaria Senese); E Basile, A Spaziani (Spoleto (PG); San Matteo degli Infermi); V Silvestri, M Favoriti, P Favoriti (Sulmona; SS Annunziata); S Novello, M Piccino, R Baldan, U Grossi, G Zanus, F Scolari, E De Leo, M Brizzolari, A Brun-Peressut, I Hoxhaj, M Scopelliti (Treviso; Ospedale Ca' Foncello - Università di Padova (DISCOG)); LB Lo Piccolo, E Montanari, D Cianflocca, A Marano, S Galati, SL Gamba, F Velluti, A Caltagirone, M Giuliano, E Potenza, B De Zolt Ponte, D Visconti, VU De Donato, L Capello, S Chaifouroosh Mamagany, C Celano (Turin; Città della Salute e della Scienza); E Donnarumma, C Saviello, R Scola (Vallo Della Lucania; Casa di Cura Prof. Dott. Luigi Cobellis); S Megna, M Berselli, N Palamara, L Liepa, E Ferri (Varese Lombardy; University of

Insubria, Ospedale di Circolo e Fondazione Macchi (Varese)); V Gentilino, M Mogiatti, G Farris, N Pasqua (Varese; Filippo Del Ponte Hospital, University of Insubria); A Iacomino (Venezia; Ospedale Civile - Santi Giovanni e Paolo); I Mondì, C Da Lio, F Sulo, E Ciccioli, D Verdi (Venice; Mirano Hospital); M Martorana, M Filardo, L Schiavone (Voghera; Ospedale Civile di Voghera); I Conversano, M Cappiello, G Scialandrone, N Petrarota, R Tumolo (andria; Lorenzo Bonomo); M Angelucci, S Valeri, G Pascarella, A Strumia, R Alloni (rome; policlinico universitario campus bio medico of rome); D Muschitiello, V Morinelli, L Bonello, SG Intini, S Moschella (udine; santa maria della misericordia).

Japan: H Kato, A Horiguchi, D Koike, Y Asano (Aichi; Fujita Health University Bantane Hospital).

Jordan: S Alananzeh, S Al Momani, M Tanashat, O Altobaishat, S Alsmadi, N Al RABADI, L Sweidan (Ajloun; Al Iman Hospital); Z Alnajjar, G Alsheikh, N Mosleh, H Alzuhd, R Alkhatib, H Al-Abdallat, M Aljarawn, T Aloqaili, I Nadi, A Abdllah, O Al-Fahel, R Khalil, M Said, A Qasem, H Al-Fahel, M Hijazi, R Rabah, A Alaqtash, DB Badwan (Amman; Al-Basheer Hospital); Y Alawneh (Amman; Ibn Al Haitham Hospital); B Alrayes, M Salah, M Al-Qannas, H Abu Obead, I Alnimer, Y Alawneh, M Almaletti (Amman; Islamic Hospital); A Khamees, R Yousef Yassin, A Alsheikh, K Al-Shami, E Abu Siam, O Sarhan, KA Sawaftah, MAM Sawaftah, MA Sawaftah, M Sabri Massadi, Z Al-sheikh ali, N Raiq, O Ibrahim, J Al Karmi, M Diab, I Aburumman , AA Altawaiha , M Hasan, A AlZu'bi, L Yasin, B Yacoub (Amman; Jordan University Hospital); R Abu Salah, M Alqedrh , S Abu khousa, MEH Albanna (Amman; Marka Specialty Hospital); R Hussam Yacoub Hattar, D Samardali, R Refaie, L Hijazein , S Hammad, M Barbarawi, S Mamduh, J Al Daradkah, S Samardali , T Alshawabkeh (Amman; Prince Hamza hospital); M Mahafdah, Q Sabbah, SAM Ba-Shammakh, A Al Hammoud, M Bani hani, M Tabaza, H Malkawi, H Haj Freej, D Kasasbeh, B Dweik, K Ayyoub, OR Mahafdah (Ar Ramtha; King Abdullah University Hospital/ Jordan University of Science and Technology); R Hiary, I Shehadeh, L Dyab, R Daradkeh, R Raddad, R Alzu'bi, T Alhaj Hasan , B Alzoubi, M Alsharayri, M Nofal, S Ellouzy, M Al-Masri, S Fakhouri, N Absy, R Suleiman, O Mansour, JS Hadidi (As-Salt; Al Hussain New Salt Hospital); A Al-fandi (Irbid; Ar Ramtha Governmental Hospital); M Al-Fraijat, N Rabai, A Al-Zubeidy, R Abd Elkareem, S Bani Amer , T Alhusban, S AL-Doghme, R Abd Alkareem, R Damseh, M Alshami , T Majed, S Al Sharie, M Araydah, R Jaba'Teh, F Haddad, O Almomani, L M Mheidat, R Haddad, S Bataineh, S Ababneh (Irbid; Princess Basma Hospital).

Kazakhstan: M Kulimbet, N Maulenov, N Lakhanov, M Ramazanov, A Kiyabayev (Almaty; City Clinical Hospital No.7, Asfendiyarov Kazakh National Medical University); D Amangaliyev, A Shamsutdinova, A Polatbekov (Almaty; JSC 'Central Clinical Hospital', Asfendiyarov Kazakh National Medical University).

Kenya: R Parker, E Irungu, A Fadipe, F Ondago, G Waiyaki (Bomet; Tenwek Hospital).

Lebanon: A Khoneisser, A Kachi, B Abboud (Beirut; Hopital Libanais Geitaoui); M Chaccour, G Bechara, N Eshak, R Hleyhel, M Barakat (Jbail ; Maritime Hospital).

Libya: N Lindi, BH Hameed, A Abaidalla, N Mosbah, M Saleh khatib (Albayda; Albayda Medical Center); MAM Elghriani, A Ali, A Fathi, S Qwyder, M Saleh (Benghazi; Al-jalaa Teaching/Trauma Hospital); M Alshamikh, A Alhammali, A Aldurssi, A Gusibat, M Suleman, S Alashhab, M Denini, F Alowjaly, MM Almihashhish, F Elkhafeefi , H Altawati (Benghazi; Benghazi Children's Hospital); S

Elfallah, F Benghalbon, R Michael , M Abosedra, A Ahmayda, H Mftah (Benghazi; Benghazi Medical Center); M Bohlala, M Muragi, S Alneihuom (Darna; Al-Wahda Hospital); A Alkaseek, A Alshiteewi, H Shames, H Bileid Bakeer (Gharyan; Gharyan Central Hospital); N Albahloul, M Abudabbous, A Belkhair, A Abdelmalik, M Assalhi, M Altajouri, A Alailesh , A Alshukre (Misurata; Misurata Central Hospital); B Alazabi, M Alazabi, G Birqeeq, AAY Almugaddami, A Egdeer (Nalut; Nalut Central Hospital); H Embarek, M Bilfaqirah (Sebha; Al-Majd Clinic); M Abdelkabir, S Abdeewi , A Abdalhadi, M Benghazi (Sebha; Aseel Alghad Clinic); H Idheiraj, M Yahmad, M Alfaid, M Abdu, E Abdu, M Khalifa, G Matroud, A Amaigl , H Aldare, E Ali, K Shwail, K Abdulrahman (Sebha; Sabha Medical Center); A Bouhuwaish, A Emran, A Abdraba (Tobruk; Tobruk Medical Center); AE Elzoubi, A Belaid (Tripoli; Alkhalil hospital); A Alragheai, D Omar, S Magrhi, H Farhat, S Alsuwiyah , S Abrayik, S Bensalem, B Algettawi (Tripoli; Metiga Hospital); S Egreara, TMA Abdulmola, I Kandil (Tripoli; Sabratha teaching hospital); F Alshreef, F Elhabishi , M Alsori, L Shawesh , S Timmalah, M Alnuwayli, A Alhamadi, K Ahmed Ibrahim, S Abdullateef, R Altayargh, A Essamei, S Altoume, A Haidar, M Khalil, A Abdulnabi, S Mohammed , A Ghummied (Tripoli; Tripoli Medical Center/ Tripoli University Hospital); E Younes, S Elfurdag , A Ali, S Ashini, M Edeeb, Z Al-azher El-hamel, M Akkawe , N AIWAER, A Khair Etareig (Zawia; Zawia Teaching Hospital); B Allbakosh, H Abusnina, A Awidan, L Alokshi, M Iqreewi , M Almahjoub , H Altounsi, M Almaqrahi (Zliten; Zliten Teaching Hospital).

Lithuania: E Dainius, S Bradulskis, E Margelis, A Mačiulaitytė, A Subocius, A Parseliunas, E Kubiliute, D Zuikyte, J Kutkevičius, J Vaitekūnas (Kaunas; LUHS Kaunas Hospital); L Venclauskas, K Jasaitis, M Jokubauskas, Z Dauksa (Kaunas; Lithuanian University of Health Sciences Kaunas Clinics); A Gulla, E Daukšaitė (Vilnius; Vilnius University Hospital).

Madagascar: MJ Rakotonaivo, CF Rahantasoa Finaritra, JB Razafindrahita, YM Razafimandimby , A Rakotondrainibe (Antananarivo; Joseph Ravoahangy Andrianavalona Hospital).

Malaysia: AD Zakaria, MIS Ismail (Kelantan; Hospital Universiti Sains Malaysia); R Noor, M Che yaacob, SF Moh Pauzi, Z Chin (Kota Bharu; Hospital Raja Perempuan Zainab II); K Voon, JH Fu, JH Lim, SA Theivendran, NN Ramli (Kuching, Sarawak; Sarawak General Hospital); M Fitri, MA Yunus, AN Ramly, A Md Yunus, AAA Anuar (Malacca; Hospital Jasin); NS Abd Ghani, AAH Ahmad Zaidi, F Ashraf, M Mahadi, AA Abdul Rahim (Serdang; Hospital Pengajar Universiti Putra Malaysia (HPUPM)).

Mali: OAA Dicko, A Dembele, H Dolo, D Kone (Ségou; District Hospital of Tominian).

Malta: C Cini, A Sultana, M Farrugia, J Schembri Higgans, S Bowman, J Psaila, P Andrejevic, S Brincat, K Muscat, M Sammut, R Abela, M Zammit Vincenti, L Casingena, J Galea, M Portelli, M Sammut, N Spiteri, K Iles, R Cachia, D Hili (Msida; Mater Dei Hospital).

Mexico: TR Ibarra-Hurtado (Guadalajara; Antiguo Hospital Civil de Guadalajara); LA Flores Chávez, JA Flores Prado, K Jasso García, NE López Bernal, EV Romo Ascencio, LM Flores Chávez, MP Mellado Tellez, SA Ibarra Camargo, G Delgado Hernandez, JA Guzman Barba, LA Rea Bocanegra, M Tello Jimenez, JA Tavares Ortega, E Gómez Mejía, I Esparza Estrada, AA Salinas Barragan, JA Jimenez Flores, SJ Vázquez-Sánchez, J Gonzalez Garcia , ZM Correa López, FJ Barbosa Camacho (Guadalajara; Clínica de Especialidades más Centro de Cirugía Simplificada); CM Nuño-Guzmán, AM Nava Franco, JF Martinez Martin del Campo , JJ Ulloa Robles , L Bravo, ME Gonzalez-Gonzalez, FD Romo Rosales, TR Ibarra-Hurtado, LG Peña Balboa, C Yanowsky-Gonzalez, R Santana Ortiz, SA

Trujillo Ponce, J Orozco-Perez, MDC Gonzalez, JE Gonzalez Aboytes, J Pizarro Lozano , JE Orozco Navarro, F Ibañez Ortiz , O Montañó Angeles, M Calderon, F Diaz , M Lazo Ramírez , JA Aguilar (Guadalajara; Hospital Civil Fray Antonio Alcalde); A Gonzalez Ojeda, C Fuentes Orozco, JM Chejfec-Ciociano, JM Carranza Rosales, MA Sánchez Audelo, CI Lupercio Figueroa, KV Ascencio Diaz, CE Gutierrez de la Rosa, F Mercado Sanchez, FY González Ponce, R Mares País, C González Baez, LÁ Pelayo Orozco, NG Barrera Lopez, A Ramírez Beas, MÁ Zaragoza Mendieta, MF Zarate casas , SL Trejo Ramos, P Salas Núñez, JA Gutiérrez Gómez (Guadalajara; Hospital de Especialidades, CMNO-IMSS); G Ambriz González, I Cabrera, HB Moya- Ambriz , FJ Silva Rivera, EM Torres De Anda, A Hernández, FJ León Frutos, VE Armenta Tapia, M Nieto Galvan, JM Alvarez Hernandez (Guadalajara; UMAE Hospital de Pediatría Centro Medico Nacional de Occidentales IMSS); AI Sánchez-Terán, N Muñoz Montes, AN Fuertes Muñoz, RL Smolinski kurek (León; Hospital Regional e Alta Especialidad del Bajío); K Bozada-Gutiérrez , A Nuñez Venzor, A Zubillaga-Mares, I Serrano (Mexico City; Hospital General Dr. Manuel Gea González); C Moreno-Licea, S Anaya Sanchez, A Trigos Díaz, CJ Pérez - Padrón, EY García-Villegas., RH Perez-Soto (Mexico City; Instituto Nacional de Ciencias Médicas y Nutrición 'Salvador Zubirán'); MJ Rueda Medécigo, A Leon - del- Angel, HDJ Pérez Baca, C Chavarría Noya , L Castro (Pachuca; Sociedad Española de Beneficencia); D Herappe , MT Barrio Renteria (Querétaro; Hospital de especialidades del niño y la mujer); A Ramos-De la Medina, L Martinez, II Durán Sánchez , DS Gonzalez , MJ Martínez (Veracruz; Hospital Español Veracruz).

Morocco: M Berrakkouch, F Hourri, A Benmansour, R Ait Ben Addi , M Melouane, A Tariq, O Boujidi, I Zerrouq , M Katif, S Amahmid, S Errami, S Jamil, O Nouhail, H Essalim, A Nidali, N Ouachou (Marrakech; Centre Hospitalier Universitaire Mohammed VI, Marrakech); A Aboumedian , S Kessab, N Lahnaoui, O Arsalan, S Kassad, A Hrora (Rabat; Centre Hospitalier Universitaire Ibn Sina Rabat).

Namibia: JT Abebrese, M Van der Colf, FW Quayson , P Shimbulu, P Nambala (Windhoek; Windhoek Central Academic Hospital).

New Zealand: S Rennie, A Herewini (Masterton; Wairarapa Hospital); S Kosna, A Olsen, J Gillingham, R Campbell, A Lin (Wellington; Wellington Regional Hospital); T Uiyapat, S Beavis, D Bardsley, M Lill (Whanganui; Whanganui Hospital); J McNab-Hand, O Ray, C Harmston (Whangarei; Whangarei Hospital).

Nigeria: A Adeyeye, A Akinmade, E Afeikhena, AI Okunlola (Ado Ekiti; Afe Babalola University Multi-System Hospital); D Idowu, J Olorunfunmi, A Olabode (Ado-Ekiti; Ekiti State University Teaching Hospital); N Oloko, KJ Bwala, A Ningi (Bauchi; Abubakar Tafawa Balewa University Teaching Hospital Bauchi); P Agbonrofo, D Osifo , P Idjerhe , BO Izedomi, O Omoike, O Irowa, S Ideh, J Enaholo, C Agbonrofo , O Emuze , PV Odigie, M Ediale, RA Eghonghon, M Edena, A Ekpetai , M Momoh, C Osime, O Osagie, A Arekhandia, M Ibadin (Benin City; University of Benin Teaching Hospital); T AbdulRahman, R Ediru, J Abutu John, O Owolanke (Bida; Federal Medical Centre Bida); U Ezomike, N Agugua-Obiany, J Ede, S Aliozor, EI Nwangwu, C Ilo, C Amah, L Onyebulu, I Ugwueke , I Obiany, C Onwuzu, U Dilibe , I Orji, V Enemuo, N Celestine , N Ekwo (Enugu; University of Nigeria Teaching Hospital); SA Sani, S Olori, I Pius Ogolekwu , O Attawodi , R Hauwa SANI, P Chimezie Andrew (Gwagwalada; University of Abuja Teaching Hospital); A Ishola, O

Ayandipo, N Akinbami, A Fakoya, TA Lawal, V Osoka, H Ogundipe (Ibadan; University College Hospital); A Ademuyiwa, F Alakaloko, O Oluseye , N Duru, M Ojo, T Olobatoke , AO Lawal, C Nwanmah , O Alaba, R Eloka, C Bode, O Elebute, J Seyi-Olajide, K Onyekachi, O Balogun, O Christianah, L Omomeji , F Akinwande, A Damola-Okesiji, J Okei (Idi Araba; Lagos University Teaching Hospital); H Abiyere, O Fatudimu, B Mustapha, O Babatunde, AI Okunlola (Ido Ekiti; Federal Teaching Hospital, Ido Ekiti); OM Williams, O Faboya, C Ónyeka, F Oni, K Shodunke, G Eke, M Abdulsalam , O Oso, A Ayodele, M Okechukwu (Ikeja; Lagos State University Teaching Hospital); C Adumah, A Talabi, O Oyinloye, O Olajide , V Agbakwuru, A Aderounmu, A Agbaje, YL Balogun, MO Ameen, DO Komolafe, O Olasehinde, O Ajiboye, M Fagbayimu, G Aduroja, O Fasoro, A Adisa, AM Olugbami, H Oyinlola, E Adebunmi (Ile-Ife; Obafemi Awolowo University Teaching Hospitals Complex); T Mohammed, A Lawal, F Bello, P Adebayo, O Salako, A Akinkuolie, MA Adetoyi, A Akeem Aderogba, T Oyeyemi , O Ojo (Ilesa; Obafemi Awolowo University Teaching Hospitals Complex Wesley Guild Hospital Unit); E Osaze, H Ekwuazi , I Ogundele, P Elemile, A Ayeni, I Okoro , C Onuoha (Ilishan-Remo; Babcock University Teaching Hospital); M Mobolaji-Ojibara , J Mohammad mohammad (Ilorin; General Hospital); NT Abdulraheem, A Jimoh , A Lawal, OK Fasiku (Ilorin; University of Ilorin Teaching Hospital); B Aminu , S Kache, G Yohanna Abrak (Kaduna; Barau Dikko Teaching Hospital); AA Sheshe, L Anyanwu, AB Muhammad, A Abubakar Abdulkarim , TN Nagwamutse, IU Garzali, S Muhammad, C Nwachukwu, IE Suleiman, M Abdullahi , SA Aji, A Dahiru, LB Abdullahi, SA Yunusa , A Yahaya, M Bello, I Wasiu, U Mohammed Bello, B Yunusa, N Umar (Kano; Aminu Kano Teaching Hospital); RE Enejo, N Nwafulume, A Oke, J Taiwo (Lokoja; Federal Teaching Hospital Lokoja); OH Ekwunife, OA Egwuonwu, OA Okoye, N Nwanne, C Ugwunne, J Ugwu, U Ezidiegwu, CD Nwosu, K Oluchukwu , V Modekwe, C Uche, EA Obiesie, C Osuigwe (Nnewi; Nnamdi Azikiwe University Teaching Hospital); OH Ekwunife, J Aseme, U Edith, J Ezeh, J Nnoli, H Willy-Chidire, D Chimkaomasiri (Onitsha; Holy Rosary Specialist Hospital); A Ojewuyi (Osogbo; UNIOSUN Teaching Hospital); I Ogundele, A Adekoya, L Amosu, A Oyedele, A Ayoade, BA Ayoade, A Asekun , A Ajayi, O Popoola, M Yinusa, AAA Oyelekan, O Oluyemi, C Nwosu, M Okudero, M Agunloye, A Ojo, C Nwokoro, I Babajimi-Joseph , A Williams, S Ogunlade (Sagamu; Olabisi Onabanjo University Teaching Hospital); M Daniyan, TT Sholadoye, M Bashir, N Oyelowo, SE Nwabuoku , A Yakubu, O Ogunsua, M Abubakar , MA Tolani, MS Aliyu (Zaria; Ahmadu Bello University Teaching Hospital).

North Macedonia: T Risteski, V Naunova, L Jovcheski (Skopje; University Clinic for Pediatric Surgery).

Oman: PW Haque, A Albatanony, A Fajardo, M Omer, DP Talreja, AM Elsayed , H Habiba, H Al-Aamri (Ibra; Ibra Hospital); F Ali, S Alsibai, B Dawud, J Albalushi , AM Al Balushi, A Alzadjali, T Al Barhi, A Alkharusi, F Alfarsi, K Al Hinai, H Al Qadhi, R Al Shehhi, H Al Miskry, O Al Hamdani, O Al Alyani, B Rehmani , D Ghosh, M Al-Attaqchi, Z Al Balushi (Muscat; Sultan Qaboos University Hospital); M Al Hinai, S Alwardi, L AlRiyami, A ALJamoudi, M Masaaod (Nizwa; Nizwa Hospital).

Pakistan: A Raza, A Hai, S Ahmed, DS Maqsood, S Chaudhary, M Farooq, M Tayyab, Z Qureshi, M Ali, S Maqbool, U Abdullah, M Aziz, A Irshad, S Said, A Zafar, U Akram, I Sadiq, A Abbas, M Siddique , I Shahbaz (Islamabad; Dr Akbar Niazi Teaching Hospital); SH Waqar, M Raheem, F Akhtar, DM

Mehmood, DN Mahmood (Islamabad; The Pakistan Institute of Medical Sciences); M Shahid, A Ali, M Ahmed, M Mansoor Iqbal, Y Lakdawala, S Otho, M Khalid, M Masood, R Kumar, S Jabeen (Karachi; PAF Faisal Hospital); S Altaf (Karachi; Patel Hospital); M Abdullah, G Shamsi, I Ahmed, N Lodhi, G Awais, L Rai (Karachi; The Indus Hospital); S Khattak, A Janjua, A Liaquat, M Saleem (Lahore; Bahria International Hospital, Bahria Orchard); MN Rafique, MM Bin Khalid, H Ahmad, MS Khalid, MH Sadiq, A Hashmi, HH Shahid, MA Sadiq, MH Chishti, M Usama, M Kashif, AM Choudhary, H Basharat, K Khalid, MA Haider, MA Bashir, H Sabir, MF Tarar, M Usama (Lahore; Services Hospital Lahore); MB Mirza, WU Rehman, W Tahir, R Khalid, CE Azmat (Lahore; The Children's Hospital & The Institute of Child Health Lahore); H Qayum, H Irfan Khan, A Mustafa (Peshawar; Mercy Teaching Hospital); HW Bhatti, MR Farooqui, F Rauf, NA Malik (Rawalpindi; Benazir Bhutto Hospital); M Usman Malik, S Hayat, D Riazhussain (Sargodha; District Headquarter & Teaching Hospital - Sargodha).

Palestine: J Najajra, N Al-Hroub, M Abu Daoud, R Jubran, M Srour, M Taamreh, A Alsalahat, H Masalma (Bethlehem, West Bank; Beit Jala Governmental Hospital (Al Hussein)); A Alwali, A Alwali, L Mohammed, M Al zebda, S Mahdi, A Shaheen, G Alranyes, L Tafesh, T M. Abubasheer, M Abu Jayyab, H Jaber, M Abuwarda, B M. J. Alhaj, T Aldirawi, F Mahmoud, M Ali, A Albhaisi, M Obaid, WJN Almadhoun, A Alroobi (Gaza; Al-Shifa Hospital); M Abo Abdo, A Abuthaher, A AbuNemer, M Abu Al Amrain, I Nasser, A Awad, A AlAgha, R Madi, D AbuNemer (Gaza; Nasser Hospital); N Kullab, H Elhallaq, A Abu Tair (Gaza; Palestine Red Crescent Society - Al-amal Hospital); H Ayesh (Hebron, West Bank; Al-Ahli Hospital); H Abu-Arish, A Zamareh, M Ahmoud, M H. Oweidat, I AlJada, M Anati, S Halabi, W Alhroub, A Abuhammad, S M. Udwan, H Yaghmour (Hebron, West Bank; Governmental Hebron Hospital-Alia); D Houmran, NA Awwad, M Abed, I HajMohammed, A Hewari, A Alqerem, E Zidan, H Abbadi, S Abed (Jenin, West Bank; The Martyr Dr. Khalil Sulaiman Hospital (Jenin Governmental Hospital)); M Shakhshir, M HajHamad, M Saifi, S Abuzahra, A Khouli, Z Shabello, Z Khraim, S Ismail, MF Dwikat (Nablus, West Bank; Rafidia Hospital); R Bassam, A Sabbah, A Gharib, R Alzughayyar, R Issa, A Abuhantash, O Matar, Y A. Omar, O Khalil, A Awwad (Tulkarm, West Bank; Martyr Thabet Thabet Governmental Hospital).

Paraguay: A Rodriguez Gonzalez, ED Sosa Ferreira, R Ferreira Acosta, MN Martínez Bareiro, JE Giubi Bobeda, A Franco (Asuncion; Hospital de Clínicas, II Cátedra de Clínica Quirúrgica, Universidad Nacional de Asunción).

Peru: A Walfor (Huánuco; Hospital Regional Hermilio Valdizán Medrano); L Poggi, L Poggi, L Fuentes Rivera Lau, MA Moreno Gonzales, F Camacho, O Ibarra, G Arredondo (Lima; British American Hospital); K Nieto Yrigoin, D Chavez Fernandez, DC Juan Carlos (Lima; Clinica Internacional); G Mendiola, A Salazar, R Casma Bustamante (Lima; Hospital Santa Rosa de Lima); G Borda-Luque, Y Carpio Colmenares, MR Li Valencia, F Palomino Escalante, K Quispe de la Roca (Lima; SANNA - Clínica El Golf); MM Caramantin Obando (Paíta; Hospital I Miguel Cruzado Vera EsSalud); R Polo, V Serna-Alarcon (Piura; Jose Cayetano Heredia III Regional Hospital).

Poland: R Mazurkiewicz, M Kołomańska, PJ Milewski (Kielce; Wojewódzki Szpital Zespólny w Kielcach); M Kisielewski, T Stefura, K Richter, W Wysocki, N Kłós, W Jabłoński, I Alsoubie, B Żaczek, T Wojewoda, J Bolanowski (Krakow; 5th Military Clinical Hospital); Z Orzeszko, M Wikar, R

Solecki, B Markowska, M Szura, T Gach (Krakow; Brothers Hospitallers Hospital); M Matyja, B Habrat, W Serebnicki (Krakow; Jagiellonian University Medical College); Z Lorenc, M Święch, M Mietła, W Krawczyk, M Nycz (Sosnowiec; Wojewódzki Szpital Specjalistyczny nr 5 im. Św Barbary); K Urbańska, K Komorowska, P Kowalewski, M Walędziak (Warsaw; Military Institute Of Medicine); JK Zajac, M Zawadzki (Wroclaw; Regional Specialist Hospital in Wroclaw); M Kusiński, M Pryt, F Brzeszczyński, H Dąbrowski, M Redynk (Łódź; Copernicus Memorial Hospital).

Portugal: J Figueiredo, B Cismasiu, R Souto, S Henriques, AL Preto Barreira, J Vaz, JM Carlos, M Trindade, L Moreira, M Palas, J Simoes (Almada; Hospital Garcia de Orta); F Ramalho de Almeida, M Vasconcelos, A Neves, J Ribeiro, F Afonso, A Pita, R Miranda Pera, M Bernardo, C Rio Ferreira, T Branco, J Fontainhas, S Pimentel Morais, B Pinto (Amadora; Hospital Prof. Doutor Fernando Fonseca, E.P.E.); S Patrocínio, L Moniz, C Rolo Santos, P Bernardo, F Nazareth (Barreiro; Centro Hospitalar Barreiro Montijo, EPE); C Silva, L Heeren, AR Mateus Loureiro, B Tinoco, A Abreu (Caldas da Rainha; Hospital das Caldas da Rainha - Centro Hospitalar do Oeste, E.P.E.); DM Gonçalves Múrias Gomes, C Figueiredo, C Agüero, M Reia, M Guerrero, MA Fernandez Romero, J Dominguez (Elvas; Hospital Santa Luzia Elvas); I Colaço, M Nunes Luís, S Andrade, S Oliveira, D Pais (Figueira da Foz; Hospital Distrital da Figueira da Foz); DG Alves, F Castro, R Ribeiro, I Mogárrio, MDC Gama Caldeira (Funchal; Hospital Dr. Nélio Mendonça); B Gama, CS Rodrigues, A Cabral, A Silva (Horta; Hospital da Horta, E.P.E.); E Borges, J Cassiano Neves, R Bernardino, P David Santos, J Secchi (Lisbon; Centro Hospitalar Lisboa Norte); M Nunes, D Tavares, M Cruz, C Quintela, C Cardoso, IM Lourenço (Matosinhos; Unidade Local de Saude de Matosinhos - Hospital Pedro Hispano); D Vaz Acosta, P Rego Ponte, R Santos Pereira (Ponta Delgada; Hospital do Divino Espírito Santo); H Capote, MB Mourato, T Mogne, N Andrade, G Fialho, F Valente Costa Pinto, B Cordeiro, M Brito, G Santos, D Rosado, C Costa, N Pratas (Portalegre; Hospital Doutor José Maria Grande); J Dias-Ferreira, AL Carreira-Marques, R Ribeiro Dias, B Carvalho, M Gomes, C Soares-Aquino, F Gomes, S Barbosa Castelo Branco, C Coutinho, JP Vieira de Sousa, D Atouguia (Porto; Centro Hospitalar e Universitário de São João); L Cidade Costa (Porto; Hospital da Prelada); D Silva, P Correia, C Henriques, AM Pinheiro Pereira, J Marques Antunes (Santa Maria da Feira; Centro Hospitalar Entre o Douro e Vouga); H Devesa, R Barradas, S Fortuna Martins, N Marcos, A Jarimba, B Louro, L Rodrigues Madeira, AR Lourenço (Santarem; Hospital de Santarem); A Ferreira, A Abreu da Silva, D Stoian, M Ferreira (Santiago do Cacém; Hospital do Litoral Alentejano); R Branquinho, JC Domingues, MI Seixo, R Lalanda, C Bôto (Tomar; Centro Hospitalar Médio Tejo); J Fernandes, P Laranjo, M Reis, I Borges da Costa, C Assis, B Lopes Patrício, NM Freitas Oliveira (Vila Franca de Xira; Hospital Vila Franca de Xira); M Carvalho, J Mendes, C Macedo Cardoso de Oliveira, B Freire, R Pinheiro Duque (Vila Nova de Famalicao; Centro Hospitalar do Medio Ave); B Vieira, U Fernandes, A Dupont, J Ribeiro, R Vaz Pereira (Vila Real; Centro Hospitalar de Trás-os-Montes e Alto Douro, E.P.E.).

Qatar: G Sarp, E Soyer Güldoğan (Doha; Turkish Hospital).

Romania: É Gáspár (Brasov; Regina Maria); A Chitul, C Bezede, E Ciofic, D Cristian (Bucharest; Coltea Clinical Hospital); EA Toma, IM Matache, O Enciu, B Bogdan-Gabriel (Bucharest; Elias Emergency Hospital); I Negoii, C Ciubotaru, I Tanase, C Dina, VM Negoita, A Perja (Bucharest;

Emergency Clinical Hospital Bucharest); R Drasovean, A Trif, D Misca, C Hossu, I Imihteev (Cluj-Napoca; Medicover Hospital Cluj).

Russian Federation: V Kakotkin, M Agapov, V Budyakova, S Dos Santos Rocha Ferreira, R Senin (Kaliningrad; Immanuel Kant Baltic Federal University, Regional Clinical Hospital); A Bedzhanyan, A Sumbaev, K Petrenko, E Bedzhanyan, E Tyurina, R Azimov, P Glushkov, K Shemyatovsky, S Husanov, A Sidorova (Moscow; Petrovsky National Research Centre of Surgery); G Yarovenko, E Shestakov, O Lisin, A Arustamyan, S Katorkin (Samara; Hospital Surgery Clinic of Samara State Medical University); J Sidorovskaia, K Choloh, I Choloh, D Kurochka (Simferopol; Municipal Emergency Hospital No.6); V Ten, Y Kudryavcev (Yuzhno-Sakhalinsk; Private healthcare institution 'RZD-Medicine').

Rwanda: JP Rugambwa, CN Nelly Rosine, N Jeannette, A Dusabimana (Huye, Gisagara; Butare university teaching hospital (CHUB)); C Seneza, C Uwakunda, L Mukamazera, G Ntwari, I Didier (Kigali; Kibagabaga Hospital); A Costas-Chavarri, M Eugene, C Nyampinga, R Munyaneza, D Muyenzi (Kigali; Rwanda Military Hospital).

Saudi Arabia: N Alzerwi, M Rayzah, A Almutairi, A Alsultan, B Ali (Al-Majmaah; King Khalid General Hospital); A Shabkah, O Ibrahim, H Said, A Alhebshi, A Mohsen, K Anaam, F Alnazawi, F Haddad, A Basalim, B Albaihani, S Al Athath, A Jowharji, A Aljahdali (Jeddah; International Medical Center); N Trabulsi, M Alharthi, A Farsi, M Ghunaim, A Nawawi, A Maghrabi (Jeddah; King Abdulaziz University Hospital); N Alzerwi, Y Aldebasi, F Al Abbood, A Elkhalifa, M Alshanwani, S Alshagrawi (Riyadh; King Salman Hospital in Riyadh); F Ahmad (Riyadh; King Saud Medical City); A Alayed, R AlQahtani, R Sugair, O Alruwaili, B Alsharari, H Albalawi, H Al SOHABI, A Alshahrani, S Asiri, M Alshehri, A Albalawi, R Alatawi, T Khewater, H Adi, J Akiely, N Musawa, F Alahmad, B Alqahtani, M Sersarah, Z Farraj, DY Alalawi (Tabuk; King Salman Armed Forces Hospital); A Alzahrani, N Al Amri, M AlThomali, D Elkafrawy (Taif City; King Faisal Medical Complex).

Serbia: J Juloski, V Cuk, V Cijan, L Milic (Belgrade; Zvezdara University Medical Center).

Slovenia: JA Košir, J Grosek, A Tomazic, T Košir Božič (Ljubljana; University Medical Centre).

South Africa: S Gumede, C Kloppers, K Booyse, S Dos Santos, M Flint (Cape Town; Groote Schuur Hospital); Z Johnson, JJ Jordaan, G Steenkamp (Cape Town; Karl Bremer Hospital); K Nieuwenhuys, J Uys, SS Verhage, A Goliath, S Gilbert (Cape Town; Khayelitsha District Hospital); M Kariem, N Karimbocus (Cape Town; Mitchell's Plain District Hospital); C Lategan, T Mabogoane (Cape Town; Victoria Hospital Wynberg); S Mewa Kinoo, R Naidoo, N Ntanzu, S Sibiya, S Ebrahim (Durban; King Edward VIII Hospital); S Govender, E Naidoo, P Moodley, K Maharaj (Durban; Stanger Hospital); H Le Roux, J Van Niekerk (East London; Cecilia Makiwane Hospital); A Sparke, P Omwansa (East London; Frere Hospital); CA Baars, S Marawu, K Sevnanan, A Szpytko (Empangeni; Ngwelezana Hospital); G Charalambous, B Van Zyl, O Pheiffer, F Roodt (George; George Hospital); D Rattray, N Rasool, M Nkogotse, R Mackay, G Urdang (Johannesburg; Edenvale); V Manchev, D Clarke, S Naidu, V Govindasamy (Pietermaritzburg; Edendale Hospital); D Montwedi (Pretoria; Kalafong Academic Hospital); LC Kolongi, C Elliot-Wilson, S Kalenga, I Serfontein, M Goga (Upington; Dr Harry Surtie Hospital); S Burger, R Duvenage (Worcester; Worcester Provincial Hospital).

Spain: H Aguado López, F Ruescas, A García Marín, M Scortechini, M Jurado Román , A Sanchez Gallego (Albacete; Hellín Hospital); E González Marín, S De la Cruz Ahufinger, M Mateu, MJ Medina, MDM Martí-Ejarque, R Soliva Domínguez, L Ruiz-Villa, E Montalbán Martínez (BARCELONA; Hospital Universitari Sagrat Cor); A Torroella, C Ginesta, G Cárdenas Rivera, VE Gonzabay, JD Acevedo Parrales (Barcelona; HM Nou Delfos); M Canals Sin, A Lombardero, B Capdevila Vilaró, M Carbonell Pradas, ME Muñoz Fernández , RA Hernandez Rodriguez, I De Haro Jorge, M Riba Martínez, L Tapia Moral, M Coronas Soucheiron, P Palazon Bellver, L Ortega Lechuga, X Tarrado, A Domenech Plana, J Prat-Ortells, M Bejarano Serrano, M Cuesta Argos, MP Martin Gimenez, SG Laura, R Ripoll i Palmés (Barcelona; Hospital Sant Joan de Deu); A Sainz Lete, JC Zevallos-Quiroz, D Gómez, B Estraviz, JM De Francisco Rios, J Barrutia Leonardo, M González de Miguel (Bizkaia; Hospital Urduliz); RL Ferlini, M Ortega Escudero, Y Galvañ Félix, C Hernandez Diaz, J Montero García (Burgos; Hospital Universitario de Burgos); Á Fernández Camuñas, EP Garcia Santos, FJ Redondo Calvo (Ciudad Real; Hospital General Universitario de Ciudad Real); M Estaire Gómez, RJ Castro Lara, A Ramos Bonilla, L Rodríguez Gómez, M Marqueta De Salas, A Alvarez Cuiñas, FM Bujalance Cabrera, MD Cancelas, A García Domínguez, G Chamoso Mialdea, EP Cagigal Ortega, D Enjuto, I Cervera (Leganés; Severo Ochoa University Hospital); R Villalobos Mori, Y Maestre González, C Gas, L Codina Corrons, C Semeraro (Lleida; Hospital Universitari Arnau de Vilanova); J García-Quijada, TW Jorgensen, L Marquez (Madrid; Hospital Central de la Cruz Roja San Jose y Santa Adela); MJ Peña Soria, N Tabatabaian , JL Garcia galocha, D Fra Corral, L Sante Serna (Madrid; Hospital Clinico San Carlos); M Díez Alonso, C Vera Mansilla, L Casalduero, S Soto Schütte, Y Allaoua (Madrid; Hospital Universitario Principe de Asturias); E Gutierrez, C Zapata Syro, F Prieto La Noire, MDM Olmedo Reinoso, S Salido, N Chavarrias, M Vicario Bravo, L Asensio Gomez, R Abad, A Gegúndez Simón, PC Arteaga Asensio (Madrid; Hospital Universitario la Paz); AM Minaya Bravo, E González , A Galvan, C Guijarro Moreno, G De la Peña González, A Sánchez Gollarte, A Robin Valle de Lersundi, MÁ García Ureña (Madrid; Hospital del Henares); JL Rodicio Miravalles, AA Suárez Álvarez, DW Silva-Cano, G Martínez Izquierdo, P Del Val Ruiz, M Moreno Gijon, S Amoza Pais, GP Ibero Casadiego , E López-Negrete Cueto, J Carrizo, S Sanz, R Rodríguez-Uria, A Cembellin, G García-Santos, A Fraile (Oviedo; Hospital Universitario Central de Asturias (HUCA)); D Córdova García, L Jiménez, J Martin Fernandez, R Alvarado Hurtado, AM Minaya Bravo, R Díaz Pedrero, N Cobeño Tamayo, V Ongil Rodríguez (Rivas-Vaciamadrid, Madrid; Hospital Universitario HM Rivas); F Aguilar del Castillo, Á De Jesús Gil, S Borrego Canovaca (Sevilla; Hospital Universitario Virgen del Rocío); JR Naranjo Fernández, Z Valera Sanchez, R Perez, M Infantes Ormad, M Sánchez Ramirez (Seville; Hospital Universitario Virgen Macarena); C Leal Ferrandis, C Esteo Verdu, S García López, J Febré, B Cuneo (Valencia; Hospital Arnau de Vilanova); C León-Espinoza, E Martí Cuñat, G Pou (Valencia; Hospital Clínico Universitario de Valencia); C Jezieniecki, S Alonso Marcos, A Vazquez Fernandez, J Beltrán de Heredia, B De Andrés-Asenjo, D Baños Méndez, JC Garcia Vera, M Ruiz Soriano, E Redondo, R Martínez Díaz , T Gómez Sanz, P Artigot, C Infante, C Ferreras García, LR Cabezudo, G Cabezudo, M Lainez Escribano, M Rodríguez-Lopez, H Nuñez Del Barrio, A Romero de Diego (Valladolid; Hospital Clínico Universitario de Valladolid); A Vazquez Melero, M Camuera, I Herrero, D Garcia López de Goicoechea, M Sánchez-Rubio (Vitoria-Gasteiz; Hospital

Universitario Araba); MDP Cebollero, JL Blas Laina (Zaragoza; Hospital Royo Villanova); V Duque Mallén, N Sánchez Fuentes, P Sancho Pardo , I Gascon Ferrer, MÁ Dobón Rascón, T Gimenez Maurel, J Chóliz, S Saudi-Moro, S Paterna -Lopez, A Martinez German, D Aparicio-López, MÁ Gascón Domínguez, P Royo Dachary (Zaragoza; Hospital Universitario Miguel Servet).

Sri Lanka: S Srishankar, SPB Thalgasipitiya, KJ Senanayake, D Wickramarathna (Anuradhapura; Teaching Hospital Anuradhapura); D Subasinghe, D Wickramasinghe (Colombo; National Hospital of Sri Lanka); M Nandasena, K Wijesinghe, H Miyasika, J Senavirathna, Y Chamara (Dehiwala; Colombo South Teaching Hospital); S Rajendra, SI Thuraisamy Sarma, B Balagobi, V Sutharshan , S Giridaran (Jaffna; Teaching Hospital, Jaffna); W Wijenayake, MT Ekanayake, S Jayatilleke, S Jayasekara, M Perera, R Perera, R Ellawala, WDD De Silva (Werahera; University Hospital, Kotelawala Defence University).

Sudan: S Alqurashi , N Rajab (Ed Dueim; Ed Dueim Teaching Hospital); TA Albushary , FA Mohammed Daoud , A Younis, MA Suliman, F Tahir Lwdie, S Ibrahim Tour Harakan, M Abdelhadi Suliman Adam , E Hegab, A Abdalla, T Almahdi , E Alabed (El Geneina; El Geneina teaching Hospital); M Ahmed, S Eldirdiri, M Salah, OA Eljizoly , A Mohammed, A Mohamed Ibrahim Mohamed (Gadarif city; Gadarif teaching hospital); A Albager, MA Ismael Alamin, M Alsalawi, MA Elgak (Kassala; Police hospital); EG Nubi Mohamed, A Eltahir, E Adel Hamdoun Aziz, I Adel, O Emadeldeen, OG Nubi, E Mohamed (Khartoum; Bashair Teaching Hospital); M Hamed, M Tageldin, E Elsheikh, U Omara, E ADAm, IMG Ahmed, GMG Ahmed, S Imam (Khartoum; Ibrahim Malik Teaching Hospital); AA Adam, S Amin Omar Alsiddig, A Ahmed, S Abdelrasoul Elnour Ismail, M Mohamedshafee, Y Mohamed (Khartoum; Omdurman Teaching Hospital); EE Abuobaida Banaga Hag El Tayeb, H Abuobaida , AS Ahmed (Khartoum; Ribat university hospital); A Elbalal, IMG Ahmed, GN El Hunjul, GMG Ahmed, A Mustafa, HA Fadlalmola (Khartoum; Soba University Hospital); A Mohammed, E Yousuf, E Hamed, S Ibrahim, O Morgan, N Omer (Wad Madani; University of Gezira Hospital).

Sweden: M Zaigham, A Al Mukhtar (Malmö; Skåne University Hospital); M Nikberg (Vasteras; Västmanlands Hospital Västerås).

Switzerland: D Fenner, D Salinovic (Frauenfeld; Spital Thurgau AG); X Papazarkadas, TV Pham, C Brassat, A Litchinko, F Ris, C Golliez, M Chevallay (Geneva; Geneva University Hospitals); J Gass, J Mühlhäusser, J Metzger, A Scheiwiller (Luzern; Luzerner Kantonsspital); A Tampakis, C Riboni, U Dietz, E Brolese, C Seiler, M Kalisvaart, JN Marx, L Eisner (Olten; Kantonsspital Olten); G Peros, F Solimene, M Gramellini, A Lareida, M Adamina, E Betz, L Dubs, K Geiger-Timm, L Gantner, N Seeger, K Richetti, K Hofmann (Winterthur; Kantonsspital Winterthur); MA Schneider, D Gero, K Lehmann, P Limani, S Hügli, S Gerdes, F Mazzola, A Hiller (Zurich; University Hospital of Zurich).

Syrian Arab Republic: MA Farho, M Mohammad, AY Arnaout, Y Nerabani, Y Maktabi, W Alsado, A Anadani (Aleppo; Abd Al Wahab Agha Hospital); M Morjan, M Nasani, W Mayo, S Kreid, M Arnaout, MN Sawas (Aleppo; Al-Shahbaa Private Hospital); M Aloulou, MK Marawy, A Kezze, I Arnaout, A Kelzia, A Ghazal (Aleppo; Aleppo Private Hospital); A Ghazal, E Dabbagh, R Masri, MH Nabhan, A Alniemi, A Alhaj, S Ward, Y Haido, N Dadoush, MK Abu albahrain, A Niazi, W Abbas, A Hasan, S Alshab, S Kamari, R Kalouk, Z Toutounji (Aleppo; Aleppo University Hospital); D Sharl Ajami, B

Alsaid, A Alusef, K Abo zaal (Damascus; Al Assad university hospital); AR Hammadieh, Z Klib, MR Mslmani, A Alhaj zain, M Klib, L Shammas, AJ Chekfa, Z Odeh, R Joumaa, H Al-zoubi (Damascus; Al-Mouwasat University Hospital); Q Mashlah, HO Odah Bashi, H Zwaraa (Damascus; Children's University Hospital); I Adham, L Hasan, A Khatib, S Jomaa, A Alfarwan, A Torbey, A Rashid, A Hawarah, A Ali, M Alhimyar, L Al-Boukhari, MA Al-yusuf, AN Aldirani (Damascus; Damascus Hospital); Y Alhammoud , Y Al-Junaidi (Homs; Al-Baseel Specialized Hospital in Karm El-Louz); M Daher, Z Asaad, A Abbas, K Khalil (Homs; The Military Hospital); J Khoury, L Hasan, J Jahjah, J Alaji, S Turkmani, S Mahfoud, A Ahmad, M Derattani, G Massarra , J Skaff, ZA Hannouneh, H Amoudi, ZA Zaher, G Zaza (Latakia; Al Saydeh Surgical Hospital); A Wassouf, A Alahmad Alismael , S Nofal, A Mansour, M Mansour, Z Alkhaier, J Suliman, M Sabboh, M Haj Hussein, M Ibrahim, ZA Abo alaros, G Alhadwah , N Kheyrbek, D Ibrahim, G Hamdan, Y Hasan, A Abo al Shamat, A Roumieh, B Khattab, H Alkhatib (Latakia; National Hospital); S Hassan, S Abdul Rahman, A Abdul Rahman, F Aliskander, J Alkharish, G Kafa, I Suleiman, A Bassma, A Alloush , N Ismaiel, J Deeb, M Alrantisi, E Salloum, A Al-Mouahhed , A Baydoun, H Yunes, S Alkadi, F Ali, D Abdulrahman, I Hussein (Latakia; Othaman Hospital); A Bakri, H Asaad, T Ashkar, H Daaboul, A Marouf, F Chahrour, B Ranjous, B Ibrahim , A Sinjab, M Alneasan, N Ali, A Alloush, J Fahed, R Attaf, S Kanaan (Latakia; Tishreen University Hospital).

Thailand: A Tansawet, W Kasetsermwiriya, I Laopeamthong, P Sukhvibul, T Techapongsatorn, N Techapongsatorn, P Kasetsermwiriya , P Leungon (Bangkok; Vajira hospital).

Togo: PS Tekam Wadje (Lomé; CHU Sylvanus Olympio).

Tunisia: MJ Kacem, R Elaifia, Y Ouadi, S Megdiche, Y Jedidi (Tunis; La Rabta Hospital).

Turkey: AÇ Bozkurt, H Tümer (Adana; Adana Seyhan State Hospital); MA Koç, A Çakmak, AF Kocaay, KY Türker, TB Türkmen (Ankara; Ankara University Medical School); O Yalkın, D Yigit (Bursa; Bursa City Hospital); B Yigit, A Aslan, S Yılmaz (Elazig; Elazig Fethi Sekin City Hospital); AN Sanli, Yİ Tandoğan, A Yıldiz, A İslar, A Ozkomec (Gaziantep; Abdulkadir Yuksel State Hospital); ME Seker, E Ay, M Erkaya, YO Koyluoglu, A Develioğlu, GM Kurtoglu, GK Kurtoglu, AF Cetişli, E Tunçcan, A Aghayeva, Z Durna, B Baca (Istanbul; Acibadem Altunizade Hospital); AE Dönmez, B Togay, E Ada, IE Yavuz , IA Bilgin, EC Karabulut, AM Uysal, Y Karataş, B Ağca, MK Aktas, F Demiral, B Duman, K Kabulov, I Hamzaoglu, T Karahasanoğlu, M Tanal, E Tuzuner (Istanbul; Acibadem Maslak Hospital); S Meriç, M Tokocin, H Yigitbas, A Barcin, G Alici, E Yavuz, OB Gülcicek, N Bugdayci , K Özdoğan, I Çakır, YE Aktimur, Y Altinel (Istanbul; Bağcilar Research And Training Hospital); ÖP Zambak Mutlu (Istanbul; Bahçelievler State Hospital); RE Sönmez, M Şermet, MS Ozsoy, H Baysal, F Buyuker (Istanbul; Istanbul Medeniyet University, School of Medicine); M Oncel, S Bektas, AE Askin, M Yashar, A İzgiş (Istanbul; Istanbul Medipol University Hospital); SS Uludağ, MF Ozcelik, S Yumurtacılar (Istanbul; Istanbul universty - Cerrahpaşa Medical faculty); A Özcan, E Somuncu, S Yılmaz, A Sapmaz, H Bolukbasi, C Özkan, E Bozdağ, MC Kizilkaya, H Telci, Y Kara, AZ Kaan, M Acar, EO Yildirim (Istanbul; Kanuni Sultan Suleyman Training and Research Hospital); G Akcakoca, C Hacialioğlu, Y Tosun, V Çalik, TE Yılmaz, H Alfakeer (Istanbul; Kartal Dr. Lutfi Kirdar Training and Research Hospital); MI Ateş, SN Karahan, M Kalender, AE Narin, D Yiğit, O Agcaoglu, S Toprak, B Celik, E Ozoran, DS Uymaz, S Yigman, E Bozkurt, A Rencuzogullari, E Balik, S Sucu

(Istanbul; Koç University Medical School); A Akmercan, K Oğur, A Hajali, QK Dolatzay (Istanbul; Marmara University, School of Medicine); E Unal (Istanbul; Sehit Prof.Dr. İlhan Varank Training and Research Hospital); N Kiziltoprak, MS Genç, B Özcan, Z Şenol, OF Ozkan, M Çuhadar, ED Terzi (Istanbul; Sultan 2. Abdulhamid Han Training and Research Hospital, University of Health Sciences); T Gülşen (Istanbul; Sultanbeyli State Hospital); MT Demirpolat (Istanbul; University of Health Science Umraniye Education and Research Hospital); B Citgez, H Ozsahin, C Ersavas (Istanbul; Uskudar University Faculty of Medicine, Memorial Hospital); C Bilir, AE Boztaş Demir, AD Hacıoglu, G Ozyuksel, H Ulman (Izmir; Bakircay University Cigli Training and Research Hospital); AM Öztürk, B Calik, AC Yaşar (Izmir; University of Health Sciences Izmir Bozyaka Training and Research Hospital); E Colak, MA Avci, E Aybar, M Gün, AB Ciftci, MS Uyanik, ME Kara, C Akgün, AC Sarı, Ö Küpçüoğlu, GO Kucuk, S Polat (Samsun; Samsun University Samsun Training and Research Hospital); N Kavak, MA Kara, G Karadeniz Cakmak, B Kum (Zonguldak; Zonguldak Bulent Ecevit University School of Medicine Research and Training Hospital); S Öztürk, B Eyduran (İzmir; University of Health Sciences Tepecik Training and Research Hospital).

Uganda: B Kigwe, M Arafat (Iganga; Iganga district hospital); M Nnabagulanyi (Kigumba; Kiryandongo Hospital); S Stonelake (Luwero; Kiwoko Hospital).

United Kingdom: ZA Fozo, A Shoker, KR Rahman, MW Saqib, R Faisal, CS Ong, A Pillai, H Unwin, A Huws, M Maybury, H Ejaz, E Daketsey, AK Lala (Bangor, North Wales; Ysbyty Gwynedd); MAK Sarker, B Chkir, MR Peris, S Khan, MA Tahir, N Sharma, R Doherty, H Alhusaini, MF Butt, H Afzal, S Handa, N Maharjan, A Mostafa, G Lee, CK Lim, A Anand, A Krishna, WT Yew, Y Lu, R Hall (Barrow in Furness; Furness General Hospital); E Mohammed (Basildon; Basildon University Hospital); F Georgiades, S Karim, K Rajaratnam (Bedford; Bedford Hospital); J Abbasy, A Bibi, S Karandikar, L Johnstone, N Fazili (Birmingham; Heartlands Hospital); A Singh, A Athanasiou, H Lidbetter, J Siby, M Kaur, A Fatima (Brighton; Royal Sussex County Hospital); SM Reddy, A Campbell, A Cardoso Almeida, K Smith, CJ Bradshaw, K Tambudze, H Delacave, I Norman (Bristol; Bristol Royal Hospital for Children); M San, S Babu, S Midya, H Bradly, S Tontus, H Chauhan, R Jurdon, M Corcos, E Jose (Camberley; Frimley Health NHS FT - Frimley Park); N Eardley, B Davies, M Ransome, S Ahmed (Chester; Countess of Chester Hospital); S Suresh, A Bavaharan, R Batir, R Sato (Colchester; Colchester Hospital University); N Chidumije, M Ahmed, TY Kwan, F Olaniru, I Parwaiz, LP Cheng (Coventry; University Hospitals Coventry and Warwickshire NHS Trust); N Gokhare Viswanath, K Nanayakkara, A Tibude (Derby; Royal Derby Hospital); O Olajumoke, M Shams, D George, A Amin, M Kausar, S Sellahewa (Dudley; Russell's Hall Hospital); H Kamal, A Kamal, M Kamal (Dundee; Ninewells Hospital); O Pryer, H Sagar, B Lulham-Robinson, S Dawo, A Nada (Durham; University Hospital North Durham); R Bethune, G Chillarge, AM Myintmo, M Horga, L Andreski, E Ruiz-Daum, T Finlay, I Rakshit (Exeter; Royal Devon and Exeter Hospital); J Bryan, D Joshi, R Marlin, M Battili, YL Aung, L Zeng, S Mathew, I Njere, S Gopaul, A Abbas (Great Yarmouth; James Paget Univeristy NHS Foundation Trust Hospital); G Singh, M Asarbakhsh, S Staight (Huddersfield; Huddersfield Royal Infirmary); R Govindaraju, M Quaunine, S Yassin, A Wilkins, J Walshaw, L Chang, A Mahendran, F Hammett, D Fairbrass, UA Kalu, E Dexter, T Nadeem, N Karunaratne, T Lo, M Pellen (Hull; Hull University Teaching Hospitals NHS Trust); SH Sarwary, J Otote, D Anbu, H Islam, N Morricone (Ilford;

King George Hospital); K Lee, E Gimson, M Wilson, C Chiam (Larbert; Forth Valley Royal Hospital); M Solkar, M Bautista, NS Blencowe, A Gupta, J Sutcliffe, A Ahmad, A Peckham-Cooper (Leeds; Leeds Teaching Hospitals); E O'Connell, KE Dey, R Lunevicius, MM Barakat, GR Goodwin, B Devkaran, IC Nzenwa, A Pilavas (Liverpool; Aintree University Hospital); K Bananis, A Alamin, S Bennett, MD Barcelona, A Sharp, F Soggiu (London; Ealing Hospital); E Baili, H Ebied, A Botha, M Haghghat Ghahfarokhi, MMT Youssef (London; Guy's and St Thomas' Hospitals); K Theodoropoulou, A Quddus, R Hegy, M Mahran, A Ghanbari (London; Homerton University Hospital); P Kapsampelis, T Chouari, J Saunders, C Boven, I Gerogiannis (London; Kingston); N Karthikeyan, C Karagianni, E Spanoudakis (London; Queen Elizabeth Hospital, Woolwich); H Younus, R Ben Hmida, D Eaton (London; The Whittington Hospital); C Seet, R Bradley, R Roberts (London; University Hospital Lewisham); FN Amir, M Durrani, S Khan, A Tahir (Macclesfield; Macclesfield District General Hospital); A Khalil, E O'Neill, S Ingley, V Bill, P Wilson, M Elmousili, E Chin, C Alphonse, P Suresh, L Devi, C Shelton (Manchester; Wythenshawe Hospital); S Bugren, M Abdelreheem, MMR Azzuz, F Gareb (Margate; Queen Elizabeth the Queen Mother Hospital Margate); DJ Dhillon, MF Khan, B Peter, I Fagiri, T Harris, P Thambi (Middlesbrough; James Cook University Hospital); M Tomlinson, C Hidalgo Salinas, A Mwanjoka, M Catterall, B Ali (Morecambe; Royal Lancaster Infirmary); S Tingle, K Waddell, F Peters, T Akharaekpanya, S Robinson (North Shields; Northumbria NHS Hospital Trust); NA Binti Yusri, A Ashiru, SA Chowdhury, J Reilly, S Malek, S Kumaran, A Doghaim, ABA Al-Hajjaj, LR Chieng, ZY Wong, R Olatunji (Nottingham; Queens Medical Centre); J Bundred, B Down, A Ang, KA Shamiyah, G Bond-Smith (Oxford; John Radcliffe Hospital); M Elmesalmi (Portsmouth; Queen Alexandra Hospital); S Monkhouse, O Mohamed, B Robertson-Jones, P Patel, BZ Hao (Redhill; East Surrey Hospital); S Ghattas, N Beharry, A Maraqa, N Uttam, S Rafiq, A Abdelhamid, E Mazumdar, M Dyer, MEEA Abdelsalam, B Johnson, M Abdelkarim, A Murtada, MMS Tora, Z Azhar (Rhyl; Glan Clwyd Hospital); O Whitehurst, K Bhatti, S Silvestre, LN Bin Aizan, A Patel (Salford; Salford Royal Hospital); Z Khan, A Maqsood-Shah, O Webster, H Reilly, S Boyes (Sheffield; Sheffield Teaching Hospital NHS Foundation Trust); S Bandyopadhyay, B McDermott, H Kynaston, B Neall, M West, G Hart, R Titcombe, H Kaur, A Ekerin, A Demetriou, N Harrison, JJQ Chen (Southampton; Southampton General Hospital); M Hammoda, L Robine-Durnell, E Mansour, A Evans, E Baker (Swansea; Morriston Hospital Swansea); I Abdullah, O Osunlusi (Walsall; Walsall Manor Hospital).

United States: S Soelling, R Askari (Boston; Brigham and Women's Hospital); M Asaad, J Leong, M Perkins, D Ozal, SS Budhwani (Chicago, IL; Mount Sinai Hospital); A Maxwell, A Shah, S Schimpke (Chicago, IL; Rush University Medical Centre); D Moris, C Nicholson Jr., HE Rice (Durham, NC; Duke University Medical Center); D Ridder, M Tsuruta, K Noguchi, D Mikami, R Kitamura, J Ng-Kamstra (Honolulu, Hawaii; The Queen's Medical Center); RJ Robitsek, K Fretwell, J Chan, R Laskowski (Jamaica; Jamaica Hospital); R Zerna Encalada (Syracuse; SUNY Upstate University Hospital).

Yemen, Rep.: R Ghaleb, N Alnamari, A Al-Bahla, B Al soudi (Hajjah; Kowaydina hospital); B Alshaikh, M Al-Shehari, M Al-Dhaheiri, YSSM Ghaleb, MI Issa (Sana'a; Al-Thawra Modern General Hospital).

Section 2: Supplementary materials, tables and figures

Supplementary materials	Pages
Development of measurement set	36
Data collection	37
Supplementary table title	
Supplementary table 1: Measurement set mapped to WHO's building blocks attributes	38
Supplementary table 2: Detailed description of key performance measures.	39
Supplementary table 3: Assumptions to calculate sample size.	40
Supplementary table 4: Included elective patients and intra-operative characteristics across income groups.	41
Supplementary table 5: Included elective, adult patients and intra-operative characteristics across income groups.	42
Supplementary table 6: Included patients eligible for day case surgery and intra-operative characteristics across income groups.	43
Supplementary table 7: Hospitals characteristics across income groups.	44
Supplementary table 8: Hospitals characteristics across income groups, based on the number of hospitals	45
Supplementary table 9: Key performance measures to evaluate access and quality.	46
Supplementary table 10: Additional descriptive measures to evaluate access and quality.	47
Supplementary figure title	
Supplementary figure 1: Flowchart of included patients according to income groups and urgency of surgery.	48
Supplementary figure 2: Global distribution of included patients.	49
References	50

Development of measurement set

We formed a diverse Study Management Group with surgeons and anaesthetists with expertise in both inguinal hernia treatment and health systems research from different income groups to ensure global representation (Appendix 1, section 1). We predicted that no single outcome would be able to evaluate performance of whole surgical systems. The aim of the consensus process therefore was to capture key outcome measures in elective system performance that would have overall face and construct validity (i.e., measure what they purport to measure) and content validity, i.e. measured all relevant attributes of the World Health Organisation's Health System Building Blocks, namely: (1) Access, (2) Coverage, (3) Quality and (4) Safety.

To achieve this, the Study Management Group conducted a three-phase consensus process to identify, prioritise, and refine a measurement set for these attributes. In brief, phase one included scoping reviews of published literature from PubMed and Google Scholar, guidelines from hernia societies, and policy documents to create a longlist of candidate measures. In phase two, the Study Management Group conducted an online consensus process. The longlist was refined into a shortlist during an online meeting using Zoom® (Zoom Corporation, Tokyo, Japan). The rationale to include and/or exclude each measure was discussed and consensus reached for each. In phase three, the measurement set and its definitions were refined into six key performance measures and seven additional descriptive measures during a face-to-face meeting in Lagos, Nigeria. The final measurement set was deemed to have face, construct, and content validity by the SMG, and to be relevant to surgical systems in any context, including all hospital types and income settings (appendix 1, p 38-39).

Data collection

Each team based in each participating hospitals followed the strategy that worked best for them to collect high quality data, which could either be on paper or electronically. Data was uploaded in REDCap either by the hospital lead or by the collaborators and all elements from the same team, including the hospital lead, had access to all data entered by their team so they could ensure it was high-quality, not duplicated, and valid data.

Some participating hospitals and teams had previous experience with this approach as they took part in previous cohort studies (GlobalSurg, CovidSurg, SurgWeek). Online training was provided by the Steering Management Group on data collection strategies, entering data in REDCap and ensuring data was valid. The Dissemination Committee members were encouraged and supported to run similar online sessions to hospitals taking part from their countries. Online materials to support data collection and data validation were available in the HIPPO webpage (<https://www.globalsurgeryunit.org/clinical-trials-holding-page/hippo/>) and in the YouTube page of the study (<https://www.youtube.com/@hipporedcap>). Members of the Steering Management Group hosted live meetings open to all collaborators taking part in the study to clarify any questions, which were recorded and available in the YouTube page too.

Supplementary table 1: Measurement set mapped to WHO's building blocks attributes.

Attributes of WHO building blocks	Key performance measures	Additional descriptive measures
Access and coverage	Emergency rate	Standardised patient pathways
	Bowel resection rate	Availability of day case surgical unit
	Waiting times with its components	Waiting list management
		Financing methods
Quality and safety	Mesh use rate	Primary operator
	Day case rate	Anaesthesia variation
	Postoperative complications	Surgical approach variation

Supplementary table 2: Detailed description of key performance measures.

Key performance measures	Rationale	Numerator	Denominator
ACCESS AND COVERAGE			
Emergency surgery rate	Indirect measure of access to elective surgery, as patients who cannot receive their care in the elective pathway, will experience a progression of their disease, increasing the risk of developing complications with time and need for emergent surgery instead. ¹	Patients undergoing emergency surgery	All patients included
Bowel resection rate	Indirect measure of access to elective surgery, theoretically related with emergency surgery, as patients are more likely to develop complications from their hernia over time, while waiting for an operation. ²	Patients where bowel resection occurred	All patients included
Waiting times	Direct measure of access to elective surgical systems that is considered relevant to patients. Longer waiting times imply poorer access. ³	NA*	NA*
QUALITY AND SAFETY			
Mesh use rate	Direct measure of quality as recommended by <i>International Guidelines</i> , ⁴ applicable to all adults undergoing elective inguinal hernia repair.	Patients where mesh was used for hernia repair	Adult patients (>16y) undergoing elective surgery
Day case rate	Direct measure of quality as recommended by the <i>International Guidelines</i> , ⁴ to adults younger than 90y and with ASA I-II undergoing elective inguinal hernia repair. Adults with ASA III with stable co-morbidities is also recommended but this distinction was not possible to be made from our dataset.	Patients discharged as planned, in the same day as the operation	Adult patients, younger than 90y, with ASA I-II
Postoperative complications	Direct measure of safety as established by the WHO. Routinely collected to monitor safety of all surgical procedures. ⁵	Patients with Clavien-Dindo grade I-V	All patients included

*NA: waiting times to elective surgery is a continuous outcome that was measured in all patients undergoing elective surgery, therefore, the denominator and numerator is not applicable to directly calculate this measure. References presented in page 50.

Supplementary table 3: Assumptions to calculate sample size.

Binary outcomes			
Outcome	Alpha	Estimated proportion	Margin of error
Emergency surgery rate	5%	6% ¹	0.47%
Bowel resection rate	5%	1% ²	0.20%
Day case rate	5%	75% ^{4,6}	0.85%
Mesh use rate	5%	95% ⁴	0.43%
Post-operative complications	5%	15% ⁴	0.70%
Continuous outcome			
Outcome	Alpha	Standard deviation	Width
Waiting time to elective surgery	5%	10	0.39

References presented in page 50.

Supplementary table 4: Included elective patients and intra-operative characteristics across income groups.

	HIC (n=9,252)	UMIC (n=93,098)	LMIC (n=3,586)	LIC (n=818)	Total (n=16,754)
Age, years					
Median	63.0	53.0	42.0	45.0	57.0
IQR	50.0 to 73.0	32.0 to 65.0	11.0 to 58.0	20.0 to 60.0	40.0 to 69.0
Age groups*					
Infants	107 (1.2%)	211 (6.8%)	330 (9.2%)	64 (7.9%)	712 (4.3%)
Children	228 (2.5%)	310 (10.0%)	614 (17.1%)	116 (14.2%)	1268 (7.6%)
Adults	8916 (96.4%)	2575 (83.2%)	2642 (73.7%)	635 (77.9%)	14768 (88.2%)
(Missing)	1	2	0	4	6
Sex					
Female	907 (9.8%)	344 (11.1%)	334 (9.3%)	74 (9.1%)	1659 (9.9%)
Male	8344 (90.2%)	2754 (88.9%)	3252 (90.7%)	741 (90.9%)	15091 (90.1%)
Missing	1	0	0	3	4
ASA grade					
ASA I-II	7530 (81.4%)	2852 (92.1%)	3462 (96.5%)	787 (96.6%)	14631 (87.3%)
ASA III-V	1661 (18.0%)	240 (7.7%)	116 (3.2%)	21 (2.6%)	2038 (12.2%)
Not recorded	61 (0.7%)	6 (0.2%)	8 (0.2%)	7 (0.9%)	82 (0.5%)
(Missing)	0	0	0	3	3
Co-morbidities					
None	6948 (75.1%)	2479 (80.1%)	3121 (87.0%)	706 (86.3%)	13254 (79.1%)
One	1714 (18.5%)	496 (16.0%)	384 (10.7%)	100 (12.2%)	2694 (16.1%)
Two	456 (4.9%)	93 (3.0%)	73 (2.0%)	9 (1.1%)	631 (3.8%)
Three or more	130 (1.4%)	26 (0.8%)	5 (0.1%)	0 (0.0%)	161 (1.0%)
(Missing)	4	3	3	3	13
Symptoms					
Asymptomatic	1312 (14.2%)	520 (16.8%)	974 (27.2%)	117 (14.3%)	2923 (17.4%)
Symptomatic	7940 (85.8%)	2578 (83.2%)	2612 (72.8%)	701 (85.7%)	13831 (82.6%)
(Missing)	0	0	0	0	0
Hernia size					
Limited to inguinal region	7869 (85.1%)	2348 (75.8%)	2247 (62.7%)	583 (71.5%)	13047 (77.9%)
Limited to scrotum	1306 (14.1%)	719 (23.2%)	1291 (36.0%)	224 (27.5%)	3540 (21.1%)
Extend to mid-thigh or beyond	77 (0.8%)	31 (1.0%)	48 (1.3%)	8 (1.0%)	164 (1.0%)
(Missing)	0	0	0	3	3
Contamination					
Clean	9194 (99.4%)	2883 (93.1%)	3479 (97.0%)	814 (99.5%)	16370 (97.7%)
Clean-Contaminated	53 (0.6%)	212 (6.8%)	104 (2.9%)	4 (0.5%)	373 (2.2%)
Contaminated	4 (0.0%)	2 (0.1%)	2 (0.1%)	0 (0.0%)	8 (0.0%)
Dirty	1 (0.0%)	0 (0.0%)	1 (0.0%)	0 (0.0%)	2 (0.0%)
(Missing)	0	1	0	0	1
Hernia defect size					
< 1.5cm	2062 (22.3%)	659 (21.3%)	932 (26.0%)	220 (26.9%)	3873 (23.1%)
1.5cm to 3cm	3689 (39.9%)	1160 (37.4%)	1306 (36.4%)	337 (41.2%)	6492 (38.8%)
>3cm	1806 (19.5%)	871 (28.1%)	1027 (28.6%)	192 (23.5%)	3896 (23.3%)
Not known	1692 (18.3%)	408 (13.2%)	321 (9.0%)	69 (8.4%)	2490 (14.9%)
(Missing)	3	0	0	0	3

Data are n (%) unless stated otherwise. Denominators for each group are presented at the top row.

HIC: high-income country, UMIC: upper-middle-income country, LMIC: lower-middle-income country, LIC: low-income country, ASA: American Society of Anaesthesiologists Physical Status Classification System grade.

Supplementary table 5: Included elective, adult patients and intra-operative characteristics across income groups

	HIC (n=8,916)	UMIC (n=2,575)	LMIC (n=2,642)	LIC (n=635)	Total (n=14,768)
Age, years					
Median	63.0	57.0	52.0	53.0	60.0
IQR	(52.0 to 73.0)	(45.0 to 67.0)	(37.0 to 63.0)	(36.0 to 63.5)	(47.0 to 70.0)
Sex					
Female	821 (9.2%)	219 (8.5%)	142 (5.4%)	46 (7.2%)	1228 (8.3%)
Male	8094 (90.8%)	2356 (91.5%)	2500 (94.6%)	589 (92.8%)	13539 (91.7%)
(Missing)	1	0	0	0	1
ASA grade					
ASA I-II	7211 (80.9%)	2337 (90.8%)	2536 (96.0%)	607 (95.6%)	12691 (85.9%)
ASA III-V	1645 (18.4%)	234 (9.1%)	105 (4.0%)	21 (3.3%)	2005 (13.6%)
Not recorded	60 (0.7%)	4 (0.2%)	1 (0.0%)	7 (1.1%)	72 (0.5%)
(Missing)	0	0	0	0	0
Co-morbidities					
None	6613 (74.2%)	1961 (76.2%)	2195 (83.1%)	526 (82.8%)	11295 (76.5%)
One	1713 (19.2%)	491 (19.1%)	367 (13.9%)	100 (15.7%)	2671 (18.1%)
Two	456 (5.1%)	93 (3.6%)	72 (2.7%)	9 (1.4%)	630 (4.3%)
Three or more	130 (1.5%)	26 (1.0%)	5 (0.2%)		161 (1.1%)
(Missing)	4	3	3	0	10
Symptoms					
Asymptomatic	1106 (12.4%)	293 (11.4%)	580 (22.0%)	60 (9.4%)	2039 (13.8%)
Symptomatic	7810 (87.6%)	2282 (88.6%)	2062 (78.0%)	575 (90.6%)	12729 (86.2%)
(Missing)	0	0	0	0	0
Hernia size					
Limited to inguinal region	7576 (85.0%)	1977 (76.8%)	1698 (64.3%)	482 (75.9%)	11733 (79.4%)
Limited to scrotum	1263 (14.2%)	569 (22.1%)	903 (34.2%)	147 (23.1%)	2882 (19.5%)
Extend to mid-thigh or beyond	77 (0.9%)	29 (1.1%)	41 (1.6%)	6 (0.9%)	153 (1.0%)
(Missing)	0	0	0	0	0
Contamination					
Clean	8858 (99.3%)	2383 (92.5%)	2546 (96.4%)	632 (99.5%)	14419 (97.6%)
Clean-Contaminated	53 (0.6%)	190 (7.4%)	93 (3.5%)	3 (0.5%)	339 (2.3%)
Contaminated	4 (0.0%)	2 (0.1%)	2 (0.1%)	0 (0.0%)	8 (0.1%)
Dirty	1 (0.0%)	0 (0.0%)	1 (0.0%)	0 (0.0%)	2 (0.0%)
(Missing)	0	0	0	0	0
Hernia defect size					
< 1.5cm	1866 (20.9%)	456 (17.7%)	425 (16.1%)	138 (21.7%)	2885 (19.5%)
1.5cm to 3cm	3654 (41.0%)	1018 (39.5%)	1063 (40.2%)	296 (46.6%)	6031 (40.8%)
>3cm	1804 (20.2%)	820 (31.8%)	920 (34.8%)	175 (27.6%)	3719 (25.2%)
Not known	1590 (17.8%)	281 (10.9%)	234 (8.9%)	26 (4.1%)	2131 (14.4%)
(Missing)	2	0	0	0	2

Data are n (%) unless stated otherwise. Denominators for each group are presented at the top row.

HIC: high-income country, UMIC: upper-middle-income country, LMIC: lower-middle-income country, LIC: low-income country, ASA: American Society of Anaesthesiologists Physical Status Classification System grade.

Supplementary table 6: Included patients eligible for day case surgery and intra-operative characteristics across income groups

	HIC (n=7,189)	UMIC (n=2,332)	LMIC (n=2,532)	LIC (n=605)	Total (n=12,658)
Age, years					
Median	61.0	56.0	51.0	52.0	58.0
IQR	(50.0 to 70.0)	(43.0 to 66.0)	(36.0 to 62.0)	(35.0 to 62.0)	(45.0 to 68.0)
Sex					
Female	681 (9.5%)	207 (8.9%)	137 (5.4%)	46 (7.6%)	1071 (8.5%)
Male	6507 (90.5%)	2125 (91.1%)	2395 (94.6%)	559 (92.4%)	11586 (91.5%)
(Missing)	1	0	0	0	1
Co-morbidities					
None	6037 (84.0%)	1882 (80.7%)	2168 (85.6%)	512 (84.6%)	10599 (83.7%)
One	1011 (14.1%)	384 (16.5%)	315 (12.4%)	85 (14.0%)	1795 (14.2%)
Two	127 (1.8%)	49 (2.1%)	44 (1.7%)	8 (1.3%)	228 (1.8%)
Three or more	10 (0.1%)	14 (0.6%)	2 (0.1%)	0 (0.0%)	26 (0.2%)
(Missing)	4 (0.1%)	2 (0.1%)	3 (0.1%)		9 (0.1%)
Symptoms					
Asymptomatic	929 (12.9%)	275 (11.8%)	566 (22.4%)	57 (9.4%)	1827 (14.4%)
Symptomatic	6260 (87.1%)	2057 (88.2%)	1966 (77.6%)	548 (90.6%)	10831 (85.6%)
(Missing)	0	0	0	0	0
Hernia size					
Limited to inguinal region	6274 (87.3%)	1813 (77.7%)	1638 (64.7%)	462 (76.4%)	10187 (80.5%)
Limited to scrotum	872 (12.1%)	494 (21.2%)	862 (34.0%)	138 (22.8%)	2366 (18.7%)
Extend to mid-thigh or beyond	43 (0.6%)	25 (1.1%)	32 (1.3%)	5 (0.8%)	105 (0.8%)
(Missing)	0	0	0	0	0
Contamination					
Clean	7147 (99.4%)	2150 (92.2%)	2442 (96.4%)	603 (99.7%)	12342 (97.5%)
Clean-Contaminated	39 (0.5%)	181 (7.8%)	88 (3.5%)	2 (0.3%)	310 (2.4%)
Contaminated	3 (0.0%)	1 (0.0%)	2 (0.1%)	0 (0.0%)	6 (0.0%)
Dirty	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
(Missing)	0	0	0	0	0
Hernia defect size					
< 1.5cm	1647 (22.9%)	417 (17.9%)	413 (16.3%)	131 (21.7%)	2608 (20.6%)
1.5cm to 3cm	2997 (41.7%)	943 (40.4%)	1019 (40.2%)	286 (47.3%)	5245 (41.4%)
>3cm	1300 (18.1%)	723 (31.0%)	876 (34.6%)	164 (27.1%)	3063 (24.2%)
Not known	1243 (17.3%)	249 (10.7%)	224 (8.8%)	24 (4.0%)	1740 (13.7%)
(Missing)	2	0	0	0	2

Data are n (%) unless stated otherwise. Denominators for each group are presented at the top row.

HIC: high-income country, UMIC: upper-middle-income country, LMIC: lower-middle-income country, LIC: low-income country, ASA: American Society of Anaesthesiologists Physical Status Classification System grade.

Supplementary table 7: Hospitals characteristics across income groups

	HIC (n=9,808)	UMIC (n=3,366)	LMIC (n=3,948)	LIC (n=936)	Total (n=18,058)
Type of hospital					
Primary level	1002 (10.3%)	129 (3.9%)	201 (5.1%)	164 (18.1%)	1496 (8.4%)
Secondary level	3562 (36.6%)	598 (18.1%)	490 (12.4%)	96 (10.6%)	4746 (26.5%)
Tertiary level	5168 (53.1%)	2572 (78.0%)	3257 (82.5%)	648 (71.4%)	11645 (65.1%)
(Missing)	76	67	0	28	171
Hospital funding					
Public	8507 (87.4%)	2651 (80.4%)	3023 (76.6%)	839 (92.4%)	15020 (84.0%)
Private	582 (6.0%)	532 (16.1%)	805 (20.4%)	69 (7.6%)	1988 (11.1%)
Public-Private	643 (6.6%)	116 (3.5%)	120 (3.0%)	0 (0.0%)	879 (4.9%)
(Missing)	76	67	0	28	171
Surgical emergency service					
Yes - surgery available 24h	8989 (92.4%)	3031 (91.9%)	3756 (95.1%)	667 (73.5%)	16443 (91.9%)
Yes - surgery only during daytime	490 (5.0%)	66 (2.0%)	142 (3.6%)	157 (17.3%)	855 (4.8%)
Yes - for assessment only	184 (1.9%)	140 (4.2%)	50 (1.3%)	12 (1.3%)	386 (2.2%)
No	69 (0.7%)	62 (1.9%)	0 (0.0%)	72 (7.9%)	203 (1.1%)
(Missing)	76	67	0	28	171

Data are n (%) unless stated otherwise. Denominators for each group are presented at the top row.

HIC: high-income country, UMIC: upper-middle-income country, LMIC: lower-middle-income country, LIC: low-income country.

Supplementary table 8: Hospitals characteristics across income groups, based on the number of hospitals.

	HIC (n=303)	UMIC (n=145)	LMIC (n=146)	LIC (n=46)	Total (n=640)
Type of hospital					
Primary level	32 (10.8%)	12 (8.7%)	17 (11.6%)	10 (22.7%)	71 (11.4%)
Secondary level	115 (38.9%)	48 (34.8%)	27 (18.5%)	8 (18.2%)	198 (31.7%)
Tertiary level	149 (50.3%)	78 (56.5%)	102 (69.9%)	26 (59.1%)	355 (56.9%)
(Missing)	7	7	0	2	16
Hospital funding					
Public	255 (86.1%)	103 (74.6%)	103 (70.5%)	39 (88.6%)	500 (80.1%)
Private	22 (7.4%)	29 (21.0%)	37 (25.3%)	5 (11.4%)	93 (14.9%)
Public-Private	19 (6.4%)	6 (4.3%)	6 (4.1%)	0 (0.0%)	31 (5.0%)
(Missing)	7	7	0	2	16
Surgical emergency service					
Yes – surgery available 24 h	270 (91.2%)	121 (87.7%)	138 (94.5%)	36 (81.8%)	565 (90.5%)
Yes - surgery only during daytime	18 (6.1%)	9 (6.5%)	6 (4.1%)	4 (9.1%)	37 (5.9%)
Yes - assessment only	3 (1.0%)	3 (2.2%)	2 (1.4%)	2 (4.5%)	10 (1.6%)
No	5 (1.7%)	5 (3.6%)	0 (0.0%)	2 (4.5%)	12 (1.9%)
(Missing)	7	7	0	2	16

Data are n (%) unless stated otherwise. Denominators for each group are presented at the top row.

HIC: high-income country, UMIC: upper-middle-income country, LMIC: lower-middle-income country, LIC: low-income country.

Supplementary table 9: Key performance measures to evaluate access and quality.

	HIC	UMIC	LMIC	LIC	Total
ACCESS AND COVERAGE					
Total	n=9,808	n=3,366	n=3,948	n=936	n=18,058
Emergency surgery					
Yes	545 (5.6%)	268 (8.0%)	360 (9.1%)	114 (12.2%)	1287 (7.1%)
No	9252 (94.4%)	3098 (92.0%)	3586 (90.9%)	818 (87.8%)	16754 (92.9%)
(Missing)	11	0	2	4	17
Bowel resection					
Yes	105 (1.1%)	49 (1.5%)	106 (2.7%)	36 (3.9%)	296 (1.6%)
No	9693 (98.9%)	3316 (98.5%)	3839 (97.3%)	896 (96.1%)	17744 (98.3%)
(Missing)	10	1	3	4	18
Total	n=9,252	n=3,098	n=35,86	n=818	n=16,754
Waiting time to elective surgery					
Median (IQR)	9.8 months (4.5 to 20.7)	7.0 months (3.0 to 18.7)	6.0 months (2.2 to 17.2)	7.0 months (2.0 to 17.9)	8.0 months (3.3 to 20.0)
(Missing)	16	3	2	3	24
Waiting time before diagnosis					
Median (IQR)	4.0 months (1.4 to 11.1)	4.0 months (1.3 to 11.8)	4.0 months (1.2 to 12.0)	5.0 months (1.6 to 11.9)	4.0 months (1.4 to 11.7)
(Missing)	1	0	0	0	1
Waiting time after diagnosis					
Median (IQR)	4.5 months (2.0 to 9.8)	2.5 months (1.0 to 6.5)	2.0 months (0.5 to 5.8)	0.9 months (0.3 to 3.8)	3.2 months (1.2 to 8.4)
(Missing)	5	2	2	3	12
Waiting time after decision for surgery					
Median (IQR)	2.7 months (1.2 to 5.6)	1.0 months (0.3 to 2.1)	0.5 months (0.1 to 1.7)	0.2 months (0.1 to 0.7)	1.6 months (0.5 to 3.9)
(Missing)	3	0	0	3	6
QUALITY AND SAFETY					
Total	n=8,916	n=2,575	n=2,642	n=635	n=14,768
Mesh use					
Yes	8842 (99.2%)	2465 (95.7%)	2231 (84.4%)	457 (72.0%)	13995 (94.8%)
No	74 (0.8%)	110 (4.3%)	411 (15.6%)	178 (28.0%)	773 (5.2%)
(Missing)	0	0	0	0	0
Total	n=7,198	n=2,332	n=2,532	n=605	n=12,658
Day case adoption					
Yes	4464 (62.2%)	1064 (45.7%)	844 (33.3%)	368 (60.8%)	6740 (53.3%)
No	2709 (37.8%)	1264 (54.3%)	1688 (66.7%)	237 (39.2%)	5898 (46.7%)
(Missing)	16	4	0	0	20
Total	n=9,808	n=3,366	n=3,948	n=936	n=18,058
Postoperative complications					
Yes	1187 (12.1%)	446 (13.3%)	634 (16.1%)	148 (15.9%)	2415 (13.4%)
Clavien-Dindo I	885 (9.1%)	316 (9.4%)	493 (12.5%)	114 (12.2%)	1808 (10.0%)
Clavien-Dindo II	208 (2.1%)	83 (2.5%)	110 (2.8%)	26 (2.8%)	427 (2.4%)
Clavien-Dindo III	84 (0.9%)	40 (1.2%)	24 (0.6%)	4 (0.4%)	152 (0.8%)
Clavien-Dindo IV	7 (0.1%)	3 (0.1%)	2 (0.1%)	0 (0.0%)	12 (0.1%)
Clavien-Dindo V	3 (0.0%)	4 (0.1%)	5 (0.1%)	4 (0.4%)	16 (0.1%)
No	8592 (87.9%)	2917 (86.7%)	3311 (83.9%)	783 (84.1%)	15603 (86.6%)
(Missing)	29	3	3	5	40

The table shows the number of patients experiencing each outcome, part of the key performance measures, except for waiting times. The rates presented are unadjusted rates. Waiting times are presented with median and interquartile range.

HIC: high-income country, UMIC: upper-middle-income country, LMIC: lower-middle-income country, LIC: low-income country.

Supplementary table 10: Additional descriptive measures to evaluate access and quality.

	HIC (n=9,808)	UMIC (n=3,366)	LMIC (n=3,948)	LIC (n=936)	Total (n=18,058)
ACCESS					
Standardised patient pathways					
Yes	8054 (82.8%)	2045 (62.0%)	2035 (51.5%)	653 (71.9%)	12787 (71.5%)
No	1678 (17.2%)	1254 (38.0%)	1913 (48.5%)	255 (28.1%)	5100 (28.5%)
(Missing)	76	67	0	28	171
Waiting list management					
Yes	9079 (93.3%)	2977 (90.2%)	3218 (81.5%)	856 (94.3%)	16130 (90.2%)
No	653 (6.7%)	322 (9.8%)	730 (18.5%)	52 (5.7%)	1757 (9.8%)
(Missing)	76	67	0	28	171
Availability of day case surgical unit					
Yes	9055 (93.0%)	2637 (79.9%)	2984 (75.6%)	760 (83.7%)	15436 (86.3%)
No	677 (7.0%)	662 (20.1%)	964 (24.4%)	148 (16.3%)	2451 (13.7%)
(Missing)	76	67	0	28	171
Financing methods					
Insurance by the government	8818 (90.6%)	2420 (73.4%)	2411 (61.1%)	648 (71.4%)	14297 (79.9%)
Other insurance ¹	605 (6.2%)	594 (18.0%)	169 (4.3%)	5 (0.6%)	1373 (7.7%)
Out of pocket	97 (1.0%)	120 (3.6%)	1081 (27.4%)	189 (20.8%)	1487 (8.3%)
Other ²	212 (2.2%)	165 (5.0%)	287 (7.3%)	66 (7.3%)	730 (4.1%)
(Missing)	76	67	0	28	171
QUALITY					
Primary operator					
Senior surgeon	7181 (73.3%)	2469 (73.4%)	2530 (64.1%)	572 (61.4%)	12752 (70.7%)
Trainee surgeon	2605 (26.6%)	847 (25.2%)	1372 (34.8%)	349 (37.4%)	5173 (28.7%)
Non surgeon, medical practitioners	12 (0.1%)	50 (1.5%)	44 (1.1%)	11 (1.2%)	117 (0.6%)
(Missing)	10	0	2	4	16
Anaesthesia variation					
General	5867 (59.9%)	1920 (57.0%)	1551 (39.3%)	320 (34.3%)	9658 (53.5%)
Spinal	2633 (26.9%)	1228 (36.5%)	1890 (47.9%)	548 (58.8%)	6299 (34.9%)
Loco-regional	239 (2.4%)	83 (2.5%)	68 (1.7%)	21 (2.3%)	411 (2.3%)
Local	1059 (10.8%)	135 (4.0%)	433 (11.0%)	43 (4.6%)	1670 (9.3%)
(Missing)	10	0	4	4	18
Surgical approach variation					
Open	7075 (72.2%)	2606 (77.4%)	3513 (89.0%)	923 (99.0%)	14117 (78.2%)
MIS / MIS converted	2723 (27.8%)	759 (22.6%)	433 (11.0%)	9 (1.0%)	3924 (21.8%)
(Missing)	10	1	2	4	17


Data are n (%) unless stated otherwise. Denominators for each group are presented at the top row.

HIC: high-income country, UMIC: upper-middle-income country, LMIC: lower-middle-income country, LIC: low-income country,

¹ *Other insurance* included insurance provided by the employer, insurance that the patients has privately arranged and paid for and insurance but unknown how it was arranged.

² *Other* included external funds or grants awarded by charities and non-governmental organisations and other methods of payment not mentioned elsewhere.


Supplementary figure 1: Flowchart of included patients according to income groups and urgency of surgery.



*invalid data included records with incomplete dates, invalid age or inconsistent data that were identified and excluded during data cleaning.

HIC: high-income country, UMIC: upper-middle-income country, LMIC: lower-middle-income country, LIC: low-income country.

Supplementary figure 2: Global distribution of included patients



References:

1. Samuel JC, Tyson AF, Mabedi C, et al. Development of a ratio of emergent to total hernia repairs as a surgical capacity metric. *Int J Surg*. 2014;12(9):906-11. doi:10.1016/j.ijsu.2014.07.019
2. Kockerling F, Heine T, Adolf D, et al. Trends in Emergent Groin Hernia Repair-An Analysis From the Herniated Registry. *Front Surg*. 2021;8:655755. doi:10.3389/fsurg.2021.655755
3. van Ginneken E, Reed S, Siciliani L, et al. *Addressing backlogs and managing waiting lists during and beyond the COVID-19 pandemic*. 2022. *European Observatory Policy Briefs*.
4. HerniaSurge Group. International guidelines for groin hernia management. *Hernia*. Feb 2018;22(1):1-165. doi:10.1007/s10029-017-1668-x
5. World Health Organization. Patient S. *WHO guidelines for safe surgery 2009 : safe surgery saves lives*. World Health Organization, Patient Safety; 2009.
6. Bailey CR, Ahuja M, Bartholomew K, et al. Guidelines for day-case surgery 2019: Guidelines from the Association of Anaesthetists and the British Association of Day Surgery. *Anaesthesia*. Jun 2019;74(6):778-792. doi:10.1111/anae.14639