








CONTEMPORARY REVIEW

# Sex-Specific Considerations in Degenerative Aortic Stenosis for Female-Tailored Transfemoral Aortic Valve Implantation Management

Giulia Masiero , MD; Valeria Paradies, MD; Anna Franzone, MD; Barbara Bellini, MD; Chiara De Biase, MD; Nicole Karam , MD; Francesca Sanguineti, MD; Mamas A. Mamas , MD; H el ene Eltchaninoff, MD; Chiara Fraccaro , MD; Battistina Castiglioni , MD; Tiziana Attisano, MD; Giovanni Esposito , MD; Alaide Chieffo , MD

**ABSTRACT:** The impact of sex on pathophysiological processes, clinical presentation, treatment options, as well as outcomes of degenerative aortic stenosis remain poorly understood. Female patients are well represented in transfemoral aortic valve implantation (TAVI) trials and appear to derive favorable outcomes with TAVI. However, higher incidences of major bleeding, vascular complications, and stroke have been reported in women following TAVI. The anatomical characteristics and pathophysiological features of aortic stenosis in women might guide a tailored planning of the percutaneous approach. We highlight whether a sex-based TAVI management strategy might impact on clinical outcomes. This review aimed to evaluate the impact of sex from diagnosis to treatment of degenerative aortic stenosis, discussing the latest evidence on epidemiology, pathophysiology, clinical presentation, therapeutic options, and outcomes. Furthermore, we focused on technical sex-oriented considerations in TAVI including the preprocedural screening, device selection, implantation strategy, and postprocedural management.

**Key Words:** aortic stenosis ■ clinical management ■ device selection ■ sex differences ■ transcatheter aortic valve implantation

**D**egenerative aortic stenosis (AS) currently represents one of the most frequent nonrheumatic valvular heart diseases worldwide.<sup>1</sup> Nevertheless, the impact of sex on pathophysiology, clinical presentation, and outcomes in AS remains poorly defined.<sup>2</sup> Female patients are well represented in the transcatheter treatment of severe AS, both in national registries and landmark trials, and several lines of evidence suggest that women might experience greater benefit than men with transfemoral aortic valve implantation (TAVI) compared with surgical treatment.<sup>3–7</sup> In this review, we aimed to describe the impact of female sex on diagnosis and management of severe AS, discussing the latest evidence on epidemiology, pathophysiology, presentation, and treatment. The novelty of this review

is its focus on technical sex-oriented considerations ranging from preprocedural screening to periprocedural tricks and postprocedural management, including the latest technological innovations. Moreover, we aimed to report whether sex-based TAVI management protocols might impact on clinical outcomes.

## IMPACT OF FEMALE SEX IN THE DIAGNOSIS OF ACQUIRED AS

### Epidemiology, Pathophysiology, and Natural History

The prevalence of symptomatic severe AS is up to 4% in elderly patients (aged >75 years), with equal frequency

Correspondence to: Alaide Chieffo, MD, FESC, Interventional Cardiology Unit, IRCCS San Raffaele Scientific Institute, 60 Via Olgettina, 20132 Milan, Italy. Email: [chieffo.alaide@hsr.it](mailto:chieffo.alaide@hsr.it)

For Sources of Funding and Disclosures, see page 12.

  2022 The Authors. Published on behalf of the American Heart Association, Inc., by Wiley. This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

JAHA is available at: [www.ahajournals.org/journal/jaha](http://www.ahajournals.org/journal/jaha)

## Nonstandard Abbreviations and Acronyms

<b>AS</b>	aortic stenosis
<b>MDCT</b>	multidetector computed tomography
<b>TAVI</b>	transfemoral aortic valve implantation
<b>THV</b>	transcatheter heart valve

in men and women and a mortality rate of over 50% at 2 years of follow-up unless promptly treated.<sup>1,8</sup> The main cause of acquired AS involves a degenerative age-related valve mineralization, a dynamic process with lipid accumulation, chronic inflammation, and active valve leaflet calcification. However, little is known about the role of sex in the progression of valve disease and in the ventricular response to the pressure overload (Figure 1). In contrast to men, who develop a fibrotic dilated cardiomyopathy, left ventricular ejection fraction (LVEF) is less likely to deteriorate in women who experience a relatively greater wall thicknesses but smaller left ventricular (LV) cavities because of concentric hypertrophy. This remodeling process reverses more often in women shortly after aortic valve replacement.<sup>9</sup> The pathological explanation for such different remodeling patterns and the extent of ventricular fibrosis partly relies on a lower expression of profibrotic and inflammatory genes (eg, collagen I-II, matrix metalloproteinase 2, transforming growth factor- $\beta$  signaling pathways), which have been found in the interstitial cells of intraoperative myocardial biopsy specimens.<sup>10</sup> Differences in valve morphology have also been described, because female sex is associated with a lower degree of aortic valve calcification measured by multidetector computed tomography (MDCT) in patients with a similar degree of hemodynamic stenosis severity evaluated at Doppler echocardiography, even after adjustment for body surface area (BSA) and echocardiographic parameters.<sup>9</sup> The histological detection of higher levels of valvular fibrosis and dense connective tissue might explain why in women a more fibrotic aortic valve remodeling is observed for a given aortic stenosis severity. In contrast, male patients have histological evidence of more pronounced calcific valve remodeling irrespective of patients' age.<sup>11-13</sup> Conflicting data exist on the natural progression of AS according to sex. Some data suggest similar progression of aortic valve disease in men and women with respect to the gradient, velocity, or valve area; whereas other studies have reported that female sex is an independent predictor of mean aortic valve gradient progression, showing a significant association with AS-related event rates at long-term follow-up.<sup>14</sup>




## Clinical Presentation

The role of sex in the modulation of the pathological processes associated with the development of

AS translates into different clinical phenotypes and decision-making strategies (Figure 1).<sup>8,12,15</sup> Almost half of women with severe AS are asymptomatic and therefore diagnosed at later stages, with a lower rate of referral to treatment.<sup>16</sup> As symptoms appear, female patients are older and experience higher rates of symptomatic heart failure with shortness of breath along with dizziness and syncope.<sup>17</sup> The heavier symptomatic burden in the context of a chronic LV pressure overload may be explained by the relative higher LV wall thickness, smaller LV cavity, and LV filling pressures in women.<sup>18</sup> In contrast, men tend to have a higher prevalence of comorbidities, in particular coronary and peripheral artery disease and diabetes; in contrast, female patients show a greater prevalence of chronic kidney disease.<sup>14,17</sup>

## Diagnostic Assessment

Consistent sex-specific guideline criteria for the grading of disease severity are still lacking. Echocardiographic assessment warrants indexed values for BSA (aortic valve area index,  $<0.6 \text{ cm}^2/\text{m}^2$ ), but the same correction has not been applied for other hemodynamic parameters such as mean gradient or peak velocity.<sup>8</sup> Transthoracic echocardiogram evaluation is often challenging, because it may underestimate the LV outflow tract area and stroke volume, resulting in discrepancies between mean gradient and aortic valve area index.<sup>19</sup> In addition, up to 40% of patients present with a severe reduction of calculated aortic valve area index with low-gradient AS. The classical low flow–low gradient because of low LVEF is more prevalent in men, whereas a paradoxical low flow–low gradient disease, where LVEF is preserved but stroke volume index is  $<35 \text{ mL/m}^2$ , is more common in elderly women because of their small, restrictive LV cavity, greater arterial stiffness, and higher ventriculo-arterial impedance. This process may be found in 10% to 15% of the AS population and may lead to paradoxical underestimation of the severity of the disease,<sup>20</sup> leading to undertreatment of female patients. Although the transthoracic echocardiogram remains the standard diagnostic test, MDCT could provide more detailed quantification of AS severity and progression, using the true MDCT measured LV outflow tract area for reclassification of aortic valve area and calculation of the aortic valve calcium load that is strongly associated with worse morbidity and mortality.<sup>21,22</sup> Women require lower levels of aortic valve calcium load to reach hemodynamically severe AS compared with men, with sex-specific calcium scores for the diagnosis of AS.<sup>8,14,23</sup> Thresholds for severe AS assessed by means of computed tomography measurement of aortic valve calcification (Agatston units) are: men  $>3000$ , women  $>1600$ =highly likely; men  $>2000$ , women  $>1200$ =likely; men  $<1600$ , women  $<800$ =unlikely. Interestingly, this discordant calcific/fibrotic patterns between men and women may be

<b>Epidemiology</b> 	<b>Prevalence of severe AS</b>	Up to 4% between elderly patients (>75 years) ≅ frequency in men and women
	<b>Natural progression</b>	Conflicting results on disease progression High rate of asymptomatic severe AS but greater symptomatic burden Lower prevalence of comorbidities
<b>Pathophysiology</b> 	<b>Patterns of AV remodeling</b>	↓ extent of AVC load but comparable degree of valvular stenosis ↑ levels of valvular fibrosis and dense connective tissue
	<b>Patterns of LV hypertrophy</b>	↓ extent of ventricular fibrosis (↓ expression of pro-fibrotic and inflammatory genes) ↑ concentric hypertrophy ↑ wall thicknesses and LVEF ↓ LV cavities
<b>Diagnostic assessment</b> 	<b>Echocardiography</b>	LV concentric remodelling ↑ paradoxical LF-LG AS
	<b>MDCT</b>	Smaller anatomic root Lower coronary take off ↓ AVC threshold for severe AS Smaller ileo-femoral vessels and BSA
	<b>cMR</b>	↓ LV mass index ↑ myocardial fibrosis

**Figure 1. Epidemiological, pathophysiological, and diagnostic peculiarities in female patients affected by degenerative AS.** AS indicates aortic stenosis; AV, aortic valve; AVC, aortic valve calcification; BSA, body surface area; cMR, cardiac magnetic resonance; LF-LG, low flow–low gradient; LV, left ventricular; LVEF, left ventricular ejection fraction; and MDCT, multidetector computed tomography.

exacerbated in specific valve morphology such as stenotic bicuspid valve that shows less aortic valve calcification for the same hemodynamic severity not only in men but also compared with women with severe tricuspid aortic valve stenosis.<sup>13</sup> Lastly, cardiac magnetic resonance could provide complementary information on LV function, because it allows identification of different patterns of hypertrophy and remodeling and quantifies the extent of LV fibrosis in late gadolinium enhancement assessment.<sup>24</sup> However, data on cardiac magnetic resonance evaluation of sex differences in myocardial LV fibrosis are still limited. Nevertheless, the sex differences in the pathophysiological process and clinical presentation of severe AS might further benefit from an integrated diagnostic approach to avoid late referral and adverse outcomes encountered in female patients.

The introduction of TAVI flattened the sex-related gap in the treatment of AS; women represent >50% of patients undergoing TAVI.<sup>26</sup> Nevertheless, although female sex is a known risk factor for perioperative mortality in both the EuroSCORE and the Society of Thoracic Surgeons risk score, current international guidelines do not provide sex-specific recommendations on treatment.<sup>8,27</sup> The heart team plays a pivotal role in the selection of the optimal modality of intervention (surgical or transcatheter) for each individual patient, based on clinical, anatomical, and procedural features.<sup>8</sup> Factors favoring the choice of TAVI over surgical aortic valve replacement in female patients include advanced age at presentation, greater frailty, lower prevalence of concomitant severe coronary artery disease, and higher likelihood of patient–prosthesis mismatching smaller aortic annulus.<sup>8</sup>

## IMPACT OF FEMALE SEX IN THE TREATMENT OF AS

### Current Guidelines Recommendations and the Role of the Heart Team

Historically, female patients with AS have been turned down for surgical aortic valve replacement more often than men, mainly because of their late presentation and perceived greater risk of perioperative complications.<sup>25</sup>

### TAVI in Female Patients: Summary of Clinical Outcomes

Current data around TAVI outcomes in female patients compared with male patients are derived from subanalyses of randomized trials and observational studies (Table 1).<sup>4–7,25,28–37</sup>

Among comparative studies, in the combined cohorts of the PARTNER (Placement of Aortic


**Table 1. Studies Comparing the Main Clinical Outcomes After Transcatheter Aortic Valve Implantation in Female and Male Patients**

Study or first author and design	Overall population, n	% Women	Follow-up	Mean age, y		Operative risk, %		Vascular complications		Bleeding		All-cause mortality	
				Women	Men	Women	Men	Women	Men	Women	Men	Women	Men
FRANCE 2 registry, prospective observational <sup>28</sup>	3972	49.5	1 y	84.0±6.6	81±6.5	21.4±13*	22.2±15.2*	4.6	1.9	4.3	1.9	19.3	23.7
Kodali et al, analysis of RCT <sup>5</sup>	2559	47.6	30d	84.9±6.9	84.1±7.3	11.9±4.2	11.1±4.0	17.3	5.08	10.5	7.7	6.5	5.9
Szerlip et al, analysis of RCT <sup>4</sup>	1661	39.5	1 y	82.5±7.2	82.0±7.1	6.8±3.0	6.3±2.8	7.2	4.2	16.7	14.6	9.4	10.4
Sanmimo et al, retrospective observational <sup>29</sup>	910	46.5	1 y	82.0±7.6	80.9±8.4	7.9±3.7	7.1±4.0	7.8	4.1	4.0	1.6	7.0	12.7
Doshi et al, retrospective observational <sup>30</sup>	41050	47.7	In-hospital	81.7±8.1	80.5±9.1	...	...	3.9	3.3	28.3	20	4.7	3.9
Stehli et al, prospective observational <sup>31</sup>	683	58	30d	84.2±5.2	83.2±6.9	5.2±3.1	4.6±3.5	8.2	5.6	3.3	1	2.4	0.3
FRAILTY-AV, prospective observational <sup>25</sup>	759	44.8	1 y	84.2±5.3	82.8±5.8	6.5±3.8	5.8±4.4	...	...	...	...	19	17
TVT registry, prospective observational <sup>32</sup>	23652	49.9	1 y	82.3±8.5	81.7±8.6	9±6	8±6	8.3	4.4	8.0	5.9	21.3	24.5
Forrest et al, analysis of RCT <sup>6</sup>	3687	46.3	1 y	84.0±7.6	82.7±7.9	9.6±4.9	8.3±4.6	9.7	4.9	42.7	31.2	21.3	24.1
D'Ascenzo et al, retrospective observational <sup>33</sup>	377	57.2	30d	82.9±5.4	81.6±5.3	5.6±3.1	7.6±6.2	12.9	9.8	44	25	7.4	8.7
Italian Multicenter CoreValve registry, prospective observational <sup>34</sup>	659	55.8	1 y	82±5	80±7	23±14*	23±14*	2.2	1.7	3.5	2.7	16	19
Humphries et al, analysis of RCT <sup>7</sup>	641	51.3	30d	83	82	7.5	7.5	12.4	5.4	21.6	15.8	6.5	11.2
Czarnecki et al, retrospective observational <sup>35</sup>	999	45.3	1 y	85	83	7.0±5.0*	8.0±7.0*	18.7	16.7	14.5	12.6	18.2	19.2
Katz et al, prospective observational <sup>36</sup>	819	51	30d	82.4±7.0	80.6±7.5	15.4±13	11.4±9.5	11.2	5.5	7.4	6.2	11.5	6.5
Tarantini et al, prospective observational <sup>37</sup>	1694	49	30d	82.7±6.3	80.8±6.9	17.9±11.7*	17.8±13.9*	3.7	2.7	...	...	36†	39.7†

FRAILTY-AVR indicates Frailty Aortic Valve Replacement study; FRANCE 2, FRANCE 2, FRANCE 2, randomized control trial; TAVI, transcatheter aortic valve implantation; and TVT, Transcatheter Valve Therapy registry.

\*Logistic EuroScore; all the other risk scores are reported as Society of Thoracic Surgeons risk score.

†14-year results.




**Figure 2. Sex-tailored TAVI planning and periprocedural management.**

3D indicates 3 dimensional; InH, in hospital; LV, left ventricular; MDCT, multidetector computed tomography; PPM, prothesis–patient mismatch; SFAR, sheath-to-femoral artery ratio; TAVI, transcatheter aortic valve implantation; THV, transcatheter heart valve; and TTE, transthoracic echocardiogram.

Transcatheter Valves) II SAPIEN 3 (S3) (Edwards Lifesciences) trial, which included high- and intermediate-risk patients, female sex was independently associated with an increased risk of major vascular complications, but no differences in 30-day survival or stroke rate were observed in women compared with men.<sup>4</sup> In the Society of Thoracic Surgeons/American College of Cardiology Transcatheter Valve Therapy registry, vascular complications were more frequent among 11 808 women compared with 11 844 men (8% versus 4%; adjusted hazard ratio [HR], 1.7 [95% CI, 1.3–2.1];  $P < 0.001$ ), whereas mortality at 1 year was lower (21% versus 25%, adjusted HR, 0.7 [95% CI, 0.6–0.9];  $P < 0.001$ ).<sup>32</sup> Likewise, in 2 large meta-analyses, women experienced higher rates of major vascular complications, major bleeding events, and stroke than men.<sup>26,38</sup> A low BSA and smaller peripheral vessel diameter increases the risk of bleeding and vascular complications in the periprocedural phase. In this context, device innovation has the potential to mitigate the burden of access-site complications in women undergoing TAVI through a progressive reduction of the size and profile of delivery systems and the extended use of expandable sheaths.<sup>39</sup> On the other hand, female sex was associated with a survival advantage at long-term follow-up (mean follow-up length of  $3.3 \pm 1.1$  years).<sup>29</sup> A lower prevalence of comorbidities

in addition to the longer life expectancy enjoyed by women compared with men in the general population may drive the higher survival reported in the longer term. Conversely, an analysis of a contemporary cohort of patients treated with the SAPIEN 3 (Edwards Lifesciences, Irvine, CA) or Corevalve Evolut R or Evolut Pro (Medtronic, Minneapolis, MN) valves found similar rates of in-hospital mortality, stroke, and pacemaker implantation in women and men.<sup>40</sup>

The WIN TAVI (Women's International Transcatheter Aortic Valve Implantation) registry was the first international, multicenter, prospective observational registry of 1019 female patients undergoing TAVI at 18 European and 1 North American center between 2013 and 2015. Baseline clinical characteristics included a mean age of 83 years and mean LVEF of 56; ~90% of patients were considered at high risk for surgery. Transfemoral TAVI was performed in 91% of cases, with 42% use of second-generation valves. At 30-days, the primary VARC-2 (Valve Academic Research Consortium 2) early safety end point (composite of mortality, stroke, major vascular complications, life-threatening bleeding, stage 2 or 3 acute kidney injury, coronary artery obstruction, or repeat procedure for valve-related dysfunction) occurred in 14% and was mainly driven by vascular complications (7.7%) or bleeding events (4.4%). Increasing age, history of stroke, LVEF  $< 30\%$ ,



**Figure 3. Unfavorable anatomical correlation between THV frame, small aortic root, and coronary ostia, both in native and in bioprosthetic valve TAVI procedures.**

**A.** Possible anatomical relationship between a self-expandable supra-annular device and a small annulus and STJ, that may hamper selective coronary artery cannulation because of a high sealing skirt. **B.** Scenario of a TAVI-in-TAVI procedure in case of a small aortic root with a narrow distance between the THV and STJ (<2 mm). Whatever is the first THV (intra-annular or supra-annular), if its commissural plane is at the level or even above the STJ, the implantation of a second THV may entail the risk of coronary obstruction because of sinus sequestration.<sup>71,72</sup> STJ indicates sinotubular junction; TAVI, transcatheter aortic valve implantation; and THV, transcatheter heart valve.

and first-generation devices were found independent predictors of the primary end point.<sup>41</sup> At 1 year, a primary efficacy composite end point (including mortality, stroke, myocardial infarction, hospitalization for valve-related symptoms or heart failure or valve-related dysfunction) occurred in 167 (16.5%) patients, with nearly half of the rates previously reported in studies including first-generation devices; rates of all-cause mortality and stroke were 13% and 2%, respectively. No associations between history of pregnancy and 1-year outcomes were observed.<sup>42</sup>

Additional evidence is expected with the results of the RHEIA (Randomized Research in Women All Comers With Aortic Stenosis) trial, a prospective, randomized multicenter study that tests noninferiority and, eventually, the superiority of TAVI versus surgical aortic valve replacement for a primary end point composed of all-cause mortality, all stroke, and rehospitalization for valve or procedure-related symptoms or worsening congestive heart failure at 1 year after the procedure among >400 women with severe aortic stenosis (NCT04160130).<sup>43</sup>

## PROPOSED SEX-ORIENTED TAVI MANAGEMENT

The distinctive anatomical characteristics and pathophysiological features of AS in women affect outcomes after TAVI and should guide tailored TAVI planning, procedural strategy, and postprocedural management (Figure 2). Several measures and technical tricks can

be implemented during standard TAVI planning and the procedure to minimize adverse events in a sex-oriented approach, taking into account the longer life expectancy of female patients. All the proposed recommendations are integrated into a contemporary lean TAVI clinical pathway.<sup>44,45</sup>

## Preprocedural Screening and Device Selection

Currently, preprocedural screening of a patient candidate to TAVI is the same for women and men and consists of a multimodality imaging evaluation including transthoracic echocardiography, coronary angiography, MDCT, according to institutional protocols.<sup>46</sup> Among them, the gold standard imaging modality is MDCT, which can also play a role in screening for coronary artery disease.<sup>47</sup> As previously highlighted, women present more often with a small anatomic root (including small annulus, sinus of Valsalva, and sinotubular junction), low coronary ostia, and small iliofemoral vessels resulting in an unfavorable sheath-to-femoral artery ratio, together with a small BSA.<sup>48,49</sup> Accordingly, careful measurement of these parameters should be performed and compared with the anatomical requirements for the different available transcatheter heart valves (THVs) and their respective sizing charts to select the most appropriate THVs and vascular access in a patient-tailored fashion.<sup>50</sup>

First of all, it is well noted that currently available THVs cover different specific ranges of annulus size, with SAPIEN (Edwards Lifesciences), Corevalve Evolut

(Medtronic), Portico (Abbott Structural Heart, Santa Clara, CA), Allegra (New Valve Technology, Hechingen, Germany), and Myval (Meril Life Sciences, Gujarat, India) as those who are approved for the small annulus size (<20mm). To date, there are no randomized trial data with regard to the superiority of one device to others in the case of a small annulus size. The SMART (Small Annuli Randomized to Evolut or SAPIEN) trial, comparing the 2 most widely available contemporary TAVI devices, the Medtronic Evolut PRO/PRO+ self-expanding and the Edwards SAPIEN 3/SAPIEN 3 Ultra balloon-expandable devices in aortic valve annulus area of  $\leq 430\text{mm}^2$ , is still ongoing.<sup>51</sup> The SMART trial will be the largest trial to enroll primarily female patients.

Second, the presence of small annulus size (generally considered as an annulus size <23mm), one of the predictors of patient–prosthesis mismatch even after THV implantation, should be taken into account for device selection.<sup>52,53</sup> Patient–prosthesis mismatch after TAVI may be associated with less regression of LV hypertrophy, LV diastolic dysfunction, LV filling pressure, less improvement in LV systolic function, and less reduction of left atrial volume.<sup>54</sup> However, clinical data to demonstrate the impact of patient–prosthesis mismatch on long-term outcomes after TAVI are not available yet.<sup>52,55</sup> In practice, one may argue that, in case of a small annulus size, it is worthwhile to pursue the largest effective orifice area and the lowest transvalvular gradient through the use of a self-expandable supra-annular device. However, the sealing skirts of supra-annular devices are particularly high, and especially in the case of associated small sinotubular junction, may hamper selective coronary artery cannulation (Figure 3).<sup>56–58</sup>









Finally, the delivery sheath profile must be obviously considered in the choice of the most appropriate THV. As discussed above, in case of suboptimal vascular access (small sheath-to-femoral artery ratio, higher tortuosity, and diffuse and circumferential calcification), a device with a low-profile delivery system has to be selected. The CoreValve Evolut R and Pro+ (Medtronic) can be implanted using the InLine sheath with 14 Fr outer diameter equivalent (minimum vessel diameter requirements, 5.0mm for sizes 23, 26, 29mm, in the absence of circumferential calcification); also, the last generation low-profile delivery system FlexNav (Abbott Structural Heart, Santa Clara, CA) designed for the Portico and Navitor valve (Abbott Structural Heart), provides the same low insertion profile (access down to 5.0-mm vessels). Other devices require at least 5.5mm minimum vessel diameter or larger. If transfemoral access is not feasible, the heart team can consider different peripheral vascular approaches such as transsubclavian/transaxillary (both CoreValve and Portico iterations have a Conformité Européenne mark of approval for this vascular approach).

## Implantation Strategy and Periprocedural Considerations in Female Patients

### Risk of Vascular and Access-Related Complications

Because female sex seems to be independently associated with an almost double risk of vascular complications after TAVI, with a relative risk of 1.6 compared with men,<sup>26</sup> it is mandatory to adopt every possible technique to minimize this risk as per Figure 4:

1. Use of 2-dimensional ultrasound guidance during access site vascular puncture, because it has been associated with a significant reduction in the rate of major vascular complications and major bleeding over conventional fluoroscopic guidance.<sup>59</sup>
2. In case of heavy, circumferential calcifications, additional tools should help to increase patient eligibility and reduce vascular complications (ie, percutaneous transluminal angioplasty, paving and cracking endovascular techniques, debulking devices). Among them, intravascular lithotripsy is a recent promising technique in which ultrasound disrupts intimal and medial calcification, altering vessel compliance to allow for the safe passage of large-bore delivery sheaths.<sup>60</sup> This expands the patient cohort that could be eligible for transfemoral access, which still remains the first choice whenever possible and is associated with decreased morbidity and mortality compared with nontransfemoral access.<sup>61</sup> However, alternative routes may be considered (transsubclavian, transcarotid, transapical, transaortic, transcaval) in case of an unfeasible transfemoral route.
3. A minimalistic approach with a radial access to guide valve implantation or manage peripheral vascular complications and an over-the-wire rapid pacing reduce the secondary access site failures.<sup>62,63</sup> In case the secondary artery access is the contralateral femoral artery, the repositioning of a crossover guidewire may help in the management of any vascular complication should it occur.
4. The hemostasis technique is of utmost importance. A balloon inflation advanced from contralateral femoral artery to create a dry field during final hemostasis may be helpful for managing TAVI vascular access sites.<sup>64</sup> Most importantly, different sutured-based and collagen plug-based vascular closure devices are now available. The use of 2 Perclose Proglide (Abbott Vascular, Santa Clara, CA) devices was associated with a lower rate of major or minor vascular complications and lower rates of acute kidney injury in patients undergoing transfemoral TAVI in comparison to Prostar XL (Abbott Vascular).<sup>65</sup> Moreover, the plug-based Manta device (Teleflex, Wayne, PA), specifically designed for large-bore arteriotomy closure, recently showed reliable safety and

FEMALE-SPECIFIC ISSUE		PERI-PROCEDURAL TAVI CONSIDERATIONS	
Small SFAR 	↑ Vascular and bleeding complications		<input type="checkbox"/> Low-profile delivery system <input type="checkbox"/> Ultrasound guided vascular puncture <input type="checkbox"/> Hemostasis technique <input type="checkbox"/> Additional tools for safety <input type="checkbox"/> Aptimal management of antithrombotic therapy
Small aortic root 	PPM		<input type="checkbox"/> THV selection according to clinical/anatomical considerations
	Acute coronary obstruction/Aortic annulus rupture		<input type="checkbox"/> THV selection <input type="checkbox"/> Simulations of prostheses implantation <input type="checkbox"/> Coronary protection <input type="checkbox"/> Avoid unnecessary oversizing
Small and hypertrophic LV 	LV perforation		<input type="checkbox"/> Dedicated pre-shaped “stiff” wire with soft spiral
	Conduction disorders		<input type="checkbox"/> Tailored device implantation depth
Long life expectancy 	Life-time strategy		<input type="checkbox"/> THV choice and implantation strategy to optimize valve performance and future valve re-intervention

**Figure 4.** Periprocedural TAVI considerations to address the specific issues associated with the treatment of female patients.

LV indicates left ventricle; PPM, patient–prosthesis mismatch; SFAR, sheath-to-femoral artery ratio; TAVI, transcatheter aortic valve implantation; and THV, transcatheter heart valve.

efficacy in selected TAVI populations. Further registry data suggested that it was associated with faster hemostasis and a lower rate of vascular complications compared with the suture-based strategy, whereas early randomized studies highlighted a higher than anticipated rate of Manta-access vascular events (up to 19%).<sup>66,67</sup> It remains to be determined whether inherent differences in the design and operation of those devices may guide a sex-tailored choice.

5. The optimal management of antithrombotic therapy during the periprocedural phase may significantly impact on the rate of vascular complications. In particular, considering the frequent low body weight of female patients, it is recommended to administer a weight-based dose of unfractionated heparin with a close monitoring of activated clotting time. In line with the most recent scientific evidence and recommendations, dual antiplatelet therapy should be administered only in case of recent percutaneous coronary intervention.<sup>68</sup>

**Risk of Left Ventricle Perforation**

A small left ventricle cavity and an hypercontractile state, frequently encountered in female patients with AS, together with a thin muscular wall and a narrow aorto-mitral angle, are potential predictors of the occurrence of left ventricle perforation during TAVI.<sup>69</sup> Left

ventricle perforation primarily occurs secondary to the valve delivery wire inadvertently being pushed too distally through the left ventricle apex in a small cavity, even if the operators perform an appropriate curve on the distal stiff end of the wire.

It is suggested to use dedicated preshaped stiff wires with soft spiral tips to mitigate this risk. For example, the Amplatzer Extra-Stiff APEX wire (Cook Medical) has a double curve design composed of a larger curve with the distal tip of the wire forming a 2-mm J bend; it can be easily maneuvered in smaller ventricles, whereas the Safari2™ wire is available in 3 different loop sizes, offering the possibility to select the curve size according to the size of the ventricle<sup>70</sup> (Figure 4).

**Risk of Acute Coronary Occlusion**

Female sex is one of the strongest baseline clinical factors associated with the risk of acute post-TAVI coronary occlusion, a rare but life-threatening complication; in several observational experiences focusing on coronary obstruction following TAVI, the vast majority (>80%) of patients were women.<sup>71</sup> Coronary obstruction following TAVI is mainly caused by the displacement of the calcified native leaflet over the coronary ostia in the presence of lower coronary ostia and/or narrow aortic root, leaving little room to accommodate the native aortic leaflets, and as previously

Downloaded from http://ahajournals.org by on July 18, 2023



described, these are frequent anatomical features in female patients.<sup>48</sup> Moreover, in patients with prior surgical or percutaneous aortic valve treatment undergoing valve-in-valve or TAVI-in-TAVI procedures, the risk of acute coronary occlusion, either directly or indirectly by sequestering the sinus of Valsalva at the sinotubular junction, is up to 6-fold higher compared with TAVI for native aortic valve disease.<sup>72</sup> Many criteria for selecting patients at high risk for obstruction have been proposed, including native or bioprosthetic aortic leaflet length that extends above the coronary ostia or above the sinotubular junction, a coronary artery height  $\leq 10$  mm, a virtual valve to coronary artery/ to sinotubular junction distance, respectively, of  $\leq 4$  and  $\leq 2$  mm, degenerative stentless bioprosthetic valve, or with externally mounted leaflets.<sup>73</sup>

Along with a tailored THV choice considering future valve reintervention in a lifetime management strategy, several periprocedural measures are useful to mitigate the risk of acute coronary obstruction, such as (Figure 3 and 4):

1. Accurate MDCT preprocedural planning with virtual simulations of prostheses implantation (the usefulness of 3-dimensional printing models is under investigation in this field).
2. If pre-TAVI balloon valvuloplasty is required, simultaneous angiography to depict coronary patency or obstruction.
3. In case of the abovementioned clinical and anatomical parameters of risk, consider leaflet modification strategies (ie, BASILICA [Bioprosthetic or Native Aortic Scallop Intentional Laceration to Prevent Iatrogenic Coronary Artery Obstruction]) and/or different technical strategies for coronary patency protection (eg, guiding catheter placement, guidewire placement, prepositioned stent, ready to be deployed after TAVI in a chimney or snorkel fashion, or orthotopic stent strategy).<sup>73</sup>
4. Use of self-expandable valves with a concave shape nitinol frame at the level of coronary arteries (with mandatory commissural alignment during implantation) rather than a balloon expandable straight cobalto-chromium frame.<sup>71</sup>
5. Use of a repositionable/retrievable THV or a valve with different anchoring mechanism (paper clip-like anchorage mechanism at the level of the leaflets).<sup>74</sup>
6. Avoid unnecessary oversizing.

### **Risk of Paravalvular Leak and Aortic Annulus Rupture**

Again, in a small hypertrophic left ventricle, any pressure or volume residual overload can impact on symptoms relief and clinical outcome. Notwithstanding, female patients less frequently experience residual

paravalvular leak; smaller annular size, greater use of TAVI oversizing, lower aortic valve calcification burden, and higher achieved true cover index typically found in female patients undergoing TAVI are likely contributing factors to this finding.<sup>75</sup> Conversely, balloon-expandable TAVI oversizing  $\geq 20\%$  of the annular area is associated with an 8-fold increased risk of annulus rupture, and this is more frequently described in women than men.<sup>76</sup>

1. It is generally recommended to optimize the procedural result by an accurate evaluation of valve hemodynamics in the catheterization laboratory (residual gradients and regurgitation). It is advisable to assess invasively any residual transprosthetic gradient.
2. Intra- or paravalvular leak has to be accurately detected by multimodality evaluation (including supraortic angiogram, intraprocedural echocardiogram evaluation, aortic and left ventricle invasive pressure measurements, aortic regurgitation index) to try to optimize the result by means of adjunctive maneuvers (ie, after dilatation).<sup>77</sup> The use of the dirotic notch index to assess for paravalvular leak avoiding left ventricle measurements is currently under investigation.<sup>44</sup>

### **Risk of Conduction Disorders and Permanent Pacemaker**

A left bundle-branch block and the dyssynchrony lead by pacing stimulation are badly tolerated in case of a small hypertrophic left ventricle.

1. A valve implant as high as possible may help to reduce the risk of conduction disorders<sup>78</sup> (Figure 4).
2. If a pacemaker is needed, the most physiological stimulation modality should be pursued, and atrial contribution should be preserved whenever possible. Within this perspective, the measurement of membranous septum length by computed tomography scan can provide a patient-specific approach to device implantation depth, because implanting the device at a depth less than the membranous septum length helps to significantly reduce permanent pacemaker rate.<sup>79</sup>

### **Lifetime Management Strategy**

Considering the longer life expectancy of women, it is particularly advisable to tailor THV choice and implantation strategy to optimize valve performance and foresee the strategy of future valve reintervention<sup>80</sup> (Figure 4). Different scenarios may arise:

1. In the case of a small sinus of Valsalva and narrow sinotubular junction, the priority should be

to preserve the easiest coronary reaccess possible, with consideration of future valve-in-valve. Accordingly, in case of short-frame THV device selection, a slightly deeper implantation into the LV outflow tract can be attempted (for example, 60:40 instead of 70:30 or 80:20) to keep the neocommissural plane below the coronary ostia plane. On the other hand, if a long-frame device is implanted, implantation strategies aiming to obtain commissural alignment should be implemented.<sup>81</sup>

2. In case of elderly patients with a larger BSA, a supra-annular device may be a good option, particularly when using commissural alignment strategies during implantation. On the other hand, in patients with known associated coronary artery disease and/or longer life expectancy (entailing an increased risk of further procedures of TAVI-in-TAVI or percutaneous coronary intervention), a balloon expandable intra-annular short frame device could be preferred, especially if implanted in a subcoronary position.<sup>82-84</sup>

The enrolling SMART trial may help to address this issue.<sup>51</sup>

## POSTPROCEDURAL MANAGEMENT

Contemporary TAVI programs have to streamline post-procedural care and discharge policy, with no distinction between sexes.<sup>44,45</sup> However, in light of the greater simplification of the TAVI procedure and shortened length of hospital stays, specific attention should be paid to the risk of complications as well as to a tailored pharmacotherapy in female patients.

The increasing clinical experience and advances in device technology have led in many centers to a paradigm shift from general anesthesia to local anesthesia or conscious sedation followed by early discharge (within 48 hours), irrespective of surgical risk.<sup>85,86</sup> Recent observational data support the benefit of a minimalist approach with reduced procedure time and shorter hospitalization, where next-day discharge will become standard practice, resulting in cost-effective quality care.<sup>44,45</sup> A reduced risk of nosocomial infections and increased chance of being discharged home is highly appealing in women who are generally older, with less comorbidities, and present later in their disease trajectory to TAVI services.<sup>15,87</sup>

Notwithstanding, the length of hospitalization after a TAVI procedure is a delicate balance between benefit of early discharge and timely detection of post-procedural complications and should be tailored to individual patient's medical needs. Advanced age, low BSA, and small vessels make women at higher risk for bleeding and vascular complications in the early period after the procedure. However, careful access-site monitoring and serial hemoglobin measurements

within 48 hours allow identification of the majority of vascular access site-related complications before discharge. The FAST TAVI (Feasibility and Safety of Early Discharge After Transfemoral Transcatheter Aortic Valve Implantation) registry demonstrated that prespecified risk criteria based on the rate of 30-day complications could identify patients (48% women) for whom early discharge was safe and effective.<sup>88</sup> Among others, stable hemoglobin values and preserved renal function are pivotal risk evaluation criteria for early discharge and should be attentively examined in women who may have greater prevalence of chronic kidney disease and anemia.<sup>17,89</sup>

The optimal periprocedural antithrombotic regimen to prevent thrombotic complications and minimize the risk of bleeding is also important.<sup>90-93</sup> Several randomized trials have investigated the safety and efficacy of different antithrombotic regimens after TAVI (Table 2). However, no specific recommendations have been indicated for women undergoing TAVI procedures. European guidelines currently recommend life-long single antiplatelet therapy or oral anticoagulation in case of patients who have other indications for oral anticoagulation (antiplatelet therapy); dual antiplatelet therapy should be administered only in cases of recent percutaneous coronary intervention.<sup>8,68,93,94</sup> Female sex representation in major randomized trials investigating antithrombotic therapies after TAVI is almost 50% on average; however, no sex-specific subanalyses have been reported to date (Table 2).<sup>95-101</sup> Observational studies described sex disparities in antiplatelet and antithrombotic management after TAVI with a higher adoption of clopidogrel and a lower use of warfarin because of lower rates of atrial fibrillation.<sup>40</sup> Two large meta-analyses, investigating TAVI outcomes by sex, reported higher rates of major bleeding, vascular complications, and stroke in women, with no impact on survival.<sup>26,40</sup> Moreover, female sex is an independent predictor of anemia in patients with severe aortic stenosis. The WIN TAVI registry showed that not only is anemia a common finding in elderly women, but also strongly correlates to the long-term prognosis.<sup>42</sup> The sex-based assessment of complications and of pharmacokinetic and pharmacodynamic responses to antithrombotic medications are essential in the periprocedural management of women undergoing TAVI. In case of oral anticoagulation therapy, particular attention has to be paid to hemoglobin values, and an accurate reevaluation of the thrombotic and bleeding risks might be suggested during follow-up. More evidence in the context of large clinical trials investigating the safety and efficacy balance of different antithrombotic strategies according to sex are needed to guide therapeutic decision making in daily clinical practice.

**Table 2. Overview of Randomized Trials Investigating Antithrombotic Therapies After TAVI**

Study	Design	n	Women, %	Treatment strategy	Main findings
Ussia et al <sup>95</sup>	Single-center RCT	79	54.0	3mo DAPT (ASA+C) followed by ASA alone vs ASA	No difference between DAPT vs ASA at 30 d (13% vs 15%; $P=0.71$ ) and 6mo (18% vs 15%; $P=0.85$ ) in composite of death, MI, stroke, life-threatening bleeding, or urgent conversion to surgery.
ARTE <sup>96</sup>	Multicenter RCT	222	36.9	3mo DAPT (ASA+C) vs ASA	The composite of death, MI, stroke or transient ischemic attack, or major or life-threatening bleeding tended to occur more frequently in the DAPT group (15.3% vs 7.2%, $P=0.065$ ) within 3mo after TAVI.
POPULAR TAVI (Cohort A) <sup>97</sup>	Multicenter RCT	665	48.7	3mo DAPT (ASA+C) followed by ASA alone vs ASA	12mo all bleeding: 15.1% with ASA vs 26.6% with DAPT (RR, 0.57 [95% CI, 0.42 to 0.77]; $P=0.001$ ); 12mo composite of death from cardiovascular causes, non-procedure-related bleeding, stroke, or MI: 23.0% with ASA vs 31.1% with DAPT (difference, -8.2 percentage points [95% CI for noninferiority, -14.9 to -1.5]; $P<0.001$ ; RR, 0.74 [95% CI for superiority, 0.57 to 0.95]; $P=0.04$ ).
POPULAR TAVI (Cohort B) <sup>98</sup>	Multicenter RCT	313	45.4	3mo OAC+C followed by OAC alone vs OAC	12m all bleeding: 21.7% with OAC vs 34.6% with OAC+C (RR, 0.63 [95% CI, 0.43 to 0.90]; $P=0.01$ ); 12mo composite of death from cardiovascular causes, non-procedure-related bleeding, stroke, or MI: 31.2% with OAC vs 45.5% with OAC+C (difference, -14.3 percentage points [95% CI for noninferiority, -25.0 to -3.6]; RR, 0.69 [95% CI for superiority, 0.51 to 0.92]).
GALILEO <sup>99</sup>	Multicenter RCT	1644	49.5	3mo rivaroxaban 10mg/qd+ASA vs 3mo DAPT (ASA+C)	Primary efficacy outcome composite of death or TE: 9.8 with rivaroxaban and 7.2 with DAPT per 100 person-years (HR, 1.35 [95% CI, 1.01 to 1.81]; $P=0.04$ ). Primary safety outcome composite of major disabling or life-threatening bleeding: 4.3 with rivaroxaban and 2.8 with DAPT per 100 person-years (HR, 1.50 [95% CI, 0.95 to 2.37]; $P=0.08$ ).
ENVISAGE TAVI AF <sup>100</sup>	Multicenter RCT	1426	47.5	Edoxaban 60mg/d±antiplatelet therapy vs VKA±antiplatelet therapy	Primary efficacy outcome composite of death from any cause, MI, ischemic stroke, TE, valve thrombosis, or major bleeding: 17.3 with edoxaban and 16.5 with VKA per 100 person-years (HR, 1.05 [95% CI, 0.85 to 1.31]; $P=0.01$ for noninferiority). Major bleeding: 9.7 with edoxaban and 7.0 with VKA per 100 person-years (HR, 1.40 [95% CI, 1.03 to 1.91]; $P=0.93$ for noninferiority)
ATLANTIS <sup>101</sup>	Multicenter RCT	1510	53.0	Apixaban 5mg BID vs single/dual APT or VKA	Primary outcome, time to death, stroke, MI, systemic emboli, intracardiac or valve thrombosis, deep vein thrombosis/pulmonary embolism, or major bleeding for apixaban vs standard of care: 18.4% vs 20.1% (HR, 0.92 [95% CI, 0.73 to 1.16]) (stratum 1: apixaban vs VKA: 21.9% vs 21.9%, stratum 2: 16.9% vs 19.3%; $P$ for interaction=0.57). Primary safety end point of life-threatening, disabling, or major bleeding 8.5% vs 8.5%.

APT indicates antiplatelet therapy; ARTE, Aspirin vs Aspirin + Clopidogrel Following Transcatheter Aortic Valve Implantation; ASA, aspirin; ATLANTIS, Anti-Thrombotic Strategy After Trans-Aortic Valve Implantation for Aortic Stenosis; C, clopidogrel; DAPT, dual antiplatelet therapy; ENVISAGE-TAVI AF, Edoxaban vs Standard of Care and Their Effects on Clinical Outcomes in Patients Having Undergone Transcatheter Aortic Valve Implantation—Atrial Fibrillation; GALILEO, Global Study Comparing a Rivaroxaban-based Antithrombotic Strategy to an Antiplatelet-based Strategy after Transcatheter Aortic Valve Replacement to Optimize Clinical Outcomes; HR, hazard ratio; MI, myocardial infarction; OAC, oral anticoagulant; POPULAR TAVI, Antiplatelet Therapy for Patients Undergoing Transcatheter Aortic-Valve Implantation; RCT, randomized controlled trial; RR, risk ratio; TAVI, transcatheter aortic valve implantation; TE, thromboembolic events; and VKA, vitamin K antagonist.

## CONCLUSIONS

Female patients represent half of the patients undergoing TAVI in clinical studies; however, sex subanalyses are limited, and the impact of sex on outcomes has been poorly investigated. Currently, the unique anatomical characteristics and pathophysiological features of AS in female patients should guide tailored TAVI planning and periprocedural management. Further prospective studies focused around optimization of the therapeutic management of women with severe AS are warranted.

## ARTICLE INFORMATION

### Affiliations

Department of Cardiac, Thoracic, Vascular Science and Public Health, University of Padova, Italy (G.M., C.F.); Department of Cardiology, Maasstad Hospital, Rotterdam, The Netherlands (V.P.); Department of Advanced Biomedical Sciences, Federico II University of Naples, Italy (A.F., G.E.); Interventional Cardiology Unit, IRCCS San Raffaele Scientific Institute, Milan, Italy (B.B., A.C.); Groupe Cardiovasculaire Interventionnel, Clinique Pasteur, Toulouse, France (C.D.B.); Cardiology Department, European Hospital Georges Pompidou, Paris, France (N.K.); Institut Cardiovasculaire Paris Sud, Hôpital Privé Jacques CARTIER, Massy, France (F.S.); Keele Cardiovascular Research Group, School of Medicine, Keele University, Stoke-on-Trent, United Kingdom (M.A.M.); Department of Cardiology, Royal Stoke University Hospital, Stoke-on-Trent, United Kingdom (M.A.M.); Department of Medicine, Thomas Jefferson University, Philadelphia, PA (M.A.M.); Cardiology Department, Rouen University Hospital, Rouen, France (H.E.); Cardiothoracovascular Department, Ospedale di Circolo, ASST Sette Laghi, Varese, Italy (B.C.); and Division of Interventional Cardiology, Cardiovascular and Thoracic Department, San Giovanni di Dio e Ruggi, Salerno, Italy (T.A.).

### Sources of Funding

This study was partially supported by the Italian Society of Interventional Cardiology (Società Italiana di Cardiologia Interventistica–Gruppo Italiano di Studi Emodinamici), which also supported the publication charge.

### Disclosures

Dr Chieffo declares consultant/speaker fees from Abbott Vascular, Abiomed, Boston Scientific, and Edwards. The remaining authors have no disclosures to report.

## REFERENCES

1. Yadgir S, Johnson CO, Aboyans V, Adebayo OM, Adedoyin RA, Afarideh M, Alahdab F, Alashi A, Alipour V, Arabloo J, et al. Global, regional, and national burden of calcific aortic valve and degenerative mitral valve diseases, 1990–2017. *Circulation*. 2020;141:1670–1680. doi: 10.1161/CIRCULATIONAHA.119.043391
2. Garcia M, Mulvagh SL, Merz CN, Buring JE, Manson JE. Cardiovascular disease in women: clinical perspectives. *Circ Res*. 2016;118:1273–1293. doi: 10.1161/CIRCRESAHA.116.307547
3. Calabrò P, Niccoli G, Gragnano F, Grove EL, Vergallo R, Mikhailidis DP, Patti G, Spaccarotella C, Katsiki N, Masiero G, et al. Are we ready for a gender-specific approach in interventional cardiology? *Int J Cardiol*. 2019;286:226–233. doi: 10.1016/j.ijcard.2018.11.022
4. Szerlip M, Gualano S, Holper E, Squiers JJ, White JM, Doshi D, Williams MR, Hahn RT, Webb JG, Svensson LG, et al. Sex-specific outcomes of transcatheter aortic valve replacement with the sapien 3 valve: insights from the PARTNER II S3 high-risk and intermediate-risk cohorts. *JACC Cardiovasc Interv*. 2018;11:13–20. doi: 10.1016/j.jcin.2017.09.035
5. Kodali S, Williams MR, Doshi D, Hahn RT, Humphries KH, Nkomo VT, Cohen DJ, Douglas PS, Mack M, Xu K, et al. Sex-specific differences at presentation and outcomes among patients undergoing transcatheter aortic valve replacement: a cohort study. *Ann Intern Med*. 2016;164:377–384. doi: 10.7326/M15-0121
6. Forrest JK, Adams DH, Popma JJ, Reardon MJ, Deeb GM, Yakubov SJ, Hermiller JB Jr, Huang J, Skelding KA, Lansky A. Transcatheter aortic valve replacement in women versus men (from the US CoreValve Trials). *Am J Cardiol*. 2016;118:396–402. doi: 10.1016/j.amjcard.2016.05.013
7. Humphries KH, Toggweiler S, Rodés-Cabau J, Nombela-Franco L, Dumont E, Wood DA, Willson AB, Binder RK, Freeman M, Lee MK, et al. Sex differences in mortality after transcatheter aortic valve replacement for severe aortic stenosis. *J Am Coll Cardiol*. 2012;60:882–886. doi: 10.1016/j.jacc.2012.06.019
8. Vahanian A, Beyersdorf F, Praz F, Milojevic M, Baldus S, Bauersachs J, Capodanno D, Conradi L, De Bonis M, De Paulis R, et al.; ESC/EACTS Scientific Document Group, ESC National Cardiac Societies. 2021 ESC/EACTS Guidelines for the management of valvular heart disease. *Eur Heart J* 2022; 43:561–632. doi: 10.1093/eurheartj/ehab395
9. Shames S, Gillam LD. Sex differences in aortic valve calcification. *Circ Cardiovasc Imaging*. 2013;6:8–10. doi: 10.1161/jcimg.2012.02.034
10. Karargas G, Dworzak E, Petrov G, Summer H, Schulze TM, Baczko I, Knosalla C, Golz S, Hetzer R, Regitz-Zagrosek V. Sex dependent regulation of fibrosis and inflammation in human left ventricular remodeling under pressure overload. *Eur J Heart Fail*. 2014;16:1160–1167. doi: 10.1002/ejhf.171
11. DesJardin JT, Chikwe J, Hahn RT, Hung JW, Delling FN. Sex differences and similarities in valvular heart disease. *Circ Res*. 2022;130:455–473. doi: 10.1161/CIRCRESAHA.121.319914
12. Summerhill VI, Moschetta D, Orekhov AN, Poggio P, Myasoedova VA. Sex-specific features of calcific aortic valve disease. *Int J Mol Sci*. 2020;21:5620. doi: 10.3390/ijms21165620
13. Voisine M, Hervault M, Shen M, Boilard AJ, Filion B, Rosa M, Bossé Y, Mathieu P, Côté N, Clavel MA. Age, sex, and valve phenotype differences in fibro-calcific remodeling of calcified aortic valve. *J Am Heart Assoc*. 2020;9:e015610. doi: 10.1161/JAHA.119.015610
14. Chandrasekhar J, Dangas G, Mehran R. Valvular heart disease in women, differential remodeling, and response to new therapies. *Curr Treat Options Cardiovasc Med*. 2017;19:74. doi: 10.1007/s11936-017-0573-z
15. Tribouilloy C, Bohbot Y, Rusinaru D, Belkhir K, Diouf M, Altes A, Delpierre Q, Serbout S, Kubala M, Levy F, et al. Excess mortality and undertreatment of women with severe aortic stenosis. *J Am Heart Assoc*. 2021;10:e018816. doi: 10.1161/JAHA.120.018816
16. Gèneux P, Stone GW, O'Gara PT, Marquis-Gravel G, Redfors B, Giustino G, Pibarot P, Bax JJ, Bonow RO, Leon MB. Natural history, diagnostic approaches, and therapeutic strategies for patients with asymptomatic severe aortic stenosis. *J Am Coll Cardiol*. 2016;67:2263–2288. doi: 10.1016/j.jacc.2016.02.057
17. Steeds RP, Messika-Zeitoun D, Thambyrajah J, Serra A, Schulz E, Maly J, Aiello M, Rudolph TK, Lloyd G, Bortone AS, et al. IMPULSE: the impact of gender on the presentation and management of aortic stenosis across Europe. *Open Heart*. 2021;8:e001443. doi: 10.1136/openhrt-2020-001443
18. Singh A, Chan DCS, Greenwood JP, Dawson DK, Sonecki P, Hogrefe K, Kelly DJ, Dhakshinamurthy V, Lang CC, Khoo JP, et al. Symptom onset in aortic stenosis: relation to sex differences in left ventricular remodeling. *JACC Cardiovasc Imaging*. 2019;12:96–105. doi: 10.1016/j.jcmg.2017.09.019
19. Chin CW, Khaw HJ, Luo E, Tan S, White AC, Newby DE, Dweck MR. Echocardiography underestimates stroke volume and aortic valve area: implications for patients with small-area low-gradient aortic stenosis. *Can J Cardiol*. 2014;30:1064–1072. doi: 10.1016/j.cjca.2014.04.021
20. Pibarot P, Dumesnil JG. Low-flow, low-gradient aortic stenosis with normal and depressed left ventricular ejection fraction. *J Am Coll Cardiol*. 2012;60:1845–1853. doi: 10.1016/j.jacc.2012.06.051
21. Aggarwal SR, Clavel MA, Messika-Zeitoun D, Cuffe C, Malouf J, Araoz PA, Mankad R, Michelena H, Vahanian A, Enriquez-Sarano M. Sex differences in aortic valve calcification measured by multidetector computed tomography in aortic stenosis. *Circ Cardiovasc Imaging*. 2013;6:40–47. doi: 10.1161/CIRCIMAGING.112.980052
22. Tuncay V, Prakken N, van Ooijen PM, Budde RP, Leiner T, Oudkerk M. Semiautomatic, quantitative measurement of aortic valve area using CTA: validation and comparison with transthoracic echocardiography. *Biomed Res Int* 2015; 2015:648283, 1, 9. doi: 10.1155/2015/648283
23. Clavel MA, Messika-Zeitoun D, Pibarot P, Aggarwal SR, Malouf J, Araoz PA, Michelena H, Cuffe C, Larose E, Capoulade R, et al. The

- complex nature of discordant severe calcified aortic valve disease grading: new insights from combined Doppler echocardiographic and computed tomographic study. *J Am Coll Cardiol*. 2013;62:2329–2338. doi: 10.1016/j.jacc.2013.08.1621
24. Tastet L, Kwieciniski J, Pibarot P, Capoulade R, Everett RJ, Newby DE, Shen M, Guzzetti E, Arsenault M, Bédard É, et al. Sex-related differences in the extent of myocardial fibrosis in patients with aortic valve stenosis. *JACC Cardiovasc Imaging*. 2020;13:699–711. doi: 10.1016/j.jcmg.2019.06.014
  25. Pighi M, Piazza N, Martucci G, Lachapelle K, Perrault LP, Asgar AW, Lauck S, Webb JG, Popma JJ, Kim DH, et al. Sex-specific determinants of outcomes after transcatheter aortic valve replacement. *Circ Cardiovasc Qual Outcomes*. 2019;12:e005363. doi: 10.1161/CIRCOUTCOMES.118.005363
  26. O'Connor SA, Morice MC, Gilard M, Leon MB, Webb JG, Dvir D, Rodés-Cabau J, Tamburino C, Capodanno D, D'Ascenzo F, et al. Revisiting sex equality with transcatheter aortic valve replacement outcomes: a collaborative, patient-level meta-analysis of 11,310 patients. *J Am Coll Cardiol*. 2015;66:221–228. doi: 10.1016/j.jacc.2015.05.024
  27. Ad N, Barnett SD, Speir AM. The performance of the EuroSCORE and the Society of Thoracic Surgeons mortality risk score: the gender factor. *Interact Cardiovasc Thorac Surg*. 2007;6:192–195. doi: 10.1510/icvts.2006.138313
  28. Bière L, Launay M, Pinaud F, Hamel JF, Eltchaninoff H, Lung B, Laskar M, Leguerrier A, Gilard M, Furber A. Influence of sex on mortality and perioperative outcomes in patients undergoing TAVR: insights from the FRANCE 2 registry. *J Am Coll Cardiol*. 2015;65:755–757. doi: 10.1016/j.jacc.2014.11.044
  29. Sannino A, Szerlip M, Harrington K, Schiattarella GG, Grayburn PA. Comparison of baseline characteristics and outcomes in men versus women with aortic stenosis undergoing transcatheter aortic valve implantation. *Am J Cardiol*. 2018;121:844–849. doi: 10.1016/j.amjcard.2017.12.020
  30. Doshi R, Shlofmitz E, Meraj P. Comparison of outcomes and complications of transcatheter aortic valve implantation in women versus men (from the National Inpatient Sample). *Am J Cardiol*. 2018;121:73–77. doi: 10.1016/j.amjcard.2017.09.015
  31. Stehli J, Dagan M, Zaman S, Koh JQS, Quine E, Gouskova N, Crawford C, Dong M, Nanayakkara S, Htun NM, et al. Impact of gender on transcatheter aortic valve implantation outcomes. *Am J Cardiol*. 2020;133:98–104. doi: 10.1016/j.amjcard.2020.07.052
  32. Chandrasekhar J, Dangas G, Yu J, Vemulapalli S, Suchindran S, Vora AN, Baber U, Mehran R, STS/ACC TVT Registry. Sex-based differences in outcomes with transcatheter aortic valve therapy: TVT Registry From 2011 to 2014. *J Am Coll Cardiol*. 2016;68:2733–2744. doi: 10.1016/j.jacc.2016.10.041
  33. D'Ascenzo F, Gonella A, Moretti C, Omedè P, Salizzoni S, La Torre M, Giordana F, Barbanti M, Ussia GP, Brambilla N, et al. Gender differences in patients undergoing TAVI: a multicentre study. *EuroIntervention*. 2013;9:367–372. doi: 10.4244/EIJV9I3A59
  34. Buja P, Napodano M, Tamburino C, Petronio AS, Ettori F, Santoro G, Ussia GP, Klugmann S, Bedogni F, Ramondo A, et al. Comparison of variables in men versus women undergoing transcatheter aortic valve implantation for severe aortic stenosis (from Italian Multicenter CoreValve registry). *Am J Cardiol*. 2013;111:88–93. doi: 10.1016/j.amjcard.2012.08.051
  35. Czarnecki A, Qiu F, Koh M, Prasad TJ, Cantor WJ, Cheema AN, Chu MWA, Feindel C, Fremes SE, Kingsbury K, et al. Clinical outcomes after trans-catheter aortic valve replacement in men and women in Ontario, Canada. *Catheter Cardiovasc Interv*. 2017;90:486–494. doi: 10.1002/ccd.26906
  36. Katz M, Carlos Bacelar Nunes Filho A, Caixeta A, Antonio Carvalho L, Sarmiento-Leite R, Alves Lemos Neto P, Eduardo Koenig São Thiago L, Dias Dourado Oliveira A, Antonio Marino M, Tadeu Tumelero R, et al.; Brazilian TAVI Registry investigators. Gender-related differences on short- and long-term outcomes of patients undergoing transcatheter aortic valve implantation. *Catheter Cardiovasc Interv*. 2017; 89:429–436. doi: 10.1002/ccd.26658
  37. Tarantini G, Baumgartner H, Frank D, Husser O, Bleiziffer S, Rudolph T, Jeger R, Fraccaro C, Hovorka T, Wendler O. Four-year mortality in women and men after transfemoral transcatheter aortic valve implantation using the SAPIEN 3. *Catheter Cardiovasc Interv*. 2021;97:876–884. doi: 10.1002/ccd.29257
  38. Saad M, Nairooz R, Pothineni NVK, Almomani A, Kovelamudi S, Sardar P, Katz M, Abdel-Wahab M, Bangalore S, Kleiman NS, et al. Long-term outcomes with transcatheter aortic valve replacement in women compared with men: evidence from a meta-analysis. *JACC Cardiovasc Interv*. 2018;11:24–35. doi: 10.1016/j.jcin.2017.08.015
  39. Glaser N, O'Sullivan CJ, Saleh N, Verouhis D, Settergren M, Linder R, Rück A. Transcatheter aortic valve replacement using the iS-leeve expandable sheath in small femoral arteries. *Open Heart*. 2021;8:e001703. doi: 10.1136/openhrt-2021-001703
  40. Wang TY, Gracia E, Callahan S, Bilfinger T, Tannous H, Pyo R, Kort S, Skopicki H, Weinstein J, Patel N, et al. Gender disparities in management and outcomes following transcatheter aortic valve implantation with newer generation transcatheter valves. *Am J Cardiol*. 2019;123:1489–1493. doi: 10.1016/j.amjcard.2019.01.048
  41. Chieffo A, Petronio AS, Mehilli J, Chandrasekhar J, Sartori S, Lefèvre T, Presbitero P, Capranzano P, Tchetché D, Iadanza A, et al. Acute and 30-day outcomes in women after TAVR: results from the WIN-TAVI (Women's International Transcatheter Aortic Valve Implantation) Real-World Registry. *JACC Cardiovasc Interv*. 2016;9:1589–1600. doi: 10.1016/j.jcin.2016.05.015
  42. Chieffo A, Petronio AS, Mehilli J, Chandrasekhar J, Sartori S, Lefèvre T, Presbitero P, Capranzano P, Tchetché D, Iadanza A, et al. 1-year clinical outcomes in women after transcatheter aortic valve replacement: results from the first WIN-TAVI registry. *JACC Cardiovasc Interv*. 2018;11:1–12. doi: 10.1016/j.jcin.2017.09.034
  43. Eltchaninoff H, Bonaros N, Prendergast B, Nietlispach F, Vasa-Nicotera M, Chieffo A, Pibarot P, Bramlage P, Sykora L, Kurucova J, et al. Rationale and design of a prospective, randomized, controlled, multicenter study to evaluate the safety and efficacy of transcatheter heart valve replacement in female patients with severe symptomatic aortic stenosis requiring aortic valve intervention (Randomized research in women with Aortic stenosis [RHEIA] trial). *Am Heart J*. 2020;228:27–35. doi: 10.1016/j.ahj.2020.06.016
  44. Van Wiechen MP, Ooms JF, Hokken TW, De Ronde-Tillmans MJ, Goudzwaard JA, Daemen J, De Jaegere PP, Mattace-Raso FU, Van Mieghem NM. Pathways towards lean TAVR. *Structural Heart*. 2020;4:284–287.
  45. Wood DA, Lauck SB, Cairns JA, Humphries KH, Cook R, Welsh R, Leipsic J, Genereux P, Moss R, Jue J, et al. The Vancouver 3M (multidisciplinary, multimodality, but minimalist) clinical pathway facilitates safe next-day discharge home at low-, medium-, and high-volume transfemoral transcatheter aortic valve replacement centers: the 3M TAVR study. *JACC Cardiovasc Interv*. 2019;12:459–469. doi: 10.1016/j.jcin.2018.12.020
  46. Otto CM, Kumbhani DJ, Alexander KP, Calhoun JH, Desai MY, Kaul S, Lee JC, Ruiz CE, Vassileva CM. 2017 ACC expert consensus decision pathway for transcatheter aortic valve replacement in the management of adults with aortic stenosis: a report of the American College of Cardiology task force on clinical expert consensus documents. *J Am Coll Cardiol*. 2017;69:1313–1346. doi: 10.1016/j.jacc.2016.12.006
  47. Francone M, Budde RPJ, Bremerich J, Dacher JN, Loewe C, Wolf F, Natale L, Pontone G, Redheuil A, Vliegenthart R, et al. CT and MR imaging prior to transcatheter aortic valve implantation: standardisation of scanning protocols, measurements and reporting—a consensus document by the European Society of Cardiovascular Radiology (ESCR). *Eur Radiol*. 2020;30:2627–2650. doi: 10.1007/s00330-019-06357-8
  48. Buellesfeld L, Stortecky S, Kalesan B, Gloekler S, Khattab AA, Nietlispach F, Delfino V, Huber C, Eberle B, Meier B, et al. Aortic root dimensions among patients with severe aortic stenosis undergoing transcatheter aortic valve replacement. *JACC Cardiovasc Interv*. 2013;6:72–83. doi: 10.1016/j.jcin.2012.09.007
  49. Dencker D, Taudorf M, Luk NH, Nielsen MB, Kofoed KF, Schroeder TV, Sondergaard L, Lönn L, De Backer O. Frequency and effect of access-related vascular injury and subsequent vascular intervention after transcatheter aortic valve replacement. *Am J Cardiol*. 2016;118:1244–1250. doi: 10.1016/j.amjcard.2016.07.045
  50. Mihos CG, Klassen SL, Yucel E. Sex-specific considerations in women with aortic stenosis and outcomes after transcatheter aortic valve replacement. *Curr Treat Options Cardiovasc Med*. 2018;20:52. doi: 10.1007/s11936-018-0651-x
  51. Herrmann HC, Abdel-Wahab M, Attizzani GF, Batchelor W, Bleiziffer S, Verdoliva S, Chang Y, Gada H, Gillam L, Guerrero M, et al. Rationale and design of the Small Annuli Randomized To Evolve or SAPIEN Trial (SMART Trial). *Am Heart J*. 2022;243:92–102. doi: 10.1016/j.ahj.2021.09.011
  52. Pibarot P, Weissman NJ, Stewart WJ, Hahn RT, Lindman BR, McAndrew T, Kodali SK, Mack MJ, Thourani VH, Miller DC, et al.

- Incidence and sequelae of prosthesis-patient mismatch in transcatheter versus surgical valve replacement in high-risk patients with severe aortic stenosis. *J Am Coll Cardiol*. 2014;64:1323–1334. doi: 10.1016/j.jacc.2014.06.1195
53. Rodés-Cabau J, Pibarot P, Suri RM, Kodali S, Thourani VH, Szeto WY, Svensson LG, Dumont E, Xu K, Hahn RT, et al. Impact of aortic annulus size on valve hemodynamics and clinical outcomes after transcatheter and surgical aortic valve replacement: insights from the PARTNER Trial. *Circ Cardiovasc Interv*. 2014;7:701–711. doi: 10.1161/CIRCINTERVENTIONS.114.001681
  54. Poulin F, Yingchoncharoen T, Wilson WM, Horlick EM, Généreux P, Tuzcu EM, Stewart W, Osten MD, Woo A, Thavendiranathan P. Impact of prosthesis-patient mismatch on left ventricular myocardial mechanics after transcatheter aortic valve replacement. *J Am Heart Assoc*. 2016;5:e002866. doi: 10.1161/JAHA.115.002866
  55. Thyregod HG, Steinbrüchel DA, Ihlemann N, Ngo TA, Nissen H, Kjeldsen BJ, Chang Y, Hansen PB, Olsen PS, Sondergaard L. No clinical effect of prosthesis-patient mismatch after transcatheter versus surgical aortic valve replacement in intermediate- and low-risk patients with severe aortic valve stenosis at mid-term follow-up: an analysis from the NOTION trial. *Eur J Cardiothorac Surg*. 2016;50:721–728. doi: 10.1093/ejcts/ezw095
  56. Akodad M, Sellers S, Gulsin GS, Tzimas G, Landes U, Chatfield AG, Chuang A, Meier D, Leipsic J, Blanke P, et al. Leaflet and neokirt height in transcatheter heart valves: implications for repeat procedures and coronary access. *JACC Cardiovasc Interv*. 2021;14:2298–2300. doi: 10.1016/j.jcin.2021.07.034
  57. Ochiai T, Oakley L, Sekhon N, Komatsu I, Flint N, Kaewkes D, Yoon SH, Raschpichler M, Patel V, Tiwana R, et al. Risk of coronary obstruction due to sinus sequestration in redo transcatheter aortic valve replacement. *JACC Cardiovasc Interv*. 2020;13:2617–2627. doi: 10.1016/j.jcin.2020.09.022
  58. Nai Fovino L, Scotti A, Massussi M, Fabris T, Cardaioli F, Rodinò G, Matsuda Y, Frigo F, Fraccaro C, Tarantini G. Incidence and feasibility of coronary access after transcatheter aortic valve replacement. *Catheter Cardiovasc Interv*. 2020;96:E535–E541. doi: 10.1002/ccd.28720
  59. Kotronias RA, Bray JJH, Rajasundaram S, Vincent F, Delhaye C, Scarsini R, Marin F, Terentes-Printzios D, Halcox J, Mamas MA, et al. Ultrasound- versus fluoroscopy-guided strategy for transfemoral transcatheter aortic valve replacement access: a systematic review and meta-analysis. *Circ Cardiovasc Interv*. 2021;14:e010742. doi: 10.1161/CIRCINTERVENTIONS.121.010742
  60. Di Mario C, Goodwin M, Ristalli F, Ravani M, Meucci F, Stolicova M, Sardella G, Salvi N, Bedogni F, Berti S, et al. A prospective registry of intravascular lithotripsy-enabled vascular access for transfemoral transcatheter aortic valve replacement. *JACC Cardiovasc Interv*. 2019;12:502–504. doi: 10.1016/j.jcin.2019.01.211
  61. Chandrasekhar J, Hibbert B, Ruel M, Lam BK, Labinaz M, Glover C. Transfemoral vs non-transfemoral access for transcatheter aortic valve implantation: a systematic review and meta-analysis. *Can J Cardiol*. 2015;31:1427–1438. doi: 10.1016/j.cjca.2015.04.023
  62. Buchanan GL, Chieffo A, Montorfano M, Maccagni D, Maisano F, Latib A, Covello RD, Grimaldi A, Alfieri O, Colombo A. A "modified crossover technique" for vascular access management in high-risk patients undergoing transfemoral transcatheter aortic valve implantation. *Catheter Cardiovasc Interv*. 2013;81:579–583. doi: 10.1002/ccd.24380
  63. Faurie B, Abdellaoui M, Wautot F, Staat P, Champagnac D, Wintzer-Wehekind J, Vanzetto G, Bertrand B, Monségu J. Rapid pacing using the left ventricular guidewire: reviving an old technique to simplify BAV and TAVI procedures. *Catheter Cardiovasc Interv*. 2016;88:988–993. doi: 10.1002/ccd.26666
  64. Sharp AS, Michev I, Maisano F, Taramasso M, Godino C, Latib A, Denti P, Dorigo E, Giacomini A, Iacì G, et al. A new technique for vascular access management in transcatheter aortic valve implantation. *Catheter Cardiovasc Interv*. 2010;75:784–793. doi: 10.1002/ccd.22238
  65. Power D, Schäfer U, Guedeney P, Claessen BE, Sartori S, Sorrentino S, Lefèvre T, Kupatt K, Tchetché D, Dumonteil N, et al. Impact of percutaneous closure device type on vascular and bleeding complications after TAVR: a post hoc analysis from the BRAVO-3 randomized trial. *Catheter Cardiovasc Interv*. 2019;93:1374–1381. doi: 10.1002/ccd.28295
  66. Masiero G, D'Angelo L, Fovino LN, Fabris T, Cardaioli F, Rodinò G, Benedetti A, Boiago M, Continisio S, Montonati C, et al. Real-world experience with a large bore vascular closure device during TAVI procedure: features and predictors of access-site vascular complications. *Front Cardiovasc Med*. 2022;9:832242. doi: 10.3389/fcvm.2022.832242
  67. Abdel-Wahab M, Hartung P, Dumpies O, Obradovic D, Wilde J, Majunke N, Boekstegers P, Müller R, Seyfarth M, Vorpahl M, et al. Comparison of a pure plug-based versus a primary suture-based vascular closure device strategy for transfemoral transcatheter aortic valve replacement: the CHOICE-CLOSURE randomized clinical trial. *Circulation*. 2021;145:170–183. doi: 10.1161/CIRCULATIONAHA.121.057856
  68. Ten Berg J, Sibbing D, Rocca B, Van Belle E, Chevalier B, Collet JP, Dudek D, Gilard M, Gorog DA, Grapsa J, et al. Management of anti-thrombotic therapy in patients undergoing transcatheter aortic valve implantation: a consensus document of the ESC Working Group on Thrombosis and the European Association of Percutaneous Cardiovascular Interventions (EAPCI), in collaboration with the ESC Council on Valvular Heart Disease. *Eur Heart J*. 2021;42:2265–2269. doi: 10.1093/eurheartj/ehab196
  69. Owais T, El Garhy M, Fuchs J, Disha K, Elkaffas S, Breuer M, Lauer B, Kuntze T. Pathophysiological factors associated with left ventricular perforation in transcatheter aortic valve implantation by transfemoral approach. *J Heart Valve Dis*. 2017;26:430–436.
  70. Rezaq A, Basavarajiah S, Latib A, Takagi K, Hasegawa T, Figini F, Cioni M, Franco A, Montorfano M, Chieffo A, et al. Incidence, management, and outcomes of cardiac tamponade during transcatheter aortic valve implantation: a single-center study. *JACC Cardiovasc Interv*. 2012;5:1264–1272. doi: 10.1016/j.jcin.2012.08.012
  71. Ribeiro HB, Webb JG, Makkar RR, Cohen MG, Kapadia SR, Kodali S, Tamburino C, Barbanti M, Chakravarty T, Jilaihawi H, et al. Predictive factors, management, and clinical outcomes of coronary obstruction following transcatheter aortic valve implantation: insights from a large multicenter registry. *J Am Coll Cardiol*. 2013;62:1552–1562. doi: 10.1016/j.jacc.2013.07.040
  72. Westermann D, Ludwig S, Kalbacher D, Spink C, Linder M, Bhadra OD, Nikorowitsch J, Waldschmidt L, Demal T, Voigtländer L, et al. Prevention of coronary obstruction in patients at risk undergoing transcatheter aortic valve implantation: the Hamburg BASILICA experience. *Clin Res Cardiol*. 2021;110:1900–1911. doi: 10.1007/s00392-021-01881-4
  73. Khan JM, Greenbaum AB, Babaliaros VC, Dvir D, Reisman M, McCabe JM, Satler L, Waksman R, Eng MH, Paone G, et al. BASILICA trial: one-year outcomes of transcatheter electrosurgical leaflet laceration to prevent TAVR coronary obstruction. *Circ Cardiovasc Interv*. 2021;14:e010238. doi: 10.1161/CIRCINTERVENTIONS.120.010238
  74. Poschner T, Werner P, Kocher A, Laufer G, Musumeci F, Andreas M, Russo M. The JenaValve pericardial transcatheter aortic valve replacement system to treat aortic valve disease. *Future Cardiol*. 2021;18:101–113. doi: 10.2217/fca-2021-0065
  75. Abdel-Wahab M, Zahn R, Horack R, Gerckens U, Schuler G, Sievert H, Eggebrecht H, Senges J, Richardt G, German transcatheter aortic valve interventions registry investigators. Aortic regurgitation after transcatheter aortic valve implantation: incidence and early outcome. Results from the German transcatheter aortic valve interventions registry. *Heart*. 2011;97:899–906. doi: 10.1136/hrt.2010.217158
  76. Barbanti M, Yang TH, Rodés Cabau J, Tamburino C, Wood DA, Jilaihawi H, Blanke P, Makkar RR, Latib A, Colombo A, et al. Anatomical and procedural features associated with aortic root rupture during balloon-expandable transcatheter aortic valve replacement. *Circulation*. 2013;128:244–253. doi: 10.1161/CIRCULATIONAHA.113.002947
  77. Mas-Peiro S, Weiler H, Papadopoulos N, Zeiher AM, Fichtlscherer S, Vasa-Nicotera M. Post/preprocedural ratio of hemodynamically assessed aortic regurgitation index as a marker for the need for corrective measures during transcatheter valve replacement: a first confirmatory study in patients receiving a new generation transcatheter self-expandable prosthesis. *Catheter Cardiovasc Interv*. 2019;93:532–537. doi: 10.1002/ccd.27847
  78. Tarantini G, Mojoli M, Purita P, Napodano M, D'Onofrio A, Frigo A, Covolo E, Facchin M, Isabella G, Gerosa G, et al. Unravelling the (arte) fact of increased pacemaker rate with the Edwards SAPIEN 3 valve. *EuroIntervention*. 2015;11:343–350. doi: 10.4244/EJY14M11\_06
  79. Jilaihawi H, Zhao Z, Du R, Staniloae C, Saric M, Neuburger PJ, Querijero M, Vainrib A, Hisamoto K, Ibrahim H, et al. Minimizing permanent pacemaker following repositionable self-expanding transcatheter aortic valve replacement. *JACC Cardiovasc Interv*. 2019;12:1796–1807. doi: 10.1016/j.jcin.2019.05.056

80. Tarantini G, Fabris T, Nai FL. TAVR-in-TAVR and coronary access: importance of preprocedural planning. *EuroIntervention*. 2020;16:e129–e132. doi: [10.4244/EIJ-D-19-01094](https://doi.org/10.4244/EIJ-D-19-01094)
81. Tarantini G, Nai Fovino L, Le Prince P, Darremont O, Urena M, Bartorelli AL, Vincent F, Hovorka T, Alcalá Navarro Y, Dumonteil N, et al. Coronary access and percutaneous coronary intervention up to 3 years after transcatheter aortic valve implantation with a balloon-expandable valve. *Circ Cardiovasc Interv*. 2020;13:e008972. doi: [10.1161/CIRCINTERVENTIONS.120.008972](https://doi.org/10.1161/CIRCINTERVENTIONS.120.008972)
82. Nai Fovino L, Scotti A, Massussi M, Cardaioli F, Rodinò G, Matsuda Y, Pavei A, Masiero G, Napodano M, Fracarro C, et al. Coronary angiography after transcatheter aortic valve replacement (TAVR) to evaluate the risk of coronary access impairment after TAVR-in-TAVR. *J Am Heart Assoc*. 2020;9:e016446. doi: [10.1161/JAHA.120.016446](https://doi.org/10.1161/JAHA.120.016446)
83. Tarantini G, Nai Fovino L. Lifetime strategy of patients with aortic stenosis: the first cut is the deepest. *JACC Cardiovasc Interv*. 2021;14:1727–1730. doi: [10.1016/j.jcin.2021.06.029](https://doi.org/10.1016/j.jcin.2021.06.029)
84. Tarantini G, Nai FL. Coronary access and TAVR-in-TAVR: don't put off until tomorrow what you can do today. *JACC Cardiovasc Interv*. 2020;13:2539–2541. doi: [10.1016/j.jcin.2020.06.065](https://doi.org/10.1016/j.jcin.2020.06.065)
85. Attizzani GF, Alkhalil A, Padaliya B, Tam CC, Lopes JP, Fares A, Bezerra HG, Medallion B, Park S, Deo S, et al. Comparison of Outcomes of transfemoral transcatheter aortic valve implantation using a minimally invasive versus conventional strategy. *Am J Cardiol*. 2015;116:1731–1736. doi: [10.1016/j.amjcard.2015.08.044](https://doi.org/10.1016/j.amjcard.2015.08.044)
86. Babaliaros V, Devireddy C, Lerakis S, Leonardi R, Iturra SA, Mavromatis K, Leshnower BG, Guyton RA, Kanitkar M, Keegan P, et al. Comparison of transfemoral transcatheter aortic valve replacement performed in the catheterization laboratory (minimalist approach) versus hybrid operating room (standard approach): outcomes and cost analysis. *JACC Cardiovasc Interv*. 2014;7:898–904. doi: [10.1016/j.jcin.2014.04.005](https://doi.org/10.1016/j.jcin.2014.04.005)
87. Alkhalil A, Lamba H, Deo S, Bezerra HG, Patel SM, Markowitz A, Simon DI, Costa MA, Davis AC, Attizzani GF. Safety of shorter length of hospital stay for patients undergoing minimalist transcatheter aortic valve replacement. *Catheter Cardiovasc Interv*. 2018;91:345–353. doi: [10.1002/ccd.27230](https://doi.org/10.1002/ccd.27230)
88. Spence MS, Baan J, Iacovelli F, Martinelli GL, Muir DF, Saia F, Bortone AS, Densem CG, Owens CG, van der Kley F, et al. Prespecified risk criteria facilitate adequate discharge and long-term outcomes after transfemoral transcatheter aortic valve implantation. *J Am Heart Assoc*. 2020;9:e016990. doi: [10.1161/JAHA.120.016990](https://doi.org/10.1161/JAHA.120.016990)
89. Nicolas J, Claessen BE, Cao D, Sartori S, Baber U, Power D, Chiarito M, Goel R, Roumeliotis A, Chandiramani R, et al. Preprocedural anemia in females undergoing transcatheter aortic valve implantation: insights from the WIN-TAVI registry. *Catheter Cardiovasc Interv*. 2021;97:E704–E715. doi: [10.1002/ccd.29276](https://doi.org/10.1002/ccd.29276)
90. Leon MB, Smith CR, Mack MJ, Makkar RR, Svensson LG, Kodali SK, Thourani VH, Tuzcu EM, Miller DC, Herrmann HC, et al. Transcatheter or surgical aortic-valve replacement in intermediate-risk patients. *N Engl J Med*. 2016;374:1609–1620. doi: [10.1056/NEJMoa1514616](https://doi.org/10.1056/NEJMoa1514616)
91. Reardon MJ, Van Mieghem NM, Popma JJ, Kleiman NS, Søndergaard L, Mumtaz M, Adams DH, Deeb GM, Maini B, Gada H, et al. Surgical or transcatheter aortic-valve replacement in intermediate-risk patients. *N Engl J Med*. 2017;376:1321–1331. doi: [10.1056/NEJMoa1700456](https://doi.org/10.1056/NEJMoa1700456)
92. Mack MJ, Leon MB, Thourani VH, Makkar R, Kodali SK, Russo M, Kapadia SR, Malaisrie SC, Cohen DJ, Pibarot P, et al. Transcatheter aortic-valve replacement with a balloon-expandable valve in low-risk patients. *N Engl J Med*. 2019;380:1695–1705.
93. Calabrò P, Gragnano F, Niccoli G, Marcucci R, Zimarino M, Spaccarotella C, Renda G, Patti G, Andò G, Moscarella E, et al. Antithrombotic therapy in patients undergoing transcatheter interventions for structural heart disease. *Circulation*. 2021;144:1323–1343. doi: [10.1161/CIRCULATIONAHA.121.054305](https://doi.org/10.1161/CIRCULATIONAHA.121.054305)
94. Capodanno D, Collet JP, Dangas G, Montalescot G, Ten Berg JM, Windecker S, Angiolillo DJ. Antithrombotic therapy after transcatheter aortic valve replacement. *JACC Cardiovasc Interv*. 2021;14:1688–1703. doi: [10.1016/j.jcin.2021.06.020](https://doi.org/10.1016/j.jcin.2021.06.020)
95. Ussia GP, Scarabelli M, Mulè M, Barbanti M, Sarkar K, Cammalleri V, Immè S, Aruta P, Pistrutto AM, Gulino S, et al. Dual antiplatelet therapy versus aspirin alone in patients undergoing transcatheter aortic valve implantation. *Am J Cardiol*. 2011;108:1772–1776. doi: [10.1016/j.amjcard.2011.07.049](https://doi.org/10.1016/j.amjcard.2011.07.049)
96. Rodés-Cabau J, Masson JB, Welsh RC, Garcia Del Blanco B, Pelletier M, Webb JG, Al-Qoofi F, Génèreux P, Maluenda G, Thoenes M, et al. Aspirin versus aspirin plus clopidogrel as antithrombotic treatment following transcatheter aortic valve replacement with a balloon-expandable valve: the ARTE (Aspirin Versus Aspirin+Clopidogrel Following Transcatheter Aortic Valve Implantation) randomized clinical trial. *JACC Cardiovasc Interv*. 2017;10:1357–1365. doi: [10.1016/j.jcin.2017.04.014](https://doi.org/10.1016/j.jcin.2017.04.014)
97. Brouwer J, Nijenhuis VJ, Delewi R, Hermanides RS, Holvoet W, Dubois CLF, Frambach P, De Bruyne B, van Houwelingen GK, Van Der Heyden JAS, et al. Aspirin with or without clopidogrel after transcatheter aortic-valve implantation. *N Engl J Med*. 2020;383:1447–1457. doi: [10.1056/NEJMoa2017815](https://doi.org/10.1056/NEJMoa2017815)
98. Nijenhuis VJ, Brouwer J, Delewi R, Hermanides RS, Holvoet W, Dubois CLF, Frambach P, De Bruyne B, van Houwelingen GK, Van Der Heyden JAS, et al. Anticoagulation with or without clopidogrel after transcatheter aortic-valve implantation. *N Engl J Med*. 2020;382:1696–1707. doi: [10.1056/NEJMoa1915152](https://doi.org/10.1056/NEJMoa1915152)
99. Dangas GD, Tijssen JGP, Wöhrle J, Søndergaard L, Gilard M, Möllmann H, Makkar RR, Herrmann HC, Giustino G, Baldus S, et al. A controlled trial of rivaroxaban after transcatheter aortic-valve replacement. *N Engl J Med*. 2020;382:120–129. doi: [10.1056/NEJMoa1911425](https://doi.org/10.1056/NEJMoa1911425)
100. Van Mieghem NM, Unverdorben M, Hengstenberg C, Möllmann H, Mehran R, López-Otero D, Nombela-Franco L, Moreno R, Nordbeck P, Thiele H, et al. Edoxaban versus vitamin K antagonist for atrial fibrillation after TAVR. *N Engl J Med*. 2021;385:2150–2160. doi: [10.1056/NEJMoa2111016](https://doi.org/10.1056/NEJMoa2111016)
101. Collet J-P. *Oral anti-Xa anticoagulation after trans-aortic valve implantation for aortic stenosis: the randomized ATLANTIS trial* Presented at: ACC 2021. May 15, 2021.