



# Super Responder Profile Under Bimekizumab Treatment in Moderate-to-Severe Psoriasis: A Short Term Real-Life Observation—IL PSO (Italian Landscape Psoriasis)

Maria Esposito<sup>1</sup> · Paolo Gisondi<sup>2</sup> · Chiara Assorgi<sup>3,4</sup> · Francesco Bellinato<sup>2</sup> · Pina Brianti<sup>5</sup> · Martina Burlando<sup>6</sup> · Giovanna Brunasso<sup>7</sup> · Stefano Caccavale<sup>8</sup> · Giacomo Caldarola<sup>9</sup> · Elena Campione<sup>10</sup> · Piergiacomo Calzavara Pinton<sup>11</sup> · Anna Campanati<sup>12</sup> · Carlo Giovanni Carrera<sup>13</sup> · Andrea Carugno<sup>14,15</sup> · Emanuele Cozzani<sup>6</sup> · Antonio Costanzo<sup>16</sup> · Francesco Cusano<sup>17</sup> · Paolo Dapavo<sup>18</sup> · Annunziata Dattola<sup>19</sup> · Clara De Simone<sup>9</sup> · Roberta Di Caprio<sup>8</sup> · Federico Diotallevi<sup>12</sup> · Maria Concetta Fagnoli<sup>1</sup> · Francesca Gaiani<sup>20</sup> · Alessandro Giunta<sup>10</sup> · Piergiorgio Malagoli<sup>20</sup> · Angelo Valerio Marzano<sup>13,21</sup> · Matteo Megna<sup>4</sup> · Santo Raffaele Mercuri<sup>5</sup> · Edoardo Mortato<sup>22</sup> · Alessandra Narcisi<sup>16</sup> · Diego Orsini<sup>3</sup> · Luca Potestio<sup>4</sup> · Pietro Quaglino<sup>18</sup> · Antonio Giovanni Richetta<sup>19</sup> · Francesca Romano<sup>23</sup> · Paolo Sena<sup>14</sup> · Emanuele Vagnozzi<sup>1</sup> · Marina Venturini<sup>11</sup> · Francesco Loconsole<sup>22</sup> · Anna Balato<sup>8</sup>

Accepted: 8 April 2025 / Published online: 15 May 2025

© The Author(s), under exclusive licence to Springer Nature Switzerland AG 2025

## Key Points

There is still a lack of knowledge on the characteristics of super responders (SRs) during treatment with bimekizumab, since genetic, clinical, and biomarkers investigations are required.

During bimekizumab treatment, a high rate of SRs was demonstrated at early clinical timepoints, with some interesting differences between SRs and non-super-responders.

## 1 Introduction

Bimekizumab is a humanized monoclonal antibody, targeting both IL-17A and IL-17F, that has been recently approved for the treatment of moderate-to-severe plaque psoriasis following encouraging results from four phase-III clinical trials [1–4]. Real-world experiences on bimekizumab are

limited and there is still limited knowledge about the super responder (SR) patient profile. There is no agreement on the definition of SR with respect to psoriasis even though this term has been used in different clinical studies [5–7].

We conducted a 16-week multicenter, retrospective, observational, real-world study aimed at assessing the SR profile under bimekizumab treatment in patients with moderate-to-severe psoriasis.

## 2 Materials and Methods

We enrolled patients from 21 Italian dermatologic Departments. Consecutive adult patients with moderate-to-severe plaque psoriasis if they had completed at least 16 weeks of bimekizumab treatment were considered eligible for inclusion in the study. Patient eligibility was assessed according to the rules of the Italian Medicine Agency (AIFA). Bimekizumab was administered at a dosage of 320 mg at weeks 0, 4, 8, 12, and 16 followed by an administration every 8 weeks, according to the summary of product characteristics. Patients whose clinical reports were not available and those who had not reached 16 weeks of follow-up were excluded. Patients signed an informed consent to allow collection and utilization of clinical data for scientific purposes. This study adhered to established standards of good clinical practice. Ethical approval number 0006349 was obtained from the Comitato Etico Territoriale (CET) Interaziendale AOU Città della Salute e della Scienza di Torino. Effectiveness

Maria Esposito and Paolo Gisondi have contributed equally to this work.

Francesco Loconsole and Anna Balato have contributed equally to this work.

Extended author information available on the last page of the article

endpoints were the improvement of 75% (PASI 75), 90% (PASI 90), and 100% (PASI 100) in Psoriasis Area and Severity Index (PASI) at different timepoints (week 4 and week 16) as compared to baseline. In this retrospective evaluation we defined as SR a patient achieving PASI 100 at each timepoint, while patients who did not achieve PASI 100 were defined as non-super-responders (NSRs).

## 2.1 Statistical Analysis

The percentage of patients who achieved PASI 100 at each timepoint (week 4 and week 16) was determined, and the percentage of these individuals with each of the following characteristics was reported: sex, age, body mass index (BMI), comorbidities, arthritis, PASI, special localization, previous systemic therapies, and previous biologic exposure. We conducted univariate analysis on each of these characteristics taking into account the achievement of PASI 100 as outcome variable. Moreover, to evaluate the possible differences in the demographic and clinical variables between the two groups of patients (SR vs. NSR), univariate analysis has been estimated. Categorical values were described by absolute and relative frequencies. Two-sided *p*-values below 0.05 were considered significant.

To evaluate the determinants of SR status achievement at week 16 we constructed a multivariable logistic regression model. The multicollinearity assumption was tested using a variance inflation factor, eliminating variables with high correlation (value > 0.9).

We included all the baseline characteristics in the multivariate regression model, odds ratios (ORs) and 95% confidence intervals (CIs) are provided. Statistical analyses were performed using SAS software v 9.3 (SAS Institute Inc., Cary, NC, USA).

## 3 Results

We enrolled 137 patients with a mean age of  $52.47 \pm 15.56$  years, mean BMI of  $27.43 \pm 5.91$  and a mean PASI at baseline of  $16.00 \pm 9.29$ . Additional demographic and clinical characteristics of our cohort at baseline are shown in Table 1.

At week 4, 72% of patients achieved PASI 75, 50% PASI 90, and 43% of patients achieved complete clearance (PASI 100), defining the SR population at week 4 (Fig. 1A).

At week 16, 93% of patients obtained PASI 75, 77% PASI 90, and 70% of patients experienced complete disease resolution (PASI 100) while only 7% of patients did not achieve PASI 75 (Fig. 1B).

**Table 1** Demographic and clinical characteristics of the study sample


	137
	Mean $\pm$ SD
<b>Number of total patients</b>	
<b>Age (years)</b>	52.47 $\pm$ 15.56
<b>BMI</b>	27.43 $\pm$ 5.91
<b>mPASI at baseline</b>	16.00 $\pm$ 9.29
<b>Males</b>	97 (70.80)
<b>Comorbidities</b>	
None	71 (65.23)
1	39 (28.47)
$\geq 2$	27 (19.71)
<b>Arthritis</b>	
Yes	17 (12.40)
No	120 (87.60)
<b>Special localization</b>	
Nails	24 (17.52)
Scalp	48 (35.04)
Genital	20 (14.60)
Palmo-plantar	7 (5.11)
<b>Previous systemic therapies</b>	
Mtx	34 (24.82)
Cys	52 (37.96)
Aci	12 (8.76)
<b>Previous biologic exposure</b>	
Naïve	62 (45.26)
< 2	42 (30.65)
$\geq 2$	33 (24.09)

*BMI* body mass index, *PASI* Psoriasis Area Severity Index, *MTX* methotrexate, *Cys* cyclosporine, *Aci* acitretin

A univariate analysis of the differences between the SR status at week 4 and week 16 was performed. All the following variables (sex, age, BMI, comorbidities, presence of arthritis, PASI, special localizations, previous systemic therapies, and previous exposure to biological drugs) were analyzed and no significant differences were encountered (data not shown).

A further univariate analysis was conducted to compare patients who achieved the SR status and patients who did not achieved the SR status (NSRs) at week 16 (Table 2). More characteristics differentiated SR from NSR. In detail, male sex ( $p = 0.0002$ ), age  $\leq 45$  years and age between 46 and 65 years ( $p = 0.0004$  and  $p = 0.003$ , respectively), BMI < 25 ( $p < 0.0001$ ), absence of comorbidities ( $p = 0.001$ ), absence of arthritis ( $p < 0.0001$ ), baseline PASI between 10 and 20 ( $p = 0.004$ ), scalp and genital localization ( $p = 0.002$  both), naïve status and prior exposure to less than two biological treatments ( $p = 0.0006$  and  $p < 0.0001$ , respectively) were more represented in SR subjects (Table 2).

## PASI response rates



**Fig. 1** Bimekizumab clinical effectiveness as assessed by Psoriasis Area Severity Index (PASI) 75, PASI 90, and PASI 100 at week 4 (A) and week 16 (B)

A multivariable logistic regression model analysis having as outcome PASI 100 at W16 was performed showing that males were more likely to achieve PASI 100 than female patients (OR: 2.692; 95% CI: 1.594–4.482;  $p < 0.001$ ). Patients aged  $> 65$  years were less likely to achieve PASI 100 response as compared to patients aged  $< 45$  years (OR: 0.2778; 95% CI: 0.1371–0.5700;  $p < 0.001$ ). No statistical differences were encountered regarding BMI. The positive predictor of PASI 100 at week 16 was the absence of comorbidities having as reference  $\geq$  two comorbidities (OR: 3; 95% CI: 1.653–5.518;  $p < 0.001$ ). Not having a concurrent arthritis also showed greater odds to achieve PASI 100 at week 16 than patients with PsA (OR: 3.5; 95% CI: 2.212–6.100;  $p < 0.001$ ). A baseline PASI score between 10 and 20 was associated with significantly higher odds compared to a baseline PASI score  $< 10$  (OR: 3.100; 95% CI: 1.900–5.920;  $p < 0.001$ ). No statistical differences were encountered regarding special localizations or previous non-biologic treatments. Conversely, previous exposure to at least two biologics was associated with significantly lower odds to achieve PASI 100 at week 16 compared with bio-naïve patients (OR: 0.2391; 95% CI: 0.1158–0.4815;  $p < 0.001$ ). Results of the multivariable logistic regression model at week 16 are shown in Fig. 2.

## 4 Discussion

Although there is no consensus on the use of the SR concept in the field of psoriasis, the raised bar of effectiveness available during bimekizumab treatment allowed us to define SRs as those reaching PASI 100 at week 4 and at week 16.

Bimekizumab demonstrated a rapid and consistent efficacy associated with a good safety profile in randomized clinical trials (RCTs) [1–4], while, to date, real-life data are limited [8–10]. As a consequence, there is still a lack of knowledge on the characteristics of SRs during treatment with bimekizumab; clinical and genetic factors might be of importance in defining this sub-group.

Some data on the SR status may be deduced from RCTs; in particular, Warren et al. [1] reported PASI 100 response in 15.4% of patients at week 4 and in 60.8% of patients at week 16, while Gordon et al. [2] described complete clearance in 18.9% and 68.2% of patients at weeks 4 and 16, respectively. Reich et al. showed PASI 100 response in 15% of patients at week 4 and in 59% of patients at week 16 in the “BE VIVID” clinical trial [3], while in the “BE RADIANT” clinical trial [4] 13.9% and 61.7% of patients obtained PASI 100 at weeks 4 and 16, respectively. Findings from our real-life experience clearly showed better results compared to RCTs, as PASI 100 was achieved by 40% of patients at week 4 and by 70% of patients at week 16, demonstrating a consistent SR population under bimekizumab treatment.

In order to investigate the dynamic process underlying achievement of the SR status, we analyzed the differences between patients who reached complete clearance at week 4 and week 16, but no significant differences were identified through the analyzed variables.

Our observation delineated a bimekizumab SR profile. The following baseline characteristics were associated with PASI 100 at week 16: male sex, age  $< 65$  years, absence of comorbidities and arthritis, baseline PASI between 10 and 20, naïve status or prior exposure to less than two biological treatments.

**Table 2** Description of the clinical and demographic variables according to the patient response profile at week 16. Univariate analysis

Population ( <i>n</i> = 137)	Week 16		<i>p</i> -value
	NSR <i>n</i> (%)	SR <i>n</i> (%)	
	41 (30)	96 (70)	
<b>Gender</b>			
Male	27 (20)	70 (51)	0.0002**
Female	14 (10)	26 (19)	0.091
<b>Age, years</b>			
≤ 45	10 (7)	36 (26)	0.0004**
≥ 46 to < 65	22 (16)	50 (37)	0.003*
≥ 65	9 (7)	10 (7)	1.000
<b>BMI</b>			
< 25	7 (5)	36 (25)	< 0.0001**
≥ 25 to < 30	25 (18)	32 (24)	0.467
≥ 30	9 (7)	28 (21)	0.005
<b>Comorbidities</b>			
None	20 (14)	51 (38)	0.001*
1	11 (8)	28 (21)	0.017
≥ 2	10 (7)	17 (12)	0.229
<b>Arthritis</b>			
Yes	7 (5)	10 (7)	0.619
No	34 (25)	86 (63)	< 0.0001**
<b>PASI</b>			
< 10	4 (3)	14 (10)	0.028
≥ 10 to < 20	26 (19)	57 (42)	0.004*
≥ 20	11 (8)	25 (18)	0.033
<b>Special localization</b>			
Nails	9 (7)	15 (11)	0.290
Scalp	12 (9)	36 (26)	0.002*
Genital	3 (2)	17 (12)	0.002*
Palmo-plantar	3 (2)	4 (3)	1.000
<b>Previous systemic therapies</b>			
Mtx	10 (7)	24 (18)	0.029
Cys	15 (11)	37 (27)	0.006
Aci	5 (4)	7 (5)	0.770
<b>Previous biologic exposure</b>			
Naïve	16 (12)	46 (34)	0.0006**
< 2	3 (2)	39 (28)	< 0.0001**
≥ 2	22 (16)	11 (8)	0.096

NSR non-super-responder, SR super responder, BMI body mass index, PASI Psoriasis Area Severity Index, MTX methotrexate, Cys cyclosporine, Aci acitretin

Most of these data have implications from a clinical point of view. Our study delineates a better response profile in male and younger patients, as recently

demonstrated by Hagino et al. [12], in the absence of comorbidities and arthritis and in patients with a PASI between 10 and 20. A possible explanation for why patients with PASI <10 do not reach statistical significance may arise from the consideration that a large proportion of patients (55%) in our sample were failure/multifailure patients, possibly representing patients with residual resistant psoriasis localizations and difficult-to-treat areas, despite a low PASI score. However, the small sample size does not allow us to confirm this hypothesis and provide statistical insights. Concerning other clinical disease characteristics, the presence of scalp or genital localization delineated a favorable clinical variant for a complete response to bimekizumab, while nail and palmoplantar psoriasis were demonstrated as more difficult-to-treat locations. A possible explanation of this latter result in an optimally responding population may be that a short-term 16-week observation period may not be sufficient to produce significant improvement on nail disease and palmoplantar localizations. A further long-term update of this analysis will be needed to confirm this clinical hypothesis. Instead, the observation of optimal results on genital psoriasis and on scalp psoriasis response patterns confirms previously reported positive bimekizumab results, also in the short term [13, 14]. Naïve status and prior exposure to less than two biologics seem to favor a positive response to bimekizumab. This observation is partially in line with a previous real-life report describing 56 patients undergoing bimekizumab treatment and demonstrating lower response in bio-experienced patients at week 4, although the observation was not confirmed at week 16, suggesting that a previous failure of biologics does not seem to affect its therapeutic effectiveness [15]. Notably, all the results were confirmed by the regression analysis.

#### 4.1 Limitations

Limitations of this study were its retrospective nature, the limited sample size, the short follow-up observation period, and the lack of uniformity of clinical assessment with a high number of involved clinicians. The most important findings of the study were the high rate of SR achieved at early clinical timepoints during bimekizumab treatment, with some interesting differences between SRs and NSRs.

In conclusion, our results suggest the optimal bimekizumab patient profile and allow us to identify the place in therapy of this new therapeutical option, although further and larger studies are needed.

# W16




Fig. 2 Results of the multivariable logistic regression model at week 16

## Declarations

Funding None.

**Conflicts of Interest** M. Esposito has served as a speaker/board member for Abbvie, Almirall, Eli Lilly, Janssen, Leo Pharma, Novartis, Sanofi, UCB. A. Balato has received honoraria for participation in advisory boards, meetings, or as a speaker for AbbVie, Celgene, Janssen-Cilag, Eli Lilly, Novartis Pharma, Pfizer, Sanofi-Genzyme, and UCB Pharma. P. Malagoli has been a speaker for AbbVie, Lilly, Novartis, Janssen-Cilag, Celgene, Leopharma, and Almirall. G. Calderola reports consulting fees or honorarium and payment for lectures from Lilly and Novartis. P. Gisondi has been a consultant and/or speaker for Abbvie, Almirall, Amgen, Janssen, Leo-Pharma, Eli-Lilly, Novartis, Pierre Fabre, Sandoz, Sanofi, and UCB. C. Carrera has served as a board participant or speaker for Abbvie, Lilly, Janssen, Novartis, Celgene, Almirall, and Leopharma. P. Dapavo has been a speaker for Novartis, Abbvie, Sanofi, UCB, Janssen, Lilly, and LeoPharma. F. Gaiani acted as a speaker or consultant for Novartis, Abbvie, Eli Lilly, Celgene, LeoPharma, and Almirall. A.Giunta received grants or is an investigator for Biogen and Lilly; and is a consultant/advisory

board/speaker for AbbVie, Almirall, Celgene, Janssen, Leo Pharma, Eli Lilly, Merck Sharpe Dohme, Novartis, Pfizer, Sandoz, and UCB. F.Loconsole served on advisory boards and/ or received honoraria for lectures from Abbvie, Janssen-Cilag, Novartis, Lilly, Sanofi. A.V. Marzano reports consultancy/advisory board disease-relevant honoraria from AbbVie, Boehringer Ingelheim, Novartis, Pfizer, Sanofi and UCB. M. Megna has served as a speaker/board member for Abbvie, Almirall, Amgen, Eli Lilly, Janssen, Leo Pharma, Novartis, UCB. M. Venturini served as a speaker or advisory board member for Abbvie, Almirall, Amgen, Eli-Lilly, Galderma, Leo Pharma, Novartis, Pierre Fabre, and UCB Pharma. A. Costanzo has served as an advisory board member, consultant and has received fees and speaker's honoraria or has participated in clinical trials for Abbvie, Almirall, Biogen, LEO Pharma, Lilly, Janssen, Novartis, Pfizer, Sanofi Genzyme, and UCB-Pharma. A. Narcisi has served on advisory boards, received honoraria for lectures and research grants from Almirall, Abbvie, Leo Pharma, Celgene, Eli Lilly, Janssen, Novartis, Sanofi-Genzyme, Amgen, and Boehringer Ingelheim. A. Carugno has served as a speaker/board member for Abbvie, Almirall, Boehringer Ingelheim, Eli Lilly, Janssen, Leo Pharma, Novartis, UCB Pharma. C. De Simone served as a speaker or advisory board member for Almirall, AbbVie, Amgen, Eli Lilly, Janssen, Leopharma, Novartis, Sanofi and UCB Pharma. M.C.

Fargnoli has served on advisory boards, received honoraria for lectures and/or research grants from AMGEN, Almirall, Abbvie, Boehringer-Ingelheim, BMS, Galderma, Kyowa Kyirin, Leo Pharma, Pierre Fabre, UCB, Lilly, Pfizer, Janssen, MSD, Novartis, Sanofi-Regeneron, Sunpharma. M. Burlando served as a speaker or advisory board member for Almirall, AbbVie, Amgen, Eli Lilly, Janssen, LeoPharma, Novartis, and UCB Pharma. D. Orsini has been a speaker and/or consultant for Abbvie, LeoPharma, UCB, Bristol-Meyer Squibb and Boehringer-Ingelheim. The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

**Ethics Approval** Patients signed an informed consent to allow collection and utilization of clinical data for scientific purposes. This study adhered to established standards of good clinical practice. Ethical approval number 0006349 was obtained from the Comitato Etico Territoriale (CET) Interaziendale AOU Città della Salute e della Scienza di Torino.

**Consent to Participate** All patients gave written informed consent for the retrospective retrieval of anonymized data. All patients gave written informed consent for the publication of anonymized data.

**Consent for Publication** Not applicable.

**Availability of Data and Material** Additional data supporting the findings of this study are available from the corresponding author on reasonable request.

**Code Availability** Not applicable.

**Authors' Contributions** Study conception and design: Maria Esposito, Paolo Gisondi, Chiara Assorgi, Francesco Bellinato, Pina Brianti, Martina Burlando, Giovanna Brunasso, Stefano Caccavale, Giacomo Caldarola, Elena Campione, Piergiacomo Calzavara Pinton, Anna Campanati, Carlo Giovanni Carrera, Andrea Carugno, Emanuele Cozzani, Antonio Costanzo, Francesco Cusano, Paolo Dapavo, Annunziata Dattola, Clara De Simone, Roberta Di Caprio, Federico Diotallevi, Maria Concetta Fargnoli, Francesca Gaiani, Alessandro Giunta, Piergiorgio Malagoli, Angelo Valerio Marzano, Matteo Megna, Santo Raffaele Mercuri, Edoardo Mortato, Alessandra Narcisi, Diego Orsini, Luca Potestio, Pietro Quaglino, Antonio Giovanni Richetta, Francesca Romano, Paolo Sena, Emanuele Vagnozzi, Marina Venturini, Francesco Loconsole, Anna Balato; manuscript editing: Maria Esposito, Paolo Gisondi, Francesco Loconsole, Anna Balato; approval to submit: Maria Esposito, Paolo Gisondi, Francesco Loconsole, Anna Balato.


**Open Access** This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article or parts of it. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

## References

- Warren RB, Blauvelt A, Bagel J, Papp KA, Yamauchi P, Armstrong A, et al. Bimekizumab versus Adalimumab in Plaque Psoriasis. *N Engl J Med.* 2021;385(2):130–41. <https://doi.org/10.1056/NEJMoa2102388>. (Epub 2021 Apr 23).
- Gordon KB, Foley P, Krueger JG, Pinter A, Reich K, Vender R, et al. Bimekizumab efficacy and safety in moderate to severe plaque psoriasis (BE READY): a multicentre, double-blind, placebo-controlled, randomised withdrawal phase 3 trial. *Lancet.* 2021;397(10273):475–86. [https://doi.org/10.1016/S0140-6736\(21\)00126-4](https://doi.org/10.1016/S0140-6736(21)00126-4). (Erratum in: *Lancet.* 2021 Mar 27;397(10280):1182. 10.1016/S0140-6736(21)00669-3).
- Reich K, Papp KA, Blauvelt A, Langley RG, Armstrong A, Warren RB, Gordon KB, et al. Bimekizumab versus ustekinumab for the treatment of moderate to severe plaque psoriasis (BE VIVID): efficacy and safety from a 52-week, multicentre, double-blind, active comparator and placebo controlled phase 3 trial. *Lancet.* 2021;397(10273):487–98. [https://doi.org/10.1016/S0140-6736\(21\)00125-2](https://doi.org/10.1016/S0140-6736(21)00125-2). (Erratum in: *Lancet.* 2021 Feb 20;397(10275):670. 10.1016/S0140-6736(21)00387-1).
- Reich K, Warren RB, Lebwohl M, Gooderham M, Strober B, Langley RG, Paul C, et al. Bimekizumab versus Secukinumab in Plaque Psoriasis. *N Engl J Med.* 2021;385(2):142–52. <https://doi.org/10.1056/NEJMoa2102383>. (Epub 2021 Apr 23).
- Talamonti M, D'Adamio S, Galluccio T, Andreani M, Pastorino R, Egan CG, et al. High-resolution HLA typing identifies a new “super responder” subgroup of HLA-C\*06:02-positive psoriatic patients: HLA-C\*06:02/HLA-C\*04, in response to ustekinumab. *J Eur Acad Dermatol Venereol.* 2019;33(10):e364–7. <https://doi.org/10.1111/jdv.15659>. (Epub 2019 May 20).
- Ruiz-Villaverde R, Vasquez-Chinchay F, Rodriguez-Fernandez-Freire L, Armario-Hita JC, Pérez-Gil A, Galán-Gutiérrez M. Super-responders in moderate-severe psoriasis under guselkumab treatment: myths, realities and future perspectives. *Life (Basel).* 2022;12(9):1412. <https://doi.org/10.3390/life12091412>.
- Loft N, Egeberg A, Rasmussen MK, Bryld LE, Nissen CV, Dam TN, et al. Prevalence and characterization of treatment-refractory psoriasis and super-responders to biologic treatment: a nationwide study. *J Eur Acad Dermatol Venereol.* 2022;36(8):1284–91. <https://doi.org/10.1111/jdv.18126>. (Epub 2022 Apr 12).
- Gargiulo L, Narcisi A, Ibba L, Balato A, Bianchi L, Brianti P, et al. Effectiveness and safety of bimekizumab for the treatment of plaque psoriasis: a real-life multicenter study-IL PSO (Italian landscape psoriasis). *Front Med (Lausanne).* 2023;8(10):1243843. <https://doi.org/10.3389/fmed.2023.1243843>.
- Valenti M, Gargiulo L, Ibba L, Pavia G, Narcisi A, Costanzo A. Sub-erythrodermic psoriasis successfully treated with bimekizumab: a case report. *Dermatol Ther.* 2022;35(12):e15952. <https://doi.org/10.1111/dth.15952>.
- Megna M, Battista T, Potestio L, Ruggiero A, Ventura V, Fabbrocini G, et al. A case of erythrodermic psoriasis rapidly and successfully treated with Bimekizumab. *J Cosmet Dermatol.* 2023;22(3):1146–8. <https://doi.org/10.1111/jocd.15543>. (Epub 2022 Nov 29).
- Lynde C, Riedl E, Maul JT, Torres T, Pinter A, Fabbrocini G et al. Comparative Effectiveness of Biologics Across Subgroups of Patients with Moderate-to-Severe Plaque Psoriasis: Results at Week 12 from the PSoHO Study in a Real-World Setting. *Adv Ther.* 2023 Mar;40(3):869–886. <https://doi.org/10.1007/s12325-022-02379-9>. (Epub 2022 Dec 14).
- Hagino T, Saeki H, Fujimoto E, Kanda N. Effectiveness of long-term bimekizumab treatment and predictive factors for responders in moderate-to-severe psoriasis: a 52-week real-world study. *J*

- Dermatol. 2025;52(2):317–28. <https://doi.org/10.1111/1346-8138.17532>. (Epub 2024 Nov 5).
13. Orsini D, Malagoli P, Balato A, Bianchi L, Brianti P, Buononato D, et al. Bimekizumab for the Treatment of Plaque Psoriasis with Involvement of Genitalia: a 16-Week Multicenter Real-World Experience - IL PSO (Italian Landscape Psoriasis). *Dermatol Pract Concept*. 2024;14(2): e2024052. <https://doi.org/10.5826/dpc.1402a52>.
  14. Hagino T, Onda M, Saeki H, Fujimoto E, Kanda N. Effectiveness of bimekizumab for genital, nail, and scalp lesions with psoriasis: a 24-week real-world study. *J Dermatol*. 2024;51(12):1658–64. <https://doi.org/10.1111/1346-8138.17427>. (Epub 2024 Aug 12).
  15. Megna M, Ruggiero A, Torta G, Martora F, Bianco M, Potestio L. Efficacy and safety of bimekizumab in real-world psoriasis management: data from patients who are biologic-naive vs. biologic-experienced. *Clin Exp Dermatol*. 2024;49(10):1186–1189. <https://doi.org/10.1093/ced/llae147>

## Authors and Affiliations

Maria Esposito<sup>1</sup>  · Paolo Gisondi<sup>2</sup> · Chiara Assorgi<sup>3,4</sup> · Francesco Bellinato<sup>2</sup> · Pina Brianti<sup>5</sup> · Martina Burlando<sup>6</sup> · Giovanna Brunasso<sup>7</sup> · Stefano Caccavale<sup>8</sup> · Giacomo Caldarola<sup>9</sup> · Elena Campione<sup>10</sup> · Piergiacomo Calzavara Pinton<sup>11</sup> · Anna Campanati<sup>12</sup> · Carlo Giovanni Carrera<sup>13</sup> · Andrea Carugno<sup>14,15</sup> · Emanuele Cozzani<sup>6</sup> · Antonio Costanzo<sup>16</sup> · Francesco Cusano<sup>17</sup> · Paolo Dapavo<sup>18</sup> · Annunziata Dattola<sup>19</sup> · Clara De Simone<sup>9</sup> · Roberta Di Caprio<sup>8</sup> · Federico Diotallevi<sup>12</sup> · Maria Concetta Fargnoli<sup>1</sup> · Francesca Gaiani<sup>20</sup> · Alessandro Giunta<sup>10</sup> · Piergiorgio Malagoli<sup>20</sup> · Angelo Valerio Marzano<sup>13,21</sup> · Matteo Megna<sup>4</sup> · Santo Raffaele Mercuri<sup>5</sup> · Edoardo Mortato<sup>22</sup> · Alessandra Narcisi<sup>16</sup> · Diego Orsini<sup>3</sup> · Luca Potestio<sup>4</sup> · Pietro Quaglino<sup>18</sup> · Antonio Giovanni Richetta<sup>19</sup> · Francesca Romano<sup>23</sup> · Paolo Sena<sup>14</sup> · Emanuele Vagnozzi<sup>1</sup> · Marina Venturini<sup>11</sup> · Francesco Loconsole<sup>22</sup> · Anna Balato<sup>8</sup>

✉ Maria Esposito  
maria.esposito3@univaq.it

<sup>1</sup> Department of Biotechnological and Applied Clinical Sciences, University of L'Aquila, via Vetoio-Coppito 2, 67100 L'Aquila, Italy

<sup>2</sup> Section of Dermatology and Venereology, Department of Medicine, University of Verona, Verona, Italy

<sup>3</sup> Clinical Dermatology Unit, San Gallicano Dermatological Institute IRCCS, Rome, Italy

<sup>4</sup> Section of Dermatology, Department of Clinical Medicine and Surgery, University of Naples Federico II, Naples, Italy

<sup>5</sup> Unit of Dermatology and Cosmetology, Vita-Salute San Raffaele University and Institute for Research and Care, Milan, Italy

<sup>6</sup> Section of Dermatology, Department of Health Sciences (DISSAL), IRCCS San Martino University Hospital, Genoa, Italy

<sup>7</sup> Department of Dermatology, Galliera Hospital, Genova, Italy

<sup>8</sup> Dermatology Unit, University of Campania L. Vanvitelli, Naples, Italy

<sup>9</sup> Section of Dermatology, Department of Translational Medicine and Surgery, Catholic University of the Sacred Heart, Rome, Italy

<sup>10</sup> Department of Dermatology, University of Rome 'Tor Vergata', Rome, Italy

<sup>11</sup> Department of Dermatology, University of Brescia, ASST Spedali Civili, Brescia, Italy

<sup>12</sup> Dermatological Clinic Department of Clinical and Molecular Sciences, Polytechnic University of the Marche Region, Ancona, Italy

<sup>13</sup> Dermatology Unit, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy

<sup>14</sup> Dermatology Unit, ASST Papa Giovanni XXIII Hospital, Bergamo, Italy

<sup>15</sup> Ph.D. Program in Molecular and Translational Medicine (DIMET), University of Milan-Bicocca, Milan, Italy

<sup>16</sup> Dermatology Unit, IRCCS Humanitas Clinical and Research Center, Rozzano (Milan), Italy

<sup>17</sup> Department of Dermatology, Gaetano Rummo Hospital, Benevento, Italy

<sup>18</sup> Department of Biomedical Science and Human Oncology, Second Dermatologic Clinic, University of Turin, Turin, Italy

<sup>19</sup> Dermatology Clinic, Department of Clinical Internal, Anesthesiological and Cardiovascular Sciences, Sapienza University, Rome, Italy

<sup>20</sup> Department of Dermatology, Dermatology Unit Azienda Ospedaliera San Donato Milanese, Milan, Italy

<sup>21</sup> Department of Pathophysiology and Transplantation, Università degli Studi di Milano, Milan, Italy

<sup>22</sup> Department of Dermatology, University of Bari, Piazza Umberto I, 1, 70121 Bari, Italy

<sup>23</sup> Dermatology Unit, AORN "A. Cardarelli", Naples, Italy