








Defining benchmarks for total and distal gastrectomy: global multicentre analysis

Marcel André Schneider¹ , Jeesun Kim², Felix Berth³, Yutaka Sugita⁴, Peter P. Grimmer³, Takeshi Sano⁴, Riccardo Rosati⁵, Gian Luca Baiocchi⁶ , Maria Bencivenga⁷, Giovanni De Manzoni⁷, Souya Nunobe⁴ , Han-Kwang Yang²  and Christian Alexander Gutschow^{1,*}  for the GastroBenchmark Consortium and the GASTRODATA Consortium

¹Department of Surgery and Transplantation, University Hospital Zurich, Zurich, Switzerland

²Department of Surgery, Seoul National University Cancer Hospital, Seoul, South Korea


³Department of General, Visceral and Transplant Surgery, University Medical Centre Mainz, Mainz, Germany

⁴Department of Gastroenterological Surgery, Cancer Institute Hospital of the Japanese Foundation for Cancer Research, Tokyo, Japan

⁵Department of Surgery, San Raffaele Hospital, Milan, Italy

⁶Department of Surgery, University Hospital of Brescia, Brescia, Italy

⁷Department of Surgery, University Hospital of Verona, Verona, Italy

*Correspondence to: Christian Alexander Gutschow, Department for General and Transplantation Surgery, University Hospital Zurich, Raemistrasse 100, Zurich, CH-8091, Switzerland (e-mail: christian.gutschow@usz.ch;  @ChristianGutsc2)

Members of the GastroBenchmark Consortium and the GASTRODATA Consortium are co-authors of this study and are listed under the heading Collaborators.

Presented to the European Surgical Association Congress (Bordeaux, France, May 2023), the Swiss Society of Surgery Congress (Basel, Switzerland, June 2023), the Swiss Society of Gastroenterology and Visceral Surgery Congress (Interlaken, Switzerland, September 2023), the Korean International Gastric Cancer Week (Seoul, Korea, September 2023) and the European Gastric Cancer Congress (Geneva, Switzerland, October 2023).

Introduction

Gastric cancer is the fifth most common cancer worldwide and ranks third for cancer-related death^{1,2}. Curative surgery with or without perioperative chemotherapy remains the therapeutic cornerstone^{3,4}; however, oncological gastric resections require particular skills and experience, have a clear correlation between case volume and outcome⁵, and a relevant learning curve⁶.

Oncological gastrectomy is associated with significant postoperative morbidity, with complication and mortality rates ranging from 11% to 46% and 3% to 20% respectively^{7–12}. The wide discrepancy in reported outcomes is due to the fact that the current evidence on morbidity after gastrectomy is derived from patients who underwent surgery in centres with variable experience and caseloads. In addition, the current definitions of postoperative morbidity are inconsistent and the documentation quality is variable. Recently, the Gastrectomy Complications Consensus Group (GCCG) curated a list of defined complications after gastrectomy⁹, which was an important step toward standardized outcome reporting. Nevertheless, it can be assumed that the available data provide only a rough guide and do not allow surgeons or centres to compare their own results.

With this in mind, this multicentre benchmark analysis of oncological gastrectomy was conceived. Benchmarking involves identifying a point of reference against which a third party's performance can be compared. By definition, a benchmark describes a 'best possible' outcome under ideal conditions¹³. Benchmarking is a well-established management tool for evaluating efficiency and productivity¹⁴. With a growing need to monitor outcomes, benchmarking is increasingly used in the

field of surgery, especially for complex and cost-intensive procedures^{15–24}. In the present study, data were derived from selected 'optimal' patients with low co-morbidity, managed by expert institutions on five continents. To account for disease-related and patient-related differences between East Asian and European/American centres, separate benchmark values were calculated per world region. These results can be implemented as a reference owing to the novelty and the relevance of the benchmark concept and the lack of comparable outcome values for oncological gastrectomy in the literature.

Methods

Ethics

This study involving data of human participants was performed in accordance with the ethical standards of the respective institutional and/or national research committees and with the 1964 Declaration of Helsinki and its later amendments or comparable ethical standards. Approval from the ethics committees of the leading centre in Zurich in Switzerland (BASEC No. 2022-00931) and each participating centre was obtained before patient inclusion.

Data collection

Study design and calculation of benchmark values followed a recommended standardized and validated methodology^{13,25–27} and was performed in collaboration with the GASTRODATA group^{9,10}.

Planned centre inclusion criteria were an average annual caseload of ≥ 20 oncological gastrectomies, the availability of a prospective database, and a special commitment to upper gastrointestinal surgery as documented by recent publications.

Received: August 14, 2023. Revised: October 04, 2023. Accepted: October 04, 2023

© The Author(s) 2024. Published by Oxford University Press on behalf of BJS Foundation Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

Table 1 Baseline demographic data of the benchmark cohort, stratified by type of surgery

	Total Gastrectomy (n = 498)	Distal Gastrectomy (n = 1071)	P	Total (n = 1569)
Age (years), median (i.q.r.)	54 (47–60)	55 (48–60)	0.137	55 (48–60)
Sex				
Female	37.3	39.9	0.348	39.1
Male	62.7	60.1		60.9
BMI (kg/m ²), median (i.q.r.)	23.5 (21.2–25.5)	23.3 (21.1–25.5)	0.313	23.3 (21.1–25.5)
ASA grade				
I	51.0	70.6	<0.001	64.4
II	49.0	29.4		35.6
World region				
America	10.2	4.5	<0.001	6.3
East Asia	45.0	79.1		68.3
Europe	44.8	16.4		25.4
pTNM staging				
0/I	37.9	65.0	<0.001	56.4
II	29.7	19.2		22.6
III	32.3	15.8		21.0
IV	0	0		0
Tumour localization				
Cardia/OGJ/fundus	33.3	0	<0.001	10.6
Corpus	52.0	48.3		49.5
Antrum/pylorus	10.6	51.2		38.3
Multiple, whole, linitis plastica	4.0	0.6		1.7
Surgical access				
Open	55.6	20.4	<0.001	31.5
Laparoscopic	37.3	66.5		57.2
Robotic	4.4	11.9		9.5
Conversion (laparoscopic to open)	2.6	1.3		1.7
Preoperative chemotherapy	41.0	11.6	<0.001	20.9

Values are % apart from those for age and BMI. i.q.r., interquartile range; OGJ, oesophagogastric junction.

Table 2 Postoperative outcomes for the benchmark cohort, stratified by type of surgery

	Total gastrectomy (n = 498)	Distal gastrectomy (n = 1071)	P	Total (n = 1569)
Duration of hospital stay (days), median (i.q.r.)	10 (8–13)	10 (8–11)	0.023	10 (8–12)
Lymph nodes resected (n), median (i.q.r.)	41 (29–54)	39 (30–50)	0.001	40 (30–51)
Resection margin				
R0	96.6	98.9	0.005	98.2
R1	3.0	0.7		1.5
R2	0.4	0.4		0.3
Blood transfusion	3.8	2.1	0.063	2.7
Escalation of care	3.2	0.7	<0.001	1.5
Reoperation	4.2	0.7	<0.001	1.8
Readmission	3.4	1.2	0.002	1.9
Overall morbidity				
None	78.7	86.2	<0.001	83.8
Minor (CD grade I–II)	10.4	7.8		8.7
Major (CD grade ≥IIIA)	10.8	6.0		7.5
CCI*, median (i.q.r.)	26.2 (20.9–29.1)	20.9 (20.9–26.2)	0.001	20.9 (20.9–26.2)
Specific complications				
Anastomotic leakage	3 (O-J)	0.5 (G-J)		2.7
Duodenal stump leakage	1	0.2		0.4
Pancreatic fistula	0.6	0.7		0.7
Lymphatic fistula	1.2	0.7		0.8
Ileus	0.6	2		1.5
Pneumonia/pulmonary complications	3.8	0.7		1.7
Wound infections	3.4	2.2		2.5
Fluid collections	2.8	1.4		1.8
Mortality				
30-day	0.8	0.1	0.038	0.3
90-day	1	0.3	0.117	0.5

Values are % apart from those for duration of hospital stay, lymph nodes resected, and CCI. *Depicted CCI values are calculated only in patients with complications. i.q.r., interquartile range; CD, Clavien–Dindo; CCI, comprehensive complication index; O-J, oesophagojejunostomy; G-J, gastrojejunostomy.


Consecutive patients undergoing gastric resections for adenocarcinoma between 1 January 2017 and 31 December 2021 were included in the overall data collection ([Appendix S1](#)).

De-identified patient-specific data were submitted via secured file transfer and then audited for completeness. In


agreement with each participating centre, no data were reported with patient or hospital identifiers. The information collected included basic demographics, ASA and WHO/Eastern Cooperative Oncology Group (ECOG) grades, tumour-specific parameters, technical details of the surgical procedure, and

a Benchmark patients**Overall complications after gastrectomy: OR (95% c.i., P value)**

Age	–	1.02 (1.01, 1.04, $P = 0.007$)
Sex	Female	–
	Male	1.34 (1.02, 1.78, $P = 0.039$)
World region	East asia	–
	Europe & America	1.17 (0.75, 1.80, $P = 0.484$)
Hospital caseload	High volume	–
	Intermediate volume	0.51 (0.31, 0.84, $P = 0.009$)
	Low volume	–
BMI	–	0.98 (0.94, 1.03, $P = 0.456$)
ASA grade	I	–
	II	0.99 (0.73, 1.35, $P = 0.960$)
TNM staging	0	–
	I	0.67 (0.31, 1.64, $P = 0.344$)
	II	0.96 (0.44, 2.35, $P = 0.929$)
	III	1.29 (0.59, 3.16, $P = 0.542$)
Type of surgery	Total gastrectomy	–
	Subtotal gastrectomy	0.68 (0.49, 0.93, $P = 0.015$)
Surgical technique	Open	–
	Laparoscopic	1.01 (0.72, 1.43, $P = 0.933$)
	Robotic	0.78 (0.44, 1.33, $P = 0.367$)

**b Non-Benchmark patients****Overall complications after gastrectomy: OR (95% c.i., P value)**

Age	–	1.01 (1.01, 1.02, $P < 0.001$)
Sex	Female	–
	Male	1.20 (1.05, 1.37, $P = 0.008$)
World region	East asia	–
	Europe & America	1.29 (1.04, 1.60, $P = 0.018$)
Hospital caseload	High volume	–
	Intermediate volume	1.05 (0.90, 1.22, $P = 0.563$)
	Low volume	0.90 (0.67, 1.20, $P = 0.468$)
BMI	–	1.02 (1.01, 1.04, $P = 0.0002$)
ASA grade	I	–
	II	1.08 (0.90, 1.31, $P = 0.411$)
	III	1.44 (1.16, 1.80, $P = 0.001$)
	IV	2.27 (1.40, 3.70, $P = 0.001$)
	V	–
TNM staging	0	–
	I	1.23 (0.83, 1.85, $P = 0.315$)
	II	1.40 (0.95, 2.11, $P = 0.098$)
	III	1.54 (1.05, 2.32, $P = 0.032$)
	IV	2.25 (1.46, 3.51, $P < 0.001$)
Type of surgery	Total gastrectomy	–
	Subtotal gastrectomy	0.56 (0.49, 0.65, $P < 0.001$)
	Pylorus-preserving gastrectomy	0.74 (0.49, 1.10, $P = 0.142$)
	Proximal gastrectomy	0.97 (0.71, 1.31, $P = 0.829$)
Surgical technique	Open	–
	Laparoscopic	0.79 (0.68, 0.92, $P = 0.003$)
	Robotic	0.51 (0.34, 0.74, $P = 0.001$)

**Fig. 1** Factors predicting occurrence of complications after gastrectomy

a OR plot of multivariable logistic regression model assessing predictive factors for occurrence of overall complications after gastrectomy in benchmark patients (1569 patients). **b** OR plot of multivariable logistic regression model assessing predictive factors for occurrence of overall complications after gastrectomy in non-benchmark patients (7787 patients).

postoperative complications. Endpoints for analysis of postoperative events were at 90 days after surgery. Complications were classified as specified in the databases of the Seoul National University Hospital (South Korea)^{28,29} and the GASTRODATA collaborative (European Chapter of the International Gastric Cancer Association)^{9,30} and were

graded according to the Clavien–Dindo (CD) classification³¹. Cumulative morbidity was assessed with the comprehensive complication index (CCI), which expresses morbidity on a continuous scale from 0 (no complications) to 100 (death) by weighing all postoperative complications according to the CD classification³².

Table 3 Benchmark values for oncological gastrectomy

	Total gastrectomy			Distal gastrectomy		
	Global (n = 498)	East Asia (n = 224)	Europe/America (n = 264)	Global (n = 1071)	East Asia (n = 847)	Europe/America (n = 222)
Duration of hospital stay (days)	≤11	≤11	≤11	≤10	≤10	≤10
R0 resection	≥91.6	≥99.4	≥91.6	≥99.8	≥99.2	100
Lymph node yield (n)	≥27	≥49	≥26	≥24	≥39	≥24
Blood transfusion	≤8.3	≤1.0	≤8.3	0	≤2.0	0
Escalation of care	≤6.6	≤1.0	≤7.1	0	≤0.6	0
Reoperation	≤7.8	0	≤8.1	0	≤0.2	0
Readmission	≤7.4	≤1.8	≤7.6	≤0.2	≤0.4	0
Any complications	≤20.0	≤21.2	≤20.0	≤15.1	≤14.8	≤14.6
Major complications	≤10.0	≤11.3	≤9.3	≤3.4	≤6.5	≤2.2
CCI*	≤33.7	≤24.8	≤51.4	≤26.2	≤24.8	≤28.0
Mortality, 30-day	0	0	0	0	0	0
Mortality, 90-day	0	0	0	0	0	0

Values are % apart from those for duration of hospital stay, lymph node yield, and CCI. *Depicted CCI values are calculated only in patients with complications. CCI, comprehensive complication index.

Table 4 Validation of benchmark values in different cohorts

	Benchmark value (TG global)	Benchmark cohort (n = 1796)	Non-benchmark (>65 years) (n = 998)	Non-benchmark (co-morbidities) (n = 505)	D3 LAD (n = 48)	HIPEC (n = 105)	Splenectomy (n = 203)	Pancreatic/multivisceral resection (n = 400)
Duration of hospital stay (days), median (i.q.r.)	≤11	10 (9–11)	10 (9–13)	10 (9–13)	13 (10–19)	14 (10–21)	12 (9–19)	12 (8–19)
R0 resection	≥91.6	98.4	98.5	99.2	97.9	74.3	91.1	85.8
Lymph node yield (n), median (i.q.r.)	≥27	40 (30–50)	36 (26–48)	34 (26–45)	31 (23–42)	38 (26–46)	46 (28–59)	33 (23–46)
Blood transfusion	≤8.3	2.4	5.6	6.1	6.3	21.0	17.2	22.3
Escalation of care	≤6.6	1.3	2.4	4.0	4.2	8.6	9.4	9.5
Reoperation	≤7.8	1.6	2.3	2.4	8.3	13.3	7.4	12.8
Readmission	≤7.4	1.7	2.3	1.8	0	10.5	3.0	7.3
Any complications	≤20.0	16.3	22.5	19.8	43.8	60.0	42.4	45.3
Major complications	≤10.0	7.5	12.2	12.5	18.8	35.2	23.2	25.0
CCI*, median (i.q.r.)	≤33.7	20.9 (20.9–26.2)	26.2 (20.9–33.5)	26.2 (20.9–26.2)	26.2 (20.9–33.7)	26.2 (20.9–41.1)	26.2 (20.9–37.1)	26.2 (20.9–42.4)
Mortality, 30-day	0	0.3	0.3	0.4	2.1	2.9	2.5	4.5
Mortality, 90-day	0	0.4	0.3	0.6	2.1	4.8	3.4	6.3

Values are % apart from those for duration of hospital stay, lymph node yield, and CCI. *Depicted CCI values are calculated only in patients with complications. TG, total gastrectomy; LAD, lymphadenectomy; HIPEC, hyperthermic intraperitoneal chemotherapy; i.q.r., interquartile range; CCI, comprehensive complication index.

Inclusion criteria for low co-morbidity

Benchmark patients were defined as aged 18–65 years, with an ASA grade ≤ II, an ECOG ≤ 1, a BMI between 18 and 30 kg/m², and no significant co-morbidities (see [Appendix S2](#)). Only patients after total gastrectomy (TG) or distal/subtotal gastrectomy (DG) with adequate lymphadenectomy (D1+, D2, or D2+) from centres that had logged at least 50 cases were eligible for the current benchmark analysis ([Table S1](#)). Exclusion criteria were histology other than adenocarcinoma, other limited types of gastric resections, concurrent major organ resections, administration of hyperthermic intraperitoneal chemotherapy (HIPEC), relevant prior abdominal surgery, and advanced tumour stage (pT4b, pM1).

Performance metrics of benchmarking

Primary outcome measures for benchmark analysis of surgical quality were duration of hospital stay, rate of R0 resection, number of lymph nodes resected, rates of blood transfusion, escalation of care, reoperation, readmission, and overall and

major (CD grade ≥ IIIA) morbidity, cumulative morbidity as measured by the CCI (all at 30 days), and rates of 30- and 90-day mortality.

According to the benchmarking methodology^{13,25–27}, median values of continuous variables and the proportions of categorical variables were calculated per participating centre. Benchmark cut-offs, indicating 'best achievable' results for each outcome indicator, were set at the 25th percentile for positive parameters (for example lymph node yield) or the 75th percentile for negative parameters (for example complications) of the centres' median values.

Statistical analysis

Statistical significance was defined as $P < 0.050$. Categorical variables are presented as number or % and were compared using Fisher's exact test and numerical variables are expressed as median and interquartile range (i.q.r.) and were compared using Wilcoxon's rank sum test. Predictive factors were identified by multivariable logistic regression with respective ORs with 95% confidence intervals. R version 4.0.2 (R Foundation

for Statistical Computing, Vienna, Austria) was used for statistical analyses and figures.

Results

Demographic characteristics of benchmark patients

From a patient cohort of 9356 oncological gastrectomies performed at 43 centres from five continents (Europe, Asia, North America, South America, and Africa; see [Appendix S3](#)), 1569 low-risk patients (16.8%) from 32 centres ([Table S1](#)) with a median age of 55 years undergoing TG (498 patients) and DG (1071 patients) were selected according to the benchmark criteria. Minimally invasive (laparoscopic or robotic) surgery (MIS) was performed in 66.7% (TG 41.7% and DG 78.4%; $P < 0.001$), with higher rates for early tumour stages, females, and East Asian patients (all $P < 0.001$). Overall and procedure-specific baseline characteristics of benchmark patients are detailed in [Table 1](#).

Perioperative outcomes for benchmark patients

The median postoperative duration of hospital stay was 10 days after both TG (i.q.r. 8–13) and DG (i.q.r. 8–11) and the median number of lymph nodes resected was 41 (i.q.r. 29–54) in TG and 39 (i.q.r. 30–50) in DG. R0 resection rates were 96.6% for TG and 98.9% for DG. Detailed perioperative outcomes for benchmark patients are shown in [Table 2](#).

In the benchmark cohort, 265 complications in 243 patients were recorded, accounting for an overall morbidity rate of 16.2% (TG 21.3% and DG 13.8%; $P < 0.001$). The rate of major complications (CD grade \geq IIIA) was higher after TG (10.8%) than after DG (6.0%) with a median CCI of 26.2 (i.q.r. 20.9–29.1) after TG and 20.9 (i.q.r. 20.9–26.2) after DG. The 30- and 90-day mortality rates were 0.3% and 0.5% respectively. The most common complications in benchmark patients were anastomotic leakage (occurring in 2.7% (3% oesophagojejunostomy after TG and 0.5% gastrojejunostomy after DG)), wound infections (occurring in 2.5%), abdominal fluid collections (occurring in 1.8%), pneumonia/pulmonary complications (occurring in 1.7%), ileus (occurring in 1.5%), and lymphatic (0.8%) and pancreatic (0.7%) fistulae. Duodenal stump leakage occurred in 0.4% (1% after TG and 0.2% after DG). [Table 2](#) shows the complication rates for the surgical procedures.

Benchmark versus non-benchmark patients

In the overall cohort, East Asian patients were younger, had a lower BMI, had fewer co-morbidities (ASA grade \leq II), and had earlier-stage tumours (all $P < 0.001$) compared with European and American patients, resulting in a higher proportion of benchmark patients from East Asia (20.6%) than from Europe and America (12.0%; [Table S2](#)). Consequently, the benchmark cohort consisted of a higher proportion of East Asian patients (1071 patients; 68.3%), patients who were younger, a higher proportion of women, and patients who had earlier-stage tumours than the non-benchmark cohort ([Table S3](#)). Relevant endpoints, such as oncological parameters and rates of readmission, escalation of care, and mortality, were significantly worse for non-benchmark patients ([Table S4](#)). Likewise, rates of specific complications, such as anastomotic leakage (overall 4.9%, TG 6.1%, and DG 1.6%) and duodenal stump leakage (overall 1.7%, TG 1.9%, and DG 1.5%), were higher in non-benchmark patients. Wound infections were the most frequent complications in the non-benchmark cohort (6.1%) and pneumonia/pulmonary complications (5.7% overall) were significantly more frequent in Europe/America (8.3%)

compared with East Asia (2.4%) and in TG (9.7%) compared with DG (2.8%).

Multivariable logistic regression revealed a significant association of overall complication rates with older age and male sex in both benchmark and non-benchmark patients. In contrast, centre localization, tumour stage, and surgical technique did not influence the morbidity of benchmark patients, whereas non-benchmark patients from East Asian centres or with early tumour stages had lower complication rates ([Fig. 1](#)). Whilst there was an overall association of MIS with reduced complication rates, a sensitivity analysis revealed that MIS was associated with decreased complications in East Asian centres only. In contrast, complication rates for European/American centres were associated with higher co-morbidity, but not with MIS ([Fig. S1](#)). Furthermore, besides fewer co-morbidities, surgery performed at an East Asian centre was the strongest predictor of favourable outcomes in terms of major complication and mortality rates ([Fig. S2](#)).

Benchmark values

In addition to procedure-specific global benchmark cut-offs for TG and DG, separate regional benchmarks for East Asia and Europe/America were calculated, to account for the demographic differences between the European/American and East Asian patient cohorts outlined above.

For TG, benchmark values for duration of hospital stay were ≤ 11 days, with relevant differences between East Asia and Europe/America regarding R0 resection rate ($\geq 99.4\%$ versus $\geq 91.6\%$ respectively) and lymph node yield (≥ 49 versus ≥ 26 respectively). Furthermore, complication-associated cut-offs, such as those for rates of blood transfusion ($\leq 8.3\%$ versus $\leq 1.0\%$ respectively), escalation of care ($\leq 7.1\%$ versus $\leq 1.0\%$ respectively), reoperation ($\leq 8.1\%$ versus 0% respectively), and readmission ($\leq 7.6\%$ versus $\leq 1.8\%$ respectively), were less demanding for European/American centres compared with East Asian cut-offs. Interestingly, the benchmark values for overall ($\leq 20.0\%$ for European/American centres versus $\leq 21.2\%$ for East Asian centres) and major ($\leq 9.3\%$ for European/American centres versus $\leq 11.3\%$ for East Asian centres) complication rates were comparable between the two regions; however, the CCI cut-off was markedly increased in European/American centres compared with East Asian centres (≤ 51.4 versus ≤ 24.8 respectively). In contrast, no prominent difference in achievable cut-offs between Eastern and Western centres was observed for DG, with the exception of a higher number of lymph nodes to be resected in East Asia (≥ 39 versus ≥ 24). Lastly, 30- and 90-day mortality rates for both interventions should be 0% globally (see [Table 3](#) for procedure-specific and subdivided benchmarks for East Asia versus Europe/America for TG and DG).

Validation of benchmark values

The global benchmark values for TG were validated in defined subgroups of the total cohort: patients fulfilling all benchmark criteria, but being older than 65 years (998 patients) or having one or more 1 major co-morbidity ([Appendix S2](#); 505 patients), undergoing D3 lymphadenectomy (48 patients), being treated with HIPEC (105 patients), undergoing simultaneous splenectomy (203 patients), or undergoing pancreatic/multivisceral resection (400 patients). Patient cohorts with extended benchmark criteria (older than 65 years or major co-morbidities) were within benchmark cut-offs, except for slightly elevated rates of overall and major complications, indicating that acceptable outcomes are achievable in high-volume centres, even for 'suboptimal' patients. In contrast,

the four patient cohorts with extended surgical procedures strongly exceeded benchmark thresholds for duration of hospital stay, R0 resection rates, morbidity-associated parameters, and mortality rates (Table 4), confirming that the benchmark values of the present study are sensitive to extended resections performed in patients with increased co-morbidity and advanced tumour stage.

Discussion

In recent years there has been growing emphasis on the assessment of surgical outcomes to evaluate the quality of care provided by hospitals, departments, and individual surgeons. These data are often made public and used to compare the performance of different providers to improve transparency and patient autonomy³³. This information has significant economic implications for healthcare providers, as it is often used by healthcare insurers, private payers, policymakers, the media, physicians, and patients to guide decisions. As a result, the assessment of the quality of patient care has become an important aspect of the healthcare sector's efforts regarding public relations. However, there are concerns that the evidence provided by large national databases, audits, and meta-analyses may be biased due to the heterogeneity of the target populations and procedures analysed^{34,35}. To increase sample sizes, inclusion criteria for these studies are often broad and cases are not risk adjusted, resulting in ranking systems that are inaccurate and misunderstood. In addition, this may lead to risk aversion among healthcare providers who may avoid highly morbid or complex cases³⁶. Other weaknesses of traditional databases include the lack of uniform data sets, the absence of consistent validation methods, and the focus on single outcome parameters, such as the 30-day mortality rate².

Benchmarking is a method for comprehensive evaluation of surgical procedures and has not previously been used in gastric cancer surgery. The aim of the present study was to address the limitations of previous research by providing benchmarks for multiple clinically relevant endpoints that can be used by third-party institutions. A particular strength of the present study is that it is based on the largest international cohort of patients undergoing oncological gastrectomy at high-volume institutions across several continents. Restriction to expert centres with comprehensive prospective gastrectomy databases and systematic classification of postoperative complications results in a high-quality data set with only minimal missing data. Using a highly selected group of patients with low co-morbidity, this approach allows the definition of global benchmark parameters, or 'best achievable' results, for TG and DG for the first time. Furthermore, as patient populations with gastric cancer show pronounced differences between East Asia and Europe/America, the calculation of separate benchmark values accounting for regional differences is an important aspect of the present study^{37,38}. The values show considerable regional differences and it is evident that, given higher rates of co-morbidities, more advanced tumour stages, and differences in caseload and centralization, the same outcomes cannot be expected in European/American centres compared with high-volume East Asian centres. The application of the respective regional benchmark is therefore advocated as a realistic and achievable outcome metric.

The postoperative morbidities of benchmark patients (overall 16.2% and major 7.5%) and the whole study cohort (overall 25.2% and major 13.3%) are consistent with the published literature. RCTs from Korea reported overall morbidity rates between 13%

and 24%^{28,29,39}, which are similar to outcome data from retrospective Western and Eastern series^{10,40}. On the other hand, three European RCTs^{41–43} reported considerably higher overall morbidity (34%–44%). Major morbidity in the benchmark cohort is lower than in European RCTs, but higher than in patients with locally advanced cancer from East Asian high-volume centres^{44,45}. Nevertheless, the global benchmark cut-offs of the present study for overall morbidity ($\leq 20\%$) and CCI (≤ 33.7) in TG are comparable to rates of postoperative morbidity in retrospective European series¹⁰, whilst benchmark values for readmission rate ($\leq 7.4\%$), lymph node yield (≥ 27), and R0 resection rate ($\geq 91.6\%$) are similar to the 9.1%–9.6% readmission rate, 29 lymph nodes, and 95% R0 resection rate reported in the LOGICA⁴¹ trial, suggesting that the global and European/American benchmarks in particular represent generally achievable goals in unselected Western patient cohorts.

Anastomotic leakage, the most frequent complication in benchmark patients, is still one of the most dreaded complications after gastrectomy, with a reported incidence between 6% and 10%^{41,42,46} and a mortality rate of up to 25%¹¹ in Western centres. With respect to overall and major morbidity, it is assumed that the lower rates for the current data set (whole cohort 4.9% and benchmark cohort 2.7%) are due to an important contribution of East Asian centres to the data set, as the rate of anastomotic insufficiency in the cohort of the present study was lower in East Asian patients (2.6%) than European patients (7.0%), similar to previous reports^{29,47}. Likewise, the low rate of pneumonia/pulmonary complications (1.7%) in benchmark patients is due to a low incidence in East Asian patients (benchmark: 0.8%, overall: 2.4%) compared with European/American patients (benchmark: 3.6%, overall: 8.3%). Importantly, however, these different rates are in line with published results of pneumonia in retrospective series of unselected European patients (8.3%)¹⁰ or prospective European RCTs, such as LOGICA (11.3%)⁴¹ and STOMACH (6.1%–8.5%)⁴², as well as published rates of high-quality prospective East Asian RCTs, such as JCOG0912 (0.9%)⁴⁴, JCOG1001 (2%)⁴⁸, KCLASS-01 (0.7%–1.6%)²⁸, and KCLASS-02 (2.7%–3.5%)²⁹. Under-reporting of these specific complications is therefore unlikely. Furthermore, rates of other specific complications, such as bowel obstruction, are within the range of published results⁴⁹.

In Europe, the mortality rate after gastrectomy ranges between 3% and 5% in expert institutions and up to 20% in low-volume centres^{8,30,41,42}. Mortality rates of the cohort in the present study are considerably lower for both benchmark and non-benchmark patients, which again are partially explained by a relevant contribution from East Asian centres where the mortality rate is typically less than 1%^{28,29,50}, even in advanced cancer. Nevertheless, the low 30- and 90-day mortality rates for the whole cohort (1.2% and 1.8% respectively) also reflect the high quality of postoperative care and effective complication management in contributing Western centres and show that optimal results can be achieved in expert centres from all over the world, regardless of tumour stage and surgical approach.

Validation of the benchmark values of the present study confirms a good correlation between surgical invasiveness and morbidity, confirming previous findings from RCTs investigating the role of splenectomy⁵¹, extended lymphadenectomy⁵², or HIPEC⁵³ in gastric cancer surgery.

There are several limitations to the present study. First, there is substantial variability in the number of cases included per centre. Due to the increased incidence of gastric cancer in East Asia and centralization of services, caseloads from the centres in Seoul

and Tokyo are considerably higher than the remaining institutions (mainly in Europe and America), which might make the analysis prone to bias. Whilst this variability may be considered a strength, better reflecting reality than a single high-volume experience, it may also indicate that differences in experience with a specific procedure and learning curve-related morbidity can have an impact on the data set. In this context, it is important to emphasize that separate benchmarks for East Asian and European/American centres were calculated, to account for underlying differences in patient demographics and centre volume. Furthermore, no correlation between morbidity and centre volume was found for benchmark patients, whereas major complication and mortality rates were lower for East Asian non-benchmark patients.

Second, misclassification or under-reporting of complications could be another source of error. CD grade 1 complications are not evenly distributed among centres, suggesting that minor morbidities, such as urinary tract infections, are under-reported. However, CD grade 1 complications only have a minimal effect on the CCI and will therefore not significantly affect the results of the present study. In addition, it cannot be excluded that different definitions of complications were used by the participating centres. Certain complications, such as ileus or lymphatic fistula, are not recorded in the GASTRODATA data set and might therefore be under-reported. The reported rates of specific complications, such as pneumonia and anastomotic insufficiency, however, compare well with rates published in the literature; therefore, it is not thought that systematic under-reporting of relevant complications occurred in the retrospective data collection of the present study.

In conclusion, this study is the first to provide benchmark values for outcomes after oncological gastrectomy with lymphadenectomy. The data were collected from a large group of patients undergoing surgery in expert centres from various countries. By analysing a subgroup of 'ideal' patients with low co-morbidity, 'best possible' results for TG and DG were obtained. It is important to note that the results of the present study represent a snapshot of the current situation and that benchmark values for gastrectomy may change upon wider adoption of minimally invasive and particularly robotic techniques. Nevertheless, the results of the present study can be used as a reference due to the novelty and the relevance of the benchmark concept and the lack of comparable outcome values in the current literature.

Collaborators

GastroBenchmark Consortium

Bas P. L. Wijnhoven (Department of Surgery, Erasmus University Medical Center, Rotterdam, The Netherlands); Hidde Overtom (Department of Surgery, Erasmus University Medical Center, Rotterdam, The Netherlands); Ines Gockel (Department of Visceral, Transplant, Thoracic and Vascular Surgery, University Hospital of Leipzig, Leipzig, Germany); René Thieme (Department of Visceral, Transplant, Thoracic and Vascular Surgery, University Hospital of Leipzig, Leipzig, Germany); Ewen A. Griffiths (Department of Upper GI Surgery, Queen Elizabeth Hospital, University Hospitals Birmingham NHS Foundation Trust, Birmingham, UK); William Butterworth (Department of Upper GI Surgery, Queen Elizabeth Hospital, University Hospitals Birmingham NHS Foundation Trust, Birmingham, UK); Henrik Nienhüser (Klinik für Allgemein-, Viszeral- und Transplantationschirurgie, Universitätsklinikum Heidelberg, Heidelberg, Germany); Beat Müller (Klinik für

Allgemein-, Viszeral- und Transplantationschirurgie, Universitätsklinikum Heidelberg, Heidelberg, Germany); Nerma Crnovrsanin (Klinik für Allgemein-, Viszeral- und Transplantationschirurgie, Universitätsklinikum Heidelberg, Heidelberg, Germany); Felix Nickel (Department of General, Visceral, and Thoracic Surgery, University Medical Center Hamburg-Eppendorf, Hamburg, Germany); Suzanne Gisbertz (Amsterdam University Medical Center & Cancer Center Amsterdam, University of Amsterdam, Department of Surgery, Amsterdam, The Netherlands); Mark van Berge Henegouwen (Amsterdam University Medical Center & Cancer Center Amsterdam, University of Amsterdam, Department of Surgery, Amsterdam, The Netherlands); Philip H. Pucher (Department of Surgery, Queen Alexandra Hospital, Portsmouth Hospitals NHS Trust, Portsmouth, UK); Kashuf Khan (Department of Surgery, Queen Alexandra Hospital, Portsmouth Hospitals NHS Trust, Portsmouth, UK); Asif Chaudry (The Royal Marsden NHS Foundation Trust, Chelsea, London, SW3 6JJ, UK); Pranav H. Patel (The Royal Marsden NHS Foundation Trust, Chelsea, London, SW3 6JJ, UK); Manuel Pera (Section of Gastrointestinal Surgery, Hospital Universitario del Mar, Universitat Autònoma de Barcelona, Barcelona, Spain); Mariagiulia Dal Cero (Section of Gastrointestinal Surgery, Hospital Universitario del Mar, Universitat Autònoma de Barcelona, Barcelona, Spain); Carlos Garcia (Hospital San Borja Arriarán, Av. Sta. Rosa 1234, Santiago, Región Metropolitana, Chile); Guillermo Martinez Salinas (Hospital San Borja Arriarán, Av. Sta. Rosa 1234, Santiago, Región Metropolitana, Chile); Paulo Kassab (Gastroesophageal and Bariatric Surgical Division, Department of Surgery, Santa Casa of São Paulo Medical School and Hospital, São Paulo, Brazil); Osvaldo Antônio Prado Castro (Gastroesophageal and Bariatric Surgical Division, Department of Surgery, Santa Casa of São Paulo Medical School and Hospital, São Paulo, Brazil); Enrique Norero (Esophagogastric Surgery Unit, Digestive Surgery Department, Hospital Dr Sotero del Rio, Pontificia Universidad Católica de Chile, Santiago, Chile); Paul Wisniowski (Division of Upper GI and General Surgery, Keck School of Medicine, University of Southern California, 1510 San Pablo St., Health Sciences Campus, Los Angeles, USA); Luke Randall Putnam (Division of Upper GI and General Surgery, Keck School of Medicine, University of Southern California, 1510 San Pablo St., Health Sciences Campus, Los Angeles, USA); Pietro Maria Lombardi (Division of Minimally Invasive Surgical Oncology, Niguarda Cancer Center, ASST Grande Ospedale Metropolitano Niguarda, Piazza Ospedale Maggiore, 3, 20162, Milan, Italy); Giovanni Ferrari (Division of Minimally Invasive Surgical Oncology, Niguarda Cancer Center, ASST Grande Ospedale Metropolitano Niguarda, Piazza Ospedale Maggiore, 3, 20162, Milan, Italy); Rita Gudaityte (Department of Surgery, Hospital of Lithuanian University of Health Sciences, Eiveniu 2, Kaunas 50161, Lithuania); Almantas Maleckas (Department of Surgery, Hospital of Lithuanian University of Health Sciences, Eiveniu 2, Kaunas 50161, Lithuania); Leanne Prodehl (Department of Surgery, Charlotte Maxeke Johannesburg Academic Hospital, University of the Witwatersrand, Johannesburg, South Africa); Antonio Castaldi (Service de Chirurgie Digestive et Cancérologie Digestive, Hôpital Universitaire Carêmeau, Nîmes, France); Michel Prudhomme (Service de Chirurgie Digestive et Cancérologie Digestive, Hôpital Universitaire Carêmeau, Nîmes, France); Simone Giacopuzzi (Department of Surgery, University Hospital of Verona, Verona, Italy); Francesco Puccetti (Department of Surgery, San Raffaele Hospital, Milano, Italy); Domenico D'Ugo (FONDAZIONE POLICLINICO UNIVERSITARIO GEMELLI-IRCCS, Roma, Italy); Daniel Gero (Department of Surgery & Transplantation,

University Hospital Zürich, Raemistrasse 100, 8091 Zurich, Switzerland); Hyuk-Joon Lee (Department of Surgery, Seoul National University Cancer Hospital, 101 Daehak-ro Jongno-gu, Seoul, South Korea).

GASTRODATA Consortium

Guillaume Piessen (Department of Surgery, University Hospital of Lille, Lille, France); Justine Lerooy (Department of Surgery, University Hospital of Lille, Lille, France); Johanna Wilhelmina van Sandick (Department of Surgical Oncology, The Netherlands Cancer Institute—Antoni van Leeuwenhoek Hospital, Postbus, 90203 1006 BE, Amsterdam, The Netherlands); John V. Reynolds (Department of Surgery, St. James's Hospital, Trinity College Dublin, Dublin, Ireland); Paolo Morgagni (GB Morgagni-L Pierantoni Hospital, Forlì, Italy); Arnulf H. Hölscher (Contilia Center for Esophageal Diseases, Elisabeth Hospital Essen, West German Tumor Center, University Medicine Essen, Germany); Martin Hemmerich (Contilia Center for Esophageal Diseases, Elisabeth Hospital Essen, West German Tumor Center, University Medicine Essen, Germany); Stefan Mönig (Department of Surgery, University Hospital of Geneva, Geneva, Switzerland); Mickael Chevally (Department of Surgery, University Hospital of Geneva, Geneva, Switzerland); Piotr Kołodziejczyk (Department of Surgery, Jagiellonian University, Kraków, Poland); Henk Hartgrink (Leiden University Medical Center, Leiden, The Netherlands); Paulo Matos da Costa (Faculdade de Medicina, Universidade de Lisboa; Lisboa, Portugal); Filipe Castro Borges (Faculdade de Medicina, Universidade de Lisboa; Lisboa, Portugal); Andrew Davies (Department of Surgery, Guy's & St Thomas' NHS Foundation Trust, London, UK); Cara Baker (Department of Surgery, Guy's & St Thomas' NHS Foundation Trust, London, UK); William Allum (The Royal Marsden NHS Foundation Trust, Chelsea, London, SW3 6JJ, UK); Sacheen Kumar (The Royal Marsden NHS Foundation Trust, Chelsea, London, SW3 6JJ, UK); Wojciech Polkowski (Medical University of Lublin, Lublin, Poland); Karol Rawicz-Pruszyński (Medical University of Lublin, Lublin, Poland); Uberto Fumagalli Romario (Digestive Surgery, European Institute of Oncology, IRCCS, Milano, Italy); Stefano De Pascale (Digestive Surgery, European Institute of Oncology, IRCCS, Milano, Italy); Antonio Tarasconi (Department of Surgery, University Hospital of Brescia, Brescia, Italy); Daniel Reim (Department of Surgery, TUM School of Medicine, Technical University of Munich, Germany); Ilaria Pergolini (Department of Surgery, TUM School of Medicine, Technical University of Munich, Germany); Lucio Lara Santos (Department of Surgery, Portuguese Institute of Oncology, Porto, Portugal); Pedro Carvalho Martins (Department of Surgery, Portuguese Institute of Oncology, Porto, Portugal); Alberto Biondi (FONDAZIONE POLICLINICO UNIVERSITARIO GEMELLI-IRCCS, Roma, Italy); Maurizio Degiuli (Department of Surgical Oncology and Digestive Surgery, San Luigi University Hospital Orbassano, School of Medicine, University of Torino, Torino, Italy); Rossella Reddavid (Department of Surgical Oncology and Digestive Surgery, San Luigi University Hospital Orbassano, School of Medicine, University of Torino, Torino, Italy); Wojciech Kielan (Wrocław Medical University, Wrocław, Poland); Paul Magnus Schneider (Digestive Oncology Tumor Center and Esophageal Cancer Center, Hirslanden Medical Center, Zurich, Switzerland); Thomas Murphy (Mercy University Hospital, Cork, Ireland).

Funding

The authors have no funding to declare.

Acknowledgements

M.A.S., J.K., and F.B. share first authorship. H.-K.Y. and C.A.G. share last authorship.

Author contributions

Marcel André Schneider (Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project administration, Software, Supervision, Validation, Visualization, Writing—original draft, Writing—review & editing), Jeusun Kim (Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project administration, Validation, Visualization, Writing—original draft, Writing—review & editing), Felix Berlth (Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project administration, Supervision, Validation, Visualization, Writing—original draft, Writing—review & editing), Yutaka Sugita (Data curation, Formal analysis, Investigation, Methodology, Project administration, Validation, Writing—review & editing), Peter P. Grimmer (Data curation, Formal analysis, Investigation, Validation, Writing—review & editing), Takeshi Sano (Data curation, Formal analysis, Investigation, Validation, Writing—review & editing), Riccardo Rosati (Data curation, Formal analysis, Investigation, Validation, Writing—review & editing), Gian Luca Baiocchi (Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project administration, Supervision, Validation, Writing—original draft, Writing—review & editing), Maria Bencivenga (Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project administration, Validation, Visualization, Writing—original draft, Writing—review & editing), Giovanni De Manzoni (Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project administration, Supervision, Validation, Visualization, Writing—original draft, Writing—review & editing), Souya Nunobe (Data curation, Formal analysis, Investigation, Methodology, Project administration, Supervision, Validation, Visualization, Writing—original draft, Writing—review & editing), Han-Kwang Yang (Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project administration, Supervision, Validation, Visualization, Writing—original draft, Writing—review & editing), Christian Alexander Gutschow (Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project administration, Supervision, Validation, Visualization, Writing—original draft, Writing—review & editing), the GastroBenchmark Consortium (Data curation, Formal analysis, Investigation, Validation, Writing—review & editing), and the GASTRODATA Consortium (Data curation, Formal analysis, Investigation, Validation, Writing—review & editing)

Disclosure

The authors declare no conflict of interest.

Supplementary material

[Supplementary material](#) is available at BJS online.

Data availability

All research data supporting this publication are available from the corresponding author upon reasonable request.

References

- Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2018;**68**:394–424
- Arnold M, Ferlay J, van Berge Henegouwen MI, Soerjomataram I. Global burden of oesophageal and gastric cancer by histology and subsite in 2018. *Gut* 2020;**69**:1564–1571
- Al-Batran SE, Hofheinz RD, Pauligk C, Kopp HG, Haag GM, Luley KB et al. Histopathological regression after neoadjuvant docetaxel, oxaliplatin, fluorouracil, and leucovorin versus epirubicin, cisplatin, and fluorouracil or capecitabine in patients with resectable gastric or gastro-oesophageal junction adenocarcinoma (FLOT4-AIO): results from the phase 2 part of a multicentre, open-label, randomised phase 2/3 trial. *Lancet Oncol* 2016;**17**:1697–1708
- Al-Batran SE, Homann N, Pauligk C, Goetze TO, Meiler J, Kasper S et al. Perioperative chemotherapy with fluorouracil plus leucovorin, oxaliplatin, and docetaxel versus fluorouracil or capecitabine plus cisplatin and epirubicin for locally advanced, resectable gastric or gastro-oesophageal junction adenocarcinoma (FLOT4): a randomised, phase 2/3 trial. *Lancet* 2019;**393**:1948–1957
- Guller U, Warschkow R, Ackermann CJ, Schmied B, Cerny T, Ess S. Lower hospital volume is associated with higher mortality after oesophageal, gastric, pancreatic and rectal cancer resection. *Swiss Med Wkly* 2017;**147**:w14473
- Brenkman HJF, Claassen L, Hannink G, van der Werf LR, Ruurda JP, Nieuwenhuizen GAP et al. Learning curve of laparoscopic gastrectomy: a multicenter study. *Ann Surg* 2023;**277**:e808–e816
- van der Werf LR, Busweiler LAD, van Sandick JW, van Berge Henegouwen MI, Wijnhoven BPL. Reporting national outcomes after esophagectomy and gastrectomy according to the Esophageal Complications Consensus Group (ECCG). *Ann Surg* 2020;**271**:1095–1101
- Voeten DM, Busweiler LAD, van der Werf LR, Wijnhoven BPL, Verhoeven RHA, van Sandick JW et al. Outcomes of esophagogastric cancer surgery during eight years of surgical auditing by the Dutch Upper Gastrointestinal Cancer Audit (DUCA). *Ann Surg* 2021;**274**:866–873
- Baiocchi GL, Giacomuzzi S, Marrelli D, Reim D, Piessen G, da Costa PM et al. International consensus on a complications list after gastrectomy for cancer. *Gastric Cancer* 2019;**22**:172–189
- Baiocchi GL, Giacomuzzi S, Reim D, Piessen G, Costa PM, Reynolds JV et al. Incidence and grading of complications after gastrectomy for cancer using the GASTRODATA registry: a European retrospective observational study. *Ann Surg* 2020;**272**:807–813
- Baiocchi GL, Giacomuzzi S, Vittimberga G, De Pascale S, Pastorelli E, Gelmini R et al. Clinical outcomes of patients with complicated post-operative course after gastrectomy for cancer: a GIRCG study using the GASTRODATA registry. *Updates Surg* 2023;**75**:419–427
- Baum P, Diers J, Lichthardt S, Kastner C, Schlegel N, Germer CT et al. Mortality and complications following visceral surgery: a nationwide analysis based on the diagnostic categories used in German hospital invoicing data. *Dtsch Arztebl Int* 2019;**116**:739–746
- Barkun J, Clavien P, Pawlik TM. The benefits of benchmarking—a new perspective on measuring quality in surgery. *JAMA Surg* 2023;**158**:341–342
- Ettorchi-Tardy A, Levif M, Michel P. Benchmarking: a method for continuous quality improvement in health. *Health Policy* 2012;**7**:e101–e119
- Breuer E, Mueller M, Doyle MB, Yang L, Darwish Murad S, Anwar IJ et al. Liver transplantation as a new standard of care in patients with perihilar cholangiocarcinoma? Results from an international benchmark study. *Ann Surg* 2022;**276**:846–853
- Gero D, Raptis DA, Vleeschouwers W, van Veldhuisen SL, Martin AS, Xiao Y et al. Defining global benchmarks in bariatric surgery: a retrospective multicenter analysis of minimally invasive Roux-en-Y gastric bypass and sleeve gastrectomy. *Ann Surg* 2019;**270**:859–867
- Gero D, Vannijvel M, Okkema S, Deleus E, Lloyd A, Lo Menzo E et al. Defining global benchmarks in elective secondary bariatric surgery comprising conversational, revisional, and reversal procedures. *Ann Surg* 2021;**274**:821–828
- Mueller M, Breuer E, Mizuno T, Bartsch F, Ratti F, Benzing C et al. Perihilar cholangiocarcinoma—novel benchmark values for surgical and oncological outcomes from 24 expert centers. *Ann Surg* 2021;**274**:780–788
- Müller PC, Breuer E, Nickel F, Zani SJ, Kauffmann E, De Franco L et al. Robotic distal pancreatectomy, a novel standard of care? Benchmark values for surgical outcomes from 16 international expert centers. *Ann Surg* 2023;**278**:253–259
- Muller X, Marcon F, Sapisochin G, Marquez M, Dondero F, Rayar M et al. Defining benchmarks in liver transplantation: a multicenter outcome analysis determining best achievable results. *Ann Surg* 2018;**267**:419–425
- Rössler F, Sapisochin G, Song G, Lin YH, Simpson MA, Hasegawa K et al. Defining benchmarks for major liver surgery: a multicenter analysis of 5202 living liver donors. *Ann Surg* 2016;**264**:492–500
- Schlegel A, van Reeve M, Croome K, Parente A, Dolcet A, Widmer J et al. A multicentre outcome analysis to define global benchmarks for donation after circulatory death liver transplantation. *J Hepatol* 2022;**76**:371–382
- Schmidt HM, Gisbertz SS, Moons J, Rouvelas I, Kauppi J, Brown A et al. Defining benchmarks for transthoracic esophagectomy: a multicenter analysis of total minimally invasive esophagectomy in low risk patients. *Ann Surg* 2017;**266**:814–821
- Staiger RD, Rössler F, Kim MJ, Brown C, Trenti L, Sasaki T et al. Benchmarks in colorectal surgery: multinational study to define quality thresholds in high and low anterior resection. *Br J Surg* 2022;**109**:1274–1281
- Gero D, Muller X, Staiger RD, Gutschow CA, Vonlanthen R, Bueter M et al. How to establish benchmarks for surgical outcomes? A checklist based on an international expert Delphi consensus. *Ann Surg* 2022;**275**:115–120
- Staiger RD, Schwandt H, Puhana MA, Clavien PA. Improving surgical outcomes through benchmarking. *Br J Surg* 2019;**106**:59–64
- Domenghino A, Walbert C, Birrer DL, Puhana MA, Clavien P-A, Heuskel D et al. Consensus recommendations on how to assess the quality of surgical interventions. *Nat Med* 2023;**29**:811–822
- Kim W, Kim H-H, Han S-U, Kim M-C, Hyung WJ, Ryu SW et al. Decreased morbidity of laparoscopic distal gastrectomy compared with open distal gastrectomy for stage I gastric cancer: short-term outcomes from a multicenter randomized controlled trial (KLASS-01). *Ann Surg* 2016;**263**:28–35
- Lee HJ, Hyung WJ, Yang HK, Han SU, Park YK, An JY et al. Short-term outcomes of a multicenter randomized controlled trial comparing laparoscopic distal gastrectomy with D2 lymphadenectomy to

- open distal gastrectomy for locally advanced gastric cancer (KLASS-02-RCT). *Ann Surg* 2019;**270**:983–991
30. Baiocchi GL, Giacomuzzi S, Marrelli D, Bencivenga M, Morgagni P, Rosa F et al. Complications after gastrectomy for cancer: Italian perspective. *Updates Surg* 2017;**69**:285–288
 31. Dindo D, Demartines N, Clavien PA. Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg* 2004;**240**:205–213
 32. Slankamenac K, Graf R, Barkun J, Puhani MA, Clavien PA. The comprehensive complication index: a novel continuous scale to measure surgical morbidity. *Ann Surg* 2013;**258**:1–7
 33. Lane-Fall MB, Neuman MD. Outcomes measures and risk adjustment. *Int Anesthesiol Clin* 2013;**51**:10–21
 34. Hemkens LG, Benchimol EI, Langan SM, Briel M, Kasenda B, Januel JM et al. The reporting of studies using routinely collected health data was often insufficient. *J Clin Epidemiol* 2016;**79**:104–111
 35. Langan SM, Benchimol EI. The need to improve reporting of routinely collected dermatology data for patient benefit. *Br J Dermatol* 2016;**174**:477–480
 36. Spray TL, Gaynor JW. A word of caution in public reporting. *Semin Thorac Cardiovasc Surg Pediatr Card Surg Annu* 2017;**20**:49–55
 37. Berlth F, Yang HK. Minimal-invasive gastrectomy: what the west can learn from the east? *Updates Surg* 2018;**70**:181–187
 38. Bollschweiler E, Berlth F, Baltin C, Mönig S, Hölscher AH. Treatment of early gastric cancer in the Western World. *World J Gastroenterol* 2014;**20**:5672–5678
 39. Hyung WJ, Yang H-K, Park Y-K, Lee H-J, An JY, Kim W et al. Long-term outcomes of laparoscopic distal gastrectomy for locally advanced gastric cancer: the KLASS-02-RCT randomized clinical trial. *J Clin Oncol* 2020;**38**:3304–3313
 40. Kim TH, Suh YS, Huh YJ, Son YG, Park JH, Yang JY et al. The comprehensive complication index (CCI) is a more sensitive complication index than the conventional Clavien–Dindo classification in radical gastric cancer surgery. *Gastric Cancer* 2018;**21**:171–181
 41. van der Veen A, Brenkman HJF, Seesing MFJ, Haverkamp L, Luyer MDP, Nieuwenhuijzen GAP et al. Laparoscopic versus open gastrectomy for gastric cancer (LOGICA): a multicenter randomized clinical trial. *J Clin Oncol* 2021;**39**:978–989
 42. van der Wielen N, Straatman J, Daams F, Rosati R, Parise P, Weitz J et al. Open versus minimally invasive total gastrectomy after neoadjuvant chemotherapy: results of a European randomized trial. *Gastric Cancer* 2021;**24**:258–271
 43. Hartgrink HH, van de Velde CJ, Putter H, Bonenkamp JJ, Klein Kranenbarg E, Songun I et al. Extended lymph node dissection for gastric cancer: who may benefit? Final results of the randomized Dutch gastric cancer group trial. *J Clin Oncol* 2004;**22**:2069–2077
 44. Katai H, Mizusawa J, Katayama H, Takagi M, Yoshikawa T, Fukagawa T et al. Short-term surgical outcomes from a phase III study of laparoscopy-assisted versus open distal gastrectomy with nodal dissection for clinical stage IA/IB gastric cancer: Japan Clinical Oncology Group Study JCOG0912. *Gastric Cancer* 2017;**20**:699–708
 45. Inaki N, Etoh T, Ohyama T, Uchiyama K, Katada N, Koeda K et al. A multi-institutional, prospective, phase II feasibility study of laparoscopy-assisted distal gastrectomy with D2 lymph node dissection for locally advanced gastric cancer (JLSSG0901). *World J Surg* 2015;**39**:2734–2741
 46. Trapani R, Rausei S, Reddavid R, Degiuli M. Risk factors for esophago-jejunal anastomosis leakage after total gastrectomy for cancer. A multicenter retrospective study of the Italian research group for gastric cancer. *Eur J Surg Oncol* 2020;**46**:2243–2247
 47. Yang HK, Hyung WJ, Han SU, Lee YJ, Park JM, Cho GS et al. Comparison of surgical outcomes among different methods of esophagojejunostomy in laparoscopic total gastrectomy for clinical stage I proximal gastric cancer: results of a single-arm multicenter phase II clinical trial in Korea, KLASS 03. *Surg Endosc* 2021;**35**:1156–1163
 48. Kurokawa Y, Doki Y, Mizusawa J, Terashima M, Katai H, Yoshikawa T et al. Bursectomy versus omentectomy alone for resectable gastric cancer (JCOG1001): a phase 3, open-label, randomised controlled trial. *Lancet Gastroenterol Hepatol* 2018;**3**:460–468
 49. Katai H, Mizusawa J, Katayama H, Morita S, Yamada T, Bando E et al. Survival outcomes after laparoscopy-assisted distal gastrectomy versus open distal gastrectomy with nodal dissection for clinical stage IA or IB gastric cancer (JCOG0912): a multicentre, non-inferiority, phase 3 randomised controlled trial. *Lancet Gastroenterol Hepatol* 2020;**5**:142–151
 50. Park DJ, Kim YW, Yang HK, Ryu KW, Han SU, Kim HH et al. Short-term outcomes of a multicentre randomized clinical trial comparing laparoscopic pylorus-preserving gastrectomy with laparoscopic distal gastrectomy for gastric cancer (the KLASS-04 trial). *Br J Surg* 2021;**108**:1043–1049
 51. Sano T, Sasako M, Mizusawa J, Yamamoto S, Katai H, Yoshikawa T et al. Randomized controlled trial to evaluate splenectomy in total gastrectomy for proximal gastric carcinoma. *Ann Surg* 2017;**265**:277–283
 52. Sasako M, Sano T, Yamamoto S, Kurokawa Y, Nashimoto A, Kurita A et al. D2 lymphadenectomy alone or with para-aortic nodal dissection for gastric cancer. *N Engl J Med* 2008;**359**:453–462
 53. Bonnot PE, Piessen G, Kepenekian V, Decullier E, Pocard M, Meunier B et al. Cytoreductive surgery with or without hyperthermic intraperitoneal chemotherapy for gastric cancer with peritoneal metastases (CYTO-CHIP study): a propensity score analysis. *J Clin Oncol* 2019;**37**:2028–2040