

ORIGINAL ARTICLE

Association between type and location of germline *BRCA1/2* pathogenic or likely pathogenic variants with phenotype and prognosis in young patients with breast cancer: results from an international cohort study

A. Toss^{1,2†}, E. Blondeaux^{3†}, E. Tenedini², L. Bonamici², R. Graffeo⁴, L. Livraghi⁵, C. Villarreal-Garza⁶, R. Bernstein Molho⁷, A. Kwong^{8,9,10}, J. Balmana¹¹, H. Wildiers¹², E. Agostinetto¹³, K. A. Phillips^{14,15}, K. Pogoda¹⁶, T. Renaud¹⁷, C. Rousset-Jablonski^{18,19,20}, A. Ferrari^{21,22}, H. C. F. Moore²³, F. A. Peccatori²⁴, S. Paluch-Shimon²⁵, R. Fruscio^{26,27}, C. Micheri²⁸, S. M. Wong²⁹, W. Cui^{14,15}, C. Vernieri^{30,31}, M. K. Lee³², L. De Marchis^{33,34}, F. J. Couch³⁵, L. Del Mastro^{36,37}, F. Puglisi^{38,39}, P. A. Meireles⁴⁰, Z. Kemp⁴¹, A. Matikas⁴², J. Plichta⁴³, M. Del Pilar Estevez-Diz⁴⁴, A. Di Meglio⁴⁵, N. Cichowska-Cwalinska⁴⁶, C. Gianni⁴⁷, R. Yerushalmi^{48,49}, R. Sanchez-Bayona⁵⁰, B. Mrinakova⁵¹, L. Matos⁵², G. Bianchini⁵³, M. Caleffi⁵⁴, A. Krivokuca⁵⁵, Y. Abdou⁵⁶, M. Mariño-Mariño⁵⁷, A. Parokonnaya⁵⁸, M. Okano^{59,60}, N. Antone⁶¹, C. Saavedra⁶², A. Sonnenblick⁶³, R. Duchnowska⁶⁴, H. L. Pais⁶⁵, N. Harbeck⁶⁶, L. Cortesi^{2,67}, V. Delucchi³, C. De Angelis^{68,69†} & M. Lambertini^{36,37*†}

¹Department of Oncology and Haematology, Azienda Ospedaliero-Universitaria di Modena, Modena; ²Department of Medical and Surgical Sciences, University of Modena and Reggio Emilia, Modena; ³Clinical Trial Unit, U.O. Epidemiologia Clinica, IRCCS Ospedale Policlinico San Martino, Genova, Italy; ⁴Oncology Institute of Southern Switzerland, EOC, Bellinzona, Switzerland; ⁵Nuovo Ospedale di Prato—Santo Stefano, Prato, Italy; ⁶Breast Cancer Center, Hospital Zambrano Hellion TecSalud, Tecnológico de Monterrey, Monterrey, Mexico; ⁷Susanne Levy Gertner Oncogenetics Unit, The Danek Gertner Institute of Human Genetics, Chaim Sheba Medical Center, affiliated to Tel Aviv University, Tel Hashomer, Israel; ⁸Hong Kong Hereditary Breast Cancer Family Registry, Hong Kong SAR; ⁹Cancer Genetics Centre and Breast Surgery Centre, Hong Kong Sanatorium and Hospital, Hong Kong SAR; ¹⁰Department of Surgery, The University of Hong Kong, Hong Kong SAR; ¹¹Hereditary Cancer Genetics Unit, Medical Oncology Department, Vall d'Hebron University Hospital, Vall d'Hebron Institute of Oncology (VHIO), Barcelona, Spain; ¹²Department of General Medical Oncology, University Hospitals Leuven, KU Leuven, Leuven; ¹³Université libre de Bruxelles (ULB), Hôpital Universitaire de Bruxelles (H.U.B.), Institut Jules Bordet, Service de Oncologie, Brussels, Belgium; ¹⁴Department of Medical Oncology, Peter MacCallum Cancer Centre, Melbourne; ¹⁵Sir Peter MacCallum Department of Oncology, The University of Melbourne, Melbourne, Australia; ¹⁶Department of Breast Cancer and Reconstructive Surgery, Maria Skłodowska-Curie National Research Institute of Oncology, Warsaw, Poland; ¹⁷Cancer Genetics Unit, Bergonie Institute, Bordeaux; ¹⁸Department of Surgery, Leon Berard Cancer Center, Lyon; ¹⁹Hôpital Femme Mère Enfant, Hospices Civils de Lyon, Lyon; ²⁰INSERM U1290 RESHAPE, Lyon, France; ²¹Hereditary Breast and Ovarian Cancer (HBOC) Unit and General Surgery 3—Senology, Breast Cancer Center, Fondazione IRCCS Policlinico San Matteo, Pavia; ²²University of Pavia, Pavia, Italy; ²³Department of Medical Oncology, Cleveland Clinic Taussig Cancer Institute, Cleveland, USA; ²⁴Gynecologic Oncology Department, European Institute of Oncology IRCCS, Milan, Italy; ²⁵Sharett Institute of Oncology, Hadassah University Hospital & Faculty of Medicine, Hebrew University, Jerusalem, Israel; ²⁶UO Gynecology, Fondazione IRCCS San Gerardo dei Tintori, Monza; ²⁷Department of Medicine and Surgery, University of Milan-Bicocca, Milan, Italy; ²⁸SUMA (Grupo Cooperativo Argentino para el Estudio y la Investigación del Cáncer de Mama), Buenos Aires, Argentina; ²⁹Stroll Cancer Prevention Centre, Jewish General Hospital, and McGill University Medical School, Montreal, Canada; ³⁰Breast Unit, Medical Oncology Department, Fondazione IRCCS Istituto Nazionale dei Tumori, Milan; ³¹Department of Oncology and Hematology-Oncology, University of Milan, Milan, Italy; ³²Memorial Sloan Kettering Cancer Center MSKCC, New York, USA; ³³Division of Medical Oncology, Department of Radiological, Oncological and Pathological Sciences, "La Sapienza" University of Rome, Rome; ³⁴Oncology Unit, Department of Hematology Oncology and Dermatology, Umberto 1 Policlinico di Roma, Rome, Italy; ³⁵Department of Laboratory Medicine and Pathology, Mayo Clinic College of Medicine, Rochester, USA; ³⁶Department of Medical Oncology, U.O. Clinica di Oncologia Medica, IRCCS Ospedale Policlinico San Martino, Genova; ³⁷Department of Internal Medicine and Medical Specialties (DIMI), School of Medicine, University of Genova, Genova; ³⁸Department of Medical Oncology, Centro di Riferimento Oncologico di Aviano (CRO) IRCCS, Aviano; ³⁹Department of Medicine, University of Udine, Udine, Italy; ⁴⁰Medical Oncology Department, Instituto Português de Oncologia de Lisboa Francisco Gentil, Lisbon, Portugal; ⁴¹Breast Medical Oncology and Cancer Genetics Units, Royal Marsden Hospital, London, UK; ⁴²Department of Oncology/Pathology, Karolinska Institute and Breast Center, Karolinska University Hospital, Stockholm, Sweden; ⁴³Department of Surgery, Duke University, Durham, USA; ⁴⁴Department of Medical Oncology, Instituto do Cancer do Estado de São Paulo, Faculdade de Medicina da Universidade de São Paulo, São Paulo, Brazil; ⁴⁵Université Paris-Saclay, Gustave Roussy, Inserm, Prédicteurs moléculaires et nouvelles cibles en oncologie, F-94805, Villejuif, France; ⁴⁶Medical University of Gdansk, Gdansk, Poland; ⁴⁷Medical Oncology, Breast & GYN Unit, IRCCS Istituto Romagnolo per lo Studio dei Tumori (IRST) "Dino Amadori", Meldola, Italy; ⁴⁸Davidoff Cancer Center, Rabin Medical Center—Beilinson, Petah Tikva; ⁴⁹Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel; ⁵⁰Hospital Universitario 12 de Octubre, Madrid, Spain; ⁵¹Department of Oncology, Faculty of Medicine, Comenius University and St. Elisabeth Cancer Institute; ⁵²Breast Unit, Champalimaud Clinical Center, Champalimaud Foundation, Lisbon, Portugal; ⁵³IRCCS Ospedale San Raffaele, Milan, Italy; ⁵⁴Nucleo Mama, Hospital Moinhos de Vento, Porto Alegre, Brazil; ⁵⁵Institute for Oncology and Radiology of Serbia, Belgrade, Serbia; ⁵⁶University of North Carolina Lineberger Comprehensive Cancer Center, Chapel Hill, USA; ⁵⁷Hospital Universitario Son Espases, Palma, Spain; ⁵⁸Blokhin National Medical Research Center of Oncology, Moscow, Russia; ⁵⁹Hoshi General Hospital, Koriyama; ⁶⁰Fukushima Medical University, Fukushima, Japan; ⁶¹Institute of Oncology "Ion Chiricuta", Cluj-Napoca, Romania; ⁶²Ramon y Cajal, IOB Institute of Oncology (Madrid), International Breast Cancer Center (IBCC), Madrid, Spain; ⁶³The Oncology Division, Tel Aviv Sourasky Medical Center, Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel; ⁶⁴Department of Oncology, Military Institute of Medicine—National Research Institute, Warsaw, Poland; ⁶⁵Hospital de Santa Maria—University of Lisbon, Lisbon, Portugal; ⁶⁶Breast Center, Department of Obstetrics and Gynecology and CCC Munich, LMU University Hospital, Munich, Germany; ⁶⁷AUSL-IRCCS Reggio Emilia, Reggio Emilia; ⁶⁸Clinical and Translational Oncology, Scuola Superiore Meridionale, Naples; ⁶⁹Istituto Nazionale Tumori Napoli IRCCS "Fondazione Pascale", Naples, Italy


*Correspondence to: Prof. Matteo Lambertini, Medical Oncology Department, U.O.C. Clinica di Oncologia Medica, IRCCS Policlinico San Martino Hospital—University of Genova, Largo Rosanna Benzi 10, 16132 Genova, Italy. Tel: +39-010-555-4254; Fax: +39-010-555-6536
E-mail: matteo.lambertini@unige.it (M. Lambertini).

†Co-first authors.

‡Co-last authors.

0923-7534/© 2025 The Author(s). Published by Elsevier Ltd on behalf of European Society for Medical Oncology. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

GRAPHICAL ABSTRACT



Available online 17 November 2025

Background: The clinical implications of specific pathogenic and likely pathogenic variant (LP/PV) types and locations in the *BRCA1* or *BRCA2* tumor-suppressor genes remain to be elucidated.

Patients and methods: The BRCA BCY Collaboration (NCT03673306) is an international, multicenter, hospital-based, retrospective cohort study that included *BRCA* carriers diagnosed with invasive breast cancer at the age of ≤ 40 years between January 2000 and December 2020. In this analysis, only patients with detailed available information on LP/PVs in the *BRCA* genes were included. Clinicopathological features and survival outcomes [disease-free survival (DFS) and overall survival (OS)] were investigated according to LP/PV type [insertion-deletion (indel) versus single-nucleotide variants versus copy number variations; truncating versus non-truncating LP/PVs; frameshift versus nonsense versus splicing versus missense LP/PVs] and location (exon involved and protein domain).

Results: Out of 5660 patients from 109 centers worldwide, 3294 were eligible for the present analysis (2080 *BRCA1* and 1214 *BRCA2*). The distribution of LP/PV types showed no meaningful associations with baseline clinicopathological features. *BRCA1* protein-truncating variants were associated with worse OS compared with non-truncating variants [hazard ratio (HR) 2.00, 95% confidence interval (CI) 1.17-3.41]. A similar, though non-significant, trend was observed for *BRCA2*. Missense variants were linked to better OS for both *BRCA1* (HR 0.48, 95% CI 0.28-0.84) and *BRCA2* carriers (HR 0.17, CI 0.03-0.96). Regarding variant location, *BRCA1* LP/PVs outside exons 2, 10, and 19 were associated with improved OS. In *BRCA2*, LP/PVs located in exons 15-26 and other regions were linked to worse DFS compared with those in exon 10, with no significant differences in OS.

Conclusions: This study advances our understanding of the influence of specific types of *BRCA* LP/PVs on breast cancer characteristics and outcomes. A deeper understanding of these variant-specific features will drive future research and support the development of tailored clinical strategies based on individual *BRCA* variant.

Key words: hereditary breast cancer, *BRCA1*, *BRCA2*, variant type, protein-truncating variant, overall survival

INTRODUCTION

BRCA1 and *BRCA2* are tumor-suppressor genes encoding for multidomain proteins, involved in the error-free repair of DNA double-strand breaks (DSBs) through homologous recombination.¹ *BRCA1* also plays a role in cell-cycle regulation, as well as participating in non-homologous end-joining and single-strand annealing.¹

The *BRCA1* gene encompasses 24 exons, 22 of which encode a protein with multiple domains including a RING finger at its N-terminal region, two nuclear localization signals (NLS), a coiled-coil domain, and two *BRCA1* C-terminal (BRCT) domains.² The *BRCA2* gene consists of 27 exons coding for a protein with four main regions: the N-terminal fragment, the central region containing eight BRC

repeats that recruit RAD51 at DSB sites (RAD51-binding domain), the DNA-binding domain (DBD) located in the C-terminal region with an alpha helical domain followed by three oligonucleotide/oligosaccharide-binding (OB) folds, and the NLS located at the extreme C-terminal region.^{3,4}

To date, thousands of sequence variants in the *BRCA* genes have been identified, with only a few subsets recognized as likely pathogenic or pathogenic variants (LP/PVs). The distribution of these LP/PVs varies greatly in frequency and types among different populations.⁵ Most known *BRCA1* and *BRCA2* LP/PVs are small insertions or deletions (indels), followed by single-nucleotide variants (SNVs). The most frequent consequence of indels is a disruption of the reading frame, namely frameshift variants, usually resulting in premature stop codons. The consequent transcription termination can trigger nonsense-mediated messenger RNA decay or produce truncated, non-functional proteins.⁶ Conversely, the substitution of one nucleotide for another can result in the replacement of one amino acid with a different one, known as missense variants, or the introduction of stop codons, known as nonsense variants.⁷ The pathogenicity of less common alterations, such as splicing variants, can impact structure, expression, or function of the encoded protein.⁸ Finally, variations in the number of copies of a particular DNA segment [copy number variations (CNVs)] are rare, accounting for ~5% of all LP/PVs and being more prevalent in *BRCA1* versus *BRCA2*.⁹

LP/PVs in the *BRCA1* or *BRCA2* genes significantly increase the risk of developing breast cancer, ovarian cancer, and other malignancies.¹⁰ Particularly, clinicopathological features of breast cancer may be influenced by the specific gene involved, with *BRCA1* more frequently associated with triple-negative breast cancer and *BRCA2* with hormone receptor-positive tumors.¹¹⁻¹³ Evidence also suggests that the clinical implications of these LP/PVs may vary depending on the specific variant type or location within *BRCA1* and *BRCA2* genes. Previous studies showed that LP/PVs occurring in distinct regions of *BRCA1* or *BRCA2* genes confer varying levels of risk of developing breast or ovarian cancer.^{14,15} Moreover, variant types that result in partially functional *BRCA1* or *BRCA2* proteins might confer reduced cancer risk and lower tumor aggressiveness compared with LP/PVs leading to transcript decay or protein truncation.¹⁶⁻¹⁸ Additionally, specific *BRCA1* or *BRCA2* variant types have been shown to influence treatment responses in patients with ovarian cancer, especially to platinum-based chemotherapy and poly (ADP-ribose) polymerase (PARP) inhibitors.^{19,20} However, no evidence exists in this regard for patients with breast cancer.

Identifying associations between the type of genetic variant and specific breast cancer characteristics, such as age at onset or tumor aggressiveness, may contribute to optimizing preventive strategies, including the intensification of surveillance protocols or the anticipation of risk-reducing surgical interventions. Furthermore, determining the prognostic significance of variant type or location could aid in refining therapeutic approaches, guiding decisions on treatment escalation or de-escalation. On these grounds,

the present study aimed to investigate the association between specific type and location of *BRCA1* or *BRCA2* LP/PVs with clinicopathological features and survival outcome in a global population of *BRCA* carriers diagnosed with breast cancer at age ≤ 40 years.

PATIENTS AND METHODS

Study design and participants

This is an international, multicenter, hospital-based, retrospective cohort study including patients diagnosed with invasive breast cancer between January 2000 and December 2020, who were ≤ 40 years and known to harbor germline LP/PVs in *BRCA1* and/or *BRCA2* genes.^{21,22} Main exclusion criteria included a history of non-invasive breast cancer, diagnosis of ovarian cancer or other malignancies with no history of invasive breast cancer, or variants of uncertain clinical significance (VUS). For this specific analysis, patients for whom the specific variant was not provided and patients with stage IV *de novo* disease, unclassifiable variants, unknown uptake or timing of risk-reducing surgeries, multiple *BRCA1* and *BRCA2* LP/PVs, and previous LP/PVs reclassified as VUS were also excluded.

Clinicopathological characteristics of patients were collected, including country of enrollment, year at breast cancer diagnosis, age at diagnosis, body mass index, timing of genetic testing, *BRCA* LP/PVs, tumor histology, grade, stage at diagnosis, hormone receptor status, human epidermal growth factor receptor 2 status, type of breast and axillary surgery, risk-reducing surgeries, systemic anti-cancer treatments, and survival outcomes.

The study was conducted in accordance with the Declaration of Helsinki. The Institut Jules Bordet (Brussels, Belgium) sponsored the study and acted as the central ethics committee. The study also obtained ethical approval from local, regional, or national institutional review boards/ethics committees of the participating centers if requested by local regulations. The first (AT and EB) and last (CDA and ML) authors guarantee for the accuracy and completeness of the data and analyses.

The STrengthening the Reporting of OBservational studies in Epidemiology (STROBE) statement was followed to report this work.²³

The study is registered at [ClinicalTrials.gov](https://clinicaltrials.gov) (NCT03673306).

Variant classification

BRCA LP/PVs were categorized as SNVs when a single-nucleotide change occurred, indels when one or more bases were lost or duplicated, and CNVs when one-copy gains and losses of single or multiple exons occurred in the coding sequence.

The nomenclature of LP/PVs was harmonized to the standards of the International Human Genome Variation Society (HGVS) for both SNVs and indels.^{24,25} CNVs were reported with no exact genomic breakpoints allowing for comparable data across centers and over time. The annotation and

classification of LP/PVs were verified by consulting supporting evidence from the literature, public databases (ENIGMA, <https://enigmaconsortium.org/>; BRCA Exchange, <https://brcaexchange.org>; ClinVar, <https://www.ncbi.nlm.nih.gov/clinvar/>; and LOVD, <https://www.lovd.nl/>), and annotation tools such as VarSome²⁶ and Franklin by Genoox (<https://franklin.genoox.com>). LP/PVs were further categorized according to ClinGene Curation Resource²⁷ and the American College of Medical Genetics and Genomics (ACMG) classification.²⁸

Genetic information for LP/PVs was compiled, including coding consequence, exon position, affected protein domain, and whether the genetic defect was predicted to truncate the coding sequence of the gene, resulting in a shorter version of the protein (protein truncation).

The coding consequences were summarized as nonsense and missense variants (indicating a stop codon or a different codon), inframe deletions (deletion of triplets preserving the gene's reading frame), frameshifts (shifting and disrupting the reading frame), and splicing (if variants are predicted to disrupt the RNA splicing).

LP/PVs were defined as protein-truncating variants if SNVs or indels were predicted to introduce premature stop codons (nonsense variants) or to disrupt the reading frame of the transcript (frameshift variants). When protein truncation could not be reliably predicted, the variant classification was considered not classifiable, indicating that the data were not evaluable rather than missing.

The affected exon was determined by the position of each variant in the coding sequence, according to the transcripts from the Matched Annotation from the NCBI and EMBL-EBI (MANE) collaboration²⁹: NM_007294.4 for *BRCA1* and NM_000059.4 for *BRCA2*. The current exon 10 in the MANE transcript of *BRCA1* is often reported as exon 11 in prior literature. This miscalculation happened because over time different transcript models and different databases were used, employing different counting schemes for the 5' untranslated exons (1a, 1b, and sometimes 2) and the tiny exon 4 (21 bp long). Moreover, the *BRCA1* transcript was sometimes numbered starting from exon 2 because exon 1 is non-coding and does not contribute to the protein sequence. However, in the current MANE reference transcript (NM_007294.4), which is the reviewed major transcript, *BRCA1* is described as having 24 exons, named exon 1 to exon 24, with numbering beginning at the first exon regardless of whether it codes for protein. The RefSeq/MANE Select transcript NM_007294.4 is the current standardized reference used for exon numbering and variant annotation according to the European Molecular Genetics Quality Network (EMQN) and HGVS guidelines.^{30,31}

Protein-level positions were inferred based on the DNA-level information. *BRCA1* and *BRCA2* protein domains were considered if well characterized in the literature.^{2,4,32,33} In the absence of a protein domain associated to the position of the variant, 'unknown' was assigned, whereas in the presence of more than one domain involved, 'multiple domains' or 'whole protein' was used.

Objectives

The aim of the present study was to investigate the role of specific type and location of *BRCA* LP/PV on the clinical features and outcomes of breast cancer in young *BRCA* carriers. Clinical features and survival outcomes were separately investigated for patients with *BRCA1* and *BRCA2* LP/PVs, according to variant category (indels versus SNVs versus CNVs), protein truncation (protein-truncating versus protein non-truncating versus unclassified variants), coding consequence (frameshift versus nonsense versus splicing versus missense versus inframe deletion variants), and LP/PV location (exon and protein domain involved).

Statistical analysis

Descriptive statistics were used to summarize continuous and categorical variables. Continuous variables were reported as median and interquartile range (IQR), while categorical variables as frequencies and percentages. The associations between LP/PV types and clinicopathological features were analyzed using chi-square test or Kruskal–Wallis–Wilcoxon rank-sum test, for categorical or continuous variables as appropriate. The median follow-up in the entire cohort and according to the LP/PV type was computed using the reverse Kaplan–Meier method.

To investigate the association between LP/PV types and survival outcomes, disease-free survival (DFS) and overall survival (OS) were considered as endpoints. DFS was defined as the time from diagnosis until locoregional recurrence, distant metastases, new contralateral or ipsilateral breast cancer, second primary malignancy, or death from any cause. OS was defined as the time from diagnosis until death from any cause.³⁴ DFS and OS event rates were computed as the ratio between the total number of events and the total observation time. Unadjusted and adjusted Cox models were used to estimate the association between each specific type and location of *BRCA* LP/PV and survival outcomes. Adjusted models included country and year at breast cancer diagnosis as stratification factors and the uptake of risk-reducing surgeries (risk-reducing mastectomy and risk-reducing salpingo-oophorectomy) as time-dependent covariates. Furthermore, to account for potential immortal time bias (i.e. patients had to survive to undergo *BRCA* testing), both unadjusted and adjusted survival models where observation times were left truncated at the time of *BRCA* testing were carried out. Considering this potential bias, results of the adjusted Cox model with left truncation were reported as the primary analysis in the manuscript, and the Kaplan–Meier method, accounting for left truncation, was utilized to plot the survival curves. In the figures, a *P* value <0.05 indicates that at least one comparison among the groups is statistically significant.

For the comparison of the location of variants in different exons, for *BRCA1*, the three exons with the highest frequency of LP/PVs (exon 2, exon 10, and exon 19) were compared with each other and other exons. Specifically, exon 2 contributes to the coding of the RING domain,

exon 10 to the NLS and the coiled-coil domains, and exon 19 to the BRCT domains. For *BRCA2*, the two exons with the highest frequency of LP/PVs (exon 10 and exon 11) and exons 15-26 were considered for the comparative analysis. In this gene, exons 10 and 11 encode the BRC repeats, whereas exons 15-26 encode the DBD.

For the comparison of different protein domains, only patients with known domain and protein non-truncating LP/PVs were included. For the comparison of different exons and type of LP/PVs in terms of frameshift versus nonsense versus splicing versus missense versus inframe deletion variants, patients with CNVs were excluded.

For the comparison of frameshift versus inframe deletion versus nonsense versus splicing versus missense LP/PVs, and for the comparison of exons, Cox models were repeated by shifting the reference group in order to identify any significant comparison.

Considering the explorative nature of the analysis, no adjustment for multiple tests was carried out. All reported *P* values are two-sided, and *P* < 0.05 was considered statistically significant. All statistical analyses were carried out using SAS software, version 9.4 (SAS Institute Inc., Cary,

NC), and R software, version 4.3.3 (R Foundation for Statistical Computing, Vienna, Austria).

RESULTS

Patient population

From a cohort of 5660 patients enrolled across 109 centers globally in the primary study, 3294 patients were included in the present analysis (Supplementary Figure S1, available at <https://doi.org/10.1016/j.annonc.2025.11.004>). Among them, 2080 harbored a *BRCA1* LP/PV, while 1214 harbored a *BRCA2* LP/PV. The distribution of LP/PV type and location within the *BRCA1* and *BRCA2* genes is detailed in Table 1. The baseline characteristics of patients included in the present study are summarized in Table 2. Overall, the median follow-up was 7.9 (IQR 4.5-12.7) years for *BRCA1* carriers and 7.9 (IQR 4.5-13.1) years for *BRCA2* carriers.

The most common LP/PVs (observed in at least 10 patients), along with their prevalence and geographic distribution, are included in Supplementary Tables S1 and S2, available at <https://doi.org/10.1016/j.annonc.2025.11.004>.

Variant type (indel, SNV, and CNV)

In *BRCA1* carriers, the distribution of variant types differed by geographic region; no other meaningful association between baseline clinicopathological variables and LP/PV types was observed (Supplementary Table S3, available at <https://doi.org/10.1016/j.annonc.2025.11.004>). The rate and type of first DFS events and the rate of OS events for *BRCA1* patients are detailed in Supplementary Table S4, available at <https://doi.org/10.1016/j.annonc.2025.11.004>. No difference in DFS was found according to the type of LP/PV (Figure 1A and Supplementary Table S5, available at <https://doi.org/10.1016/j.annonc.2025.11.004>). Accordingly, no significant difference in OS was observed among SNV, CNV, and indel variants (Figure 1B and Supplementary Table S6, available at <https://doi.org/10.1016/j.annonc.2025.11.004>).

In *BRCA2* carriers, the distribution of variant types differed by geographic region; no other meaningful association between baseline clinicopathological variables and LP/PV types was observed (Supplementary Table S7, available at <https://doi.org/10.1016/j.annonc.2025.11.004>). The rate and type of first DFS events and the rate of OS events for *BRCA2* patients are detailed in Supplementary Table S8, available at <https://doi.org/10.1016/j.annonc.2025.11.004>. No difference in DFS (Figure 1C) or in OS (Figure 1D and Supplementary Tables S9 and S10, available at <https://doi.org/10.1016/j.annonc.2025.11.004>) was found according to the type of LP/PV.

Protein truncation (protein-truncating variants and non-truncating variants)

In *BRCA1* carriers, the distribution of protein-truncating variants differed according to geographic region. No other meaningful associations between baseline clinicopathological variables and protein truncation were detected (Supplementary Table S11, available at <https://doi.org/10.1016/j.annonc.2025.11.004>). The rate and type of first DFS

Table 1. Distribution of LP/PV type and location in *BRCA1* and *BRCA2* genes

	<i>BRCA1</i> <i>n</i> = 2080 (%)	<i>BRCA2</i> <i>n</i> = 1214 (%)	Total <i>n</i> = 3294 (%)
Variant type			
Indel	1248 (60.0)	772 (63.6)	2020 (61.3)
SNV	667 (32.1)	409 (33.7)	1076 (32.7)
CNV	165 (7.9)	33 (2.7)	198 (6.0)
	(143 ODel- 22 OCDup)	(32 ODel- 1 OCDup)	(175 ODel- 23 OCDup)
Protein truncation			
Yes	1514 (72.8)	1007 (83.0)	2521 (76.5)
No	227 (10.9)	50 (4.1)	277 (8.4)
NC	339 (16.3)	157 (12.9)	496 (15.1)
Coding consequence ^a	<i>n</i> = 1915 (%)	<i>n</i> = 1181 (%)	<i>n</i> = 3096 (%)
Frameshift	1158 (60.5)	710 (60.1)	1868 (60.3)
Inframe deletion	14 (0.7)	2 (0.2)	16 (0.5)
Missense	213 (11.1)	48 (4.1)	261 (8.4)
Nonsense	356 (18.6)	299 (25.3)	655 (21.2)
Splicing	174 (9.1)	122 (10.3)	296 (9.6)
Exon ^a	<i>n</i> = 1915 (%)	<i>n</i> = 1181 (%)	<i>n</i> = 3096 (%)
Exon 2	243 (12.7)	—	243 (7.8)
Exon 10	623 (32.5)	—	623 (20.1)
Exon 19	346 (18.1)	—	346 (11.2)
Other exons	703 (36.7)	—	703 (22.7)
Exon 10	—	107 (9.1)	107 (3.5)
Exon 11	—	564 (47.8)	564 (18.2)
Exon 15-26	—	312 (26.4)	312 (10.1)
Other exons	—	198 (16.8)	198 (6.4)
Protein domain ^b	<i>n</i> = 227 (%)	<i>n</i> = 50 (%)	<i>n</i> = 277 (%)
RING domain	117 (51.5)	—	117 (42.2)
BRCT domain	108 (47.6)	—	108 (39.0)
OB domain	—	44 (88.0)	44 (15.9)
BRC domain	—	2 (4.0)	2 (0.7)
Alpha helix	—	1 (2.0)	1 (0.4)
Unknown	2 (0.9)	3 (6.0)	5 (1.8)

BRCT, *BRCA1* C-terminal; CNV, copy number variations; indel, small deletions or insertions; NC, not classifiable; OB, oligonucleotide/oligosaccharide-binding; ODel, one-copy deletion; OCDup, one-copy duplication; SNV, single-nucleotide variants.

^aOnly in patients without CNV.

^bOnly in patients with no protein truncation.

Table 2. Baseline characteristics of patients

	BRCA1 n = 2080 (%)	BRCA2 n = 1214 (%)	Total n = 3294 (%)
Patients' characteristics			
Region			
Central/South America	153 (7.4)	70 (5.8)	223 (6.8)
Australia/Oceania	105 (5.1)	74 (6.1)	179 (5.4)
Northern Europe	269 (12.9)	179 (14.7)	448 (13.6)
Eastern Europe	189 (9.1)	59 (4.9)	248 (7.5)
North America	255 (12.3)	131 (10.8)	386 (11.7)
Southern Europe	781 (37.5)	534 (44.0)	1315 (39.9)
Asia	325 (15.6)	164 (13.5)	489 (14.9)
Africa	3 (0.1)	3 (0.2)	6 (0.2)
Year of diagnosis			
2000-2004	269 (12.9)	173 (14.3)	442 (13.4)
2005-2008	321 (15.4)	180 (14.8)	501 (15.2)
2009-2012	419 (20.1)	222 (18.3)	641 (19.5)
2013-2016	477 (22.9)	283 (23.3)	760 (23.1)
2017-2020	594 (28.6)	356 (29.3)	950 (28.8)
Median age (years) at diagnosis (IQR)	35.0 (31.0-37.0)	35.0 (32.0-38.0)	35.0 (31.0-38.0)
Age at diagnosis, years			
≤30	478 (23.0)	190 (15.7)	668 (20.3)
31-35	716 (34.4)	446 (36.7)	1162 (35.3)
36-40	886 (42.6)	578 (47.6)	1464 (44.4)
Median BMI at diagnosis (IQR)	23.0 (20.6-26.2)	22.8 (20.3-25.9)	23.0 (20.4-26.0)
BMI			
Underweight	108 (5.2)	71 (5.8)	179 (5.4)
Normal weight	1059 (50.9)	658 (54.2)	1717 (52.1)
Overweight	423 (20.3)	216 (17.8)	639 (19.4)
Obese	201 (9.7)	105 (8.7)	306 (9.3)
Unknown	289 (13.9)	164 (13.5)	453 (13.8)
Median time (years) to BRCA test (IQR)	0.4 (0.1-2.2)	0.7 (0.1-3.3)	0.5 (0.1-2.6)
Timing of BRCA testing^a			
Test before diagnosis	227 (10.9)	90 (7.4)	317 (9.6)
Test at diagnosis	851 (40.9)	446 (36.7)	1297 (39.4)
Test after diagnosis	919 (44.2)	638 (52.6)	1557 (47.3)
Unknown date of BRCA testing	83 (4.0)	40 (3.3)	123 (3.7)
Tumor characteristics			
Tumor histology			
Ductal	1775 (85.3)	988 (81.4)	2763 (83.9)
Other	286 (13.8)	217 (17.9)	503 (15.3)
Unknown	19 (0.9)	9 (0.7)	28 (0.9)
Tumor grade			
G1	21 (1.0)	45 (3.7)	66 (2.0)
G2	304 (14.6)	476 (39.2)	780 (23.7)
G3	1634 (78.6)	628 (51.7)	2262 (68.7)
Unknown	121 (5.8)	65 (5.4)	186 (5.6)
Tumor size			
T0	3 (0.1)	0 (0.0)	3 (0.1)
T1	741 (35.6)	470 (38.7)	1211 (36.8)
T2	988 (47.5)	526 (43.3)	1514 (46.0)
T3/T4	271 (13.0)	169 (13.9)	440 (13.4)
Unknown	77 (3.7)	49 (4.0)	126 (3.8)
Nodal status			
N0	1173 (56.4)	478 (39.4)	1651 (50.1)
N1	647 (31.1)	472 (38.9)	1119 (34.0)
N2/N3	207 (10.0)	217 (17.9)	424 (12.9)
Unknown	53 (2.5)	47 (3.9)	100 (3.0)
Hormone receptor status			
Negative	1534 (73.8)	189 (15.6)	1723 (52.3)
Positive	536 (25.8)	1021 (84.1)	1557 (47.3)
Unknown	10 (0.5)	4 (0.3)	14 (0.4)
HER2 status			
Negative	1907 (91.7)	1031 (84.9)	2938 (89.2)
Positive	100 (4.8)	136 (11.2)	236 (7.2)
Unknown	73 (3.5)	47 (3.9)	120 (3.6)

Continued

Table 2. Continued

	BRCA1 n = 2080 (%)	BRCA2 n = 1214 (%)	Total n = 3294 (%)
Treatment			
Breast surgery			
None	6 (0.3)	5 (0.4)	11 (0.3)
Breast conserving	864 (41.5)	354 (29.2)	1218 (37.0)
Mastectomy	1201 (57.7)	850 (70.0)	2051 (62.3)
Unknown	9 (0.4)	5 (0.4)	14 (0.4)
Axillary surgery			
No axillary surgery	46 (2.2)	39 (3.2)	85 (2.6)
Sentinel node biopsy only	971 (46.7)	420 (34.6)	1391 (42.2)
Axillary dissection	1031 (49.6)	741 (61.0)	1772 (53.8)
Unknown	32 (1.5)	14 (1.1)	46 (1.4)
Use of CT			
No	86 (4.1)	132 (10.9)	218 (6.6)
Yes	1981 (95.2)	1068 (88.0)	3049 (92.6)
Unknown	13 (0.6)	14 (1.1)	27 (0.8)
Type of CT ^b			
Antra + taxane	1473 (74.4)	744 (69.7)	2217 (72.7)
Antra only	300 (15.1)	179 (16.8)	479 (15.7)
Other	208 (10.5)	145 (13.6)	353 (11.6)
Use of platinum ^c			
No	1023 (73.2)	105 (69.5)	1128 (72.9)
Yes	317 (22.7)	38 (25.2)	355 (22.9)
Unknown	57 (4.1)	8 (5.3)	65 (4.2)
Use of ET ^d			
No	51 (9.5)	25 (2.4)	76 (4.9)
Yes	477 (89.0)	982 (96.2)	1459 (93.7)
Unknown	8 (1.5)	14 (1.4)	22 (1.4)
Type of ET ^e			
Tamoxifen alone	165 (34.6)	307 (31.3)	472 (32.4)
Tamoxifen + LHRHa	131 (27.5)	259 (26.4)	390 (26.7)
LHRHa alone	16 (3.4)	16 (1.6)	32 (2.2)
AI + LHRHa	90 (18.9)	210 (21.4)	300 (20.6)
Tamoxifen → AI	61 (12.8)	166 (16.9)	227 (15.6)
Other	7 (1.5)	15 (1.5)	22 (1.5)
Unknown	7 (1.5)	9 (0.9)	16 (1.1)

AI, aromatase inhibitor; BMI, body mass index; CT, chemotherapy; ET, endocrine therapy; HER2, human epidermal growth factor receptor 2; IQR, interquartile range; LHRHa, luteinizing hormone-releasing hormone analogue.

^aTest before diagnosis, carried out any time up to 2 months before the diagnosis of breast cancer; test at diagnosis, carried out from 2 months before and up to 6 months after the diagnosis of breast cancer; test after diagnosis, carried out >6 months after the diagnosis of breast cancer.

^bOnly patients with use of chemotherapy.

^cOnly patients with use of chemotherapy and triple-negative breast cancer.

^dOnly patients with hormone receptor-positive tumors.

^eOnly patients with hormone receptor-positive tumors treated with ET.

events and the rate of OS events for *BRCA1* patients are detailed in [Supplementary Table S12](https://doi.org/10.1016/j.annonc.2025.11.004), available at <https://doi.org/10.1016/j.annonc.2025.11.004>. No difference in DFS was found according to protein truncation ([Figure 2A](https://doi.org/10.1016/j.annonc.2025.11.004) and [Supplementary Table S5](https://doi.org/10.1016/j.annonc.2025.11.004), available at <https://doi.org/10.1016/j.annonc.2025.11.004>). Nonetheless, patients with *BRCA1* protein-truncating variants showed worse OS than patients with non-truncating variants [hazard ratio (HR) 2.00, 95% confidence interval (CI) 1.17-3.41; [Figure 2B](https://doi.org/10.1016/j.annonc.2025.11.004) and [Supplementary Table S6](https://doi.org/10.1016/j.annonc.2025.11.004), available at <https://doi.org/10.1016/j.annonc.2025.11.004>].

In *BRCA2* carriers, a different distribution of protein-truncating variants was observed according to geographic region. No other meaningful associations between baseline clinicopathological variables and protein truncation were observed ([Supplementary Table S13](https://doi.org/10.1016/j.annonc.2025.11.004), available at <https://doi.org/10.1016/j.annonc.2025.11.004>). The rate and type of first DFS events and the rate of OS events for *BRCA2* patients are detailed in [Supplementary Table S14](https://doi.org/10.1016/j.annonc.2025.11.004), available at <https://doi.org/10.1016/j.annonc.2025.11.004>.

[10.1016/j.annonc.2025.11.004](https://doi.org/10.1016/j.annonc.2025.11.004). Protein-truncating variants were associated with numerically worse DFS, without statistical significance ([Figure 2C](https://doi.org/10.1016/j.annonc.2025.11.004) and [Supplementary Table S9](https://doi.org/10.1016/j.annonc.2025.11.004), available at <https://doi.org/10.1016/j.annonc.2025.11.004>). Patients with *BRCA2* protein-truncating variants exhibited numerically worse OS than those with non-truncating variants (HR 6.27, 95% CI 0.86-45.87; [Figure 2D](https://doi.org/10.1016/j.annonc.2025.11.004) and [Supplementary Table S10](https://doi.org/10.1016/j.annonc.2025.11.004), available at <https://doi.org/10.1016/j.annonc.2025.11.004>), although not statistically significant.

Coding consequence (frameshift, inframe deletion, missense, nonsense, splicing variants)

In *BRCA1* carriers, a different distribution of coding consequences was observed according to geographic region. No other meaningful associations between baseline clinicopathological variables and coding consequence were detected ([Supplementary Table S15](https://doi.org/10.1016/j.annonc.2025.11.004), available at <https://doi.org/10.1016/j.annonc.2025.11.004>). The rate and type of first




Figure 1. Disease-free survival (DFS) and overall survival (OS) according to variant type. (A) DFS in *BRCA1* carriers. (B) OS in *BRCA1* carriers. (C) DFS in *BRCA2* carriers. (D) OS in *BRCA2* carriers.

CNV, copy number variations; indel, small deletions or insertions; SNV, single-nucleotide variants.

DFS events and the rate of OS events for *BRCA1* patients are detailed in [Supplementary Table S16](https://doi.org/10.1016/j.annonc.2025.11.004), available at <https://doi.org/10.1016/j.annonc.2025.11.004>. No differences in DFS were identified according to coding consequence ([Figure 3A](https://doi.org/10.1016/j.annonc.2025.11.004) and [Supplementary Table S5](https://doi.org/10.1016/j.annonc.2025.11.004), available at <https://doi.org/10.1016/j.annonc.2025.11.004>). Nevertheless, *BRCA1* missense variants were associated with better OS compared with frameshift variants (HR 0.48, 95% CI 0.28-0.84; [Figure 3B](https://doi.org/10.1016/j.annonc.2025.11.004) and [Supplementary Table S6](https://doi.org/10.1016/j.annonc.2025.11.004), available at <https://doi.org/10.1016/j.annonc.2025.11.004>).

In *BRCA2* carriers, a different distribution of coding consequences was observed according to geographic region. No other meaningful associations between baseline clinicopathological variables and coding consequence were detected ([Supplementary Table S17](https://doi.org/10.1016/j.annonc.2025.11.004), available at <https://doi.org/10.1016/j.annonc.2025.11.004>). The rate and type of first DFS events and the rate of OS events for *BRCA2* patients are detailed in [Supplementary Table S18](https://doi.org/10.1016/j.annonc.2025.11.004), available at <https://doi.org/10.1016/j.annonc.2025.11.004>. No statistically significant differences in DFS were observed according to coding consequence ([Figure 3C](https://doi.org/10.1016/j.annonc.2025.11.004) and [Supplementary Table S9](https://doi.org/10.1016/j.annonc.2025.11.004), available at <https://doi.org/10.1016/j.annonc.2025.11.004>). Nevertheless, *BRCA2* missense variants were associated with better OS compared with splicing variants (HR 0.12, 95% CI 0.02-0.90;

[Figure 3D](https://doi.org/10.1016/j.annonc.2025.11.004) and [Supplementary Table S10](https://doi.org/10.1016/j.annonc.2025.11.004), available at <https://doi.org/10.1016/j.annonc.2025.11.004>).

Variant location (exon and protein domain)

In *BRCA1* carriers, a different distribution of exons involved was observed according to geographic region. No other meaningful association between baseline clinicopathological variables and exons was identified ([Supplementary Table S19](https://doi.org/10.1016/j.annonc.2025.11.004), available at <https://doi.org/10.1016/j.annonc.2025.11.004>). The rate and type of first DFS events and the rate of OS events for *BRCA1* patients are detailed in [Supplementary Table S20](https://doi.org/10.1016/j.annonc.2025.11.004), available at <https://doi.org/10.1016/j.annonc.2025.11.004>. No differences in DFS were observed based on variant location ([Figure 4A](https://doi.org/10.1016/j.annonc.2025.11.004) and [Supplementary Table S5](https://doi.org/10.1016/j.annonc.2025.11.004), available at <https://doi.org/10.1016/j.annonc.2025.11.004>). However, LP/PVs located outside exons 2, 10, and 19 were associated with better OS compared with patients with variants in exon 2 (HR 0.56, 95% CI 0.34-0.94) and exon 10 (HR 0.53, 95% CI 0.38-0.75; [Figure 4B](https://doi.org/10.1016/j.annonc.2025.11.004) and [Supplementary Table S6](https://doi.org/10.1016/j.annonc.2025.11.004), available at <https://doi.org/10.1016/j.annonc.2025.11.004>).

In *BRCA2* carriers, patients with LP/PV located in exon 10 had earlier diagnosis (≤ 30 years) compared with patients with variants in exon 11, exons 15-26, and other exons




Figure 2. Disease-free survival (DFS) and overall survival (OS) according to protein truncation. (A) DFS in *BRCA1* carriers. (B) OS in *BRCA1* carriers. (C) DFS in *BRCA2* carriers. (D) OS in *BRCA2* carriers. NC, not classifiable.




Figure 3. Disease-free survival (DFS) and overall survival (OS) according to coding consequence. (A) DFS in *BRCA1* carriers. (B) OS in *BRCA1* carriers. (C) DFS in *BRCA2* carriers. (D) OS in *BRCA2* carriers.

(26.2% versus 13.7%, 15.4%, 17.2%, respectively) (Supplementary Table S21, available at <https://doi.org/10.1016/j.annonc.2025.11.004>). The rate and type of first DFS events and the rate of OS events for *BRCA2* patients are detailed in Supplementary Table S22, available at <https://doi.org/10.1016/j.annonc.2025.11.004>.

Patients with LP/PV in exons 15-26 showed worse DFS compared with those with LP/PV in exon 10 (HR 1.74, 95% CI 1.02-2.98). Patients with LP/PV in the 'other exons' showed worse DFS compared with those with LP/PV in exon 10 (HR 1.91, 95% CI 1.09-3.35; Figure 4C and Supplementary Table S9, available at <https://doi.org/10.1016/j.annonc.2025.11.004>). No differences in OS were observed according to the exon involved (Figure 4D and Supplementary Table S10, available at <https://doi.org/10.1016/j.annonc.2025.11.004>).

Considering patients with only non-truncating variants, no differences in terms of baseline characteristics, DFS, or OS (Supplementary Tables S5, S6, S23, and S24, available at <https://doi.org/10.1016/j.annonc.2025.11.004>) were detected according to the most represented protein domains of *BRCA1* (BRCT domain versus RING domain). Among *BRCA2* carriers, most of the patients harbored LP/PVs in the OB domain (88%); therefore, no comparisons were carried out.

DISCUSSION

In this global study, the specific type and location of *BRCA* LP/PV seemed not to influence tumor clinicopathological features, although they might have a prognostic role in a cohort of *BRCA* carriers affected by early-onset breast cancer.

The distribution of variant type among patients included in the present study is consistent with that observed in prior breast cancer studies,³⁵ with the majority being indel, followed by SNVs. Regarding the coding consequences, the majority of LP/PVs were frameshift variants in both the *BRCA1* and *BRCA2* genes, with protein-truncating variants being the most common in this study population. Furthermore, as previously described,³⁶ in this study the majority of LP/PVs were located in *BRCA1* exon 10 (formerly exon 11) and *BRCA2* exon 11, which are the largest exons of both *BRCA* genes. Overall, the most frequent variants were the founder mutations *BRCA1* c.5266dup, *BRCA1* c.68_69del, *BRCA2* c.5946del, and *BRCA2* c.2808_2811del, along with *BRCA1* c.181T>G and *BRCA2* c.6275_6276del.³⁷ *BRCA1* c.5266dup, likely originating from Northern Europe 1500-1800 years ago and then entering the Ashkenazi Jewish population more recently, was the most common variant, particularly among carriers from South and East Europe and Israel.³⁸ *BRCA1* c.68_69del was enriched in




Figure 4. Disease-free survival (DFS) and overall survival (OS) according to exon involved. (A) DFS in *BRCA1* carriers. (B) OS in *BRCA1* carriers. (C) DFS in *BRCA2* carriers. (D) OS in *BRCA2* carriers.

North America and Asia, reflecting both Ashkenazi Jewish-associated haplotypes and independent origins in populations such as Pakistanis and Malaysians.³⁹⁻⁴³ *BRCA1* c.181T>G, predominantly observed in South and East Europe, supports its classification as an Eastern European founder variant.⁴⁴ *BRCA2* c.5946del was consistent with its role as a founder variant in Ashkenazi Jewish and Northern European populations, including Iceland,^{37,45,46} whereas *BRCA2* c.2808_2811del showed a recurrent mutational origin across geographically distant populations.⁴⁷⁻⁴⁹ Finally, *BRCA2* c.6275_6276del, previously described in Western populations,³⁷ was highly represented in Asian carriers in our dataset. Notably, we found significantly different distributions of variant type and location across ethnic and geographic groups. This could be partially attributable to population characteristics such as founder effects, but it can also be influenced by the wide heterogeneity in the methods of genetic testing available in each country. It is noteworthy that this study was conducted on a population diagnosed with breast cancer between 2000 and 2020; during this period, the methods for variant detection have significantly evolved, shifting from 'hotspot' searches to increasingly larger panels, and the detection of CNVs has spread only in the last decade.

In this large cohort of *BRCA* carriers with breast cancer, variant type (indel, SNV, CNV) seemed not to be associated with differences in prognosis (DFS or OS). Prior studies showed that *BRCA* CNVs were correlated to a higher breast cancer risk versus non-CNV LP/PVs (i.e. small variants such as nonsense, missense, and frameshift).⁵⁰ Moreover, in a small cohort of young women with triple-negative breast cancer, *BRCA1* CNVs were associated with superior OS rates compared with other LP/PVs.⁵¹ Nevertheless, to date, no other data on their impact on breast cancer phenotype and outcome have been documented.

To the best of our knowledge, this is the first study describing a possible correlation between *BRCA* protein truncation and worse survival in patients with breast cancer. Indeed, the present study showed that patients with protein-truncating variants of *BRCA1* gene display worse OS compared with patients with non-truncating variants. Interestingly, despite comparable overall DFS, patients with *BRCA1* truncating variants showed a higher rate of distant relapses and second primary tumors compared with non-truncating variants, which might have impacted OS. On the other hand, the divergence between DFS and OS among *BRCA1* carriers might also suggest a possible role of *BRCA1* truncating variants in influencing other determinants of life expectancy, such as other health conditions. Nonetheless, data on causes of death other than breast cancer were not collected in the present study; therefore, further research is needed to confirm this hypothesis.

Interestingly, the present study showed that patients with missense variants display better OS compared with patients with frameshift variants in *BRCA1* gene and better OS compared with patients with splicing variants in *BRCA2* gene. In these cases as well, absolute DFS was comparable;

however, patients with *BRCA1* missense LP/PVs showed a lower rate of distant relapses compared with those with *BRCA1* frameshift LP/PVs. Similarly, *BRCA2* missense LP/PVs were associated with lower rates of both locoregional and distant relapses compared with *BRCA2* splicing LP/PVs. The favorable prognosis associated with missense variants was already described by Corso et al.,¹⁷ who found that missense and splicing LP/PVs of *BRCA1/BRCA2* were independently associated with a lower incidence of locoregional recurrence after primary surgery, when compared with indels, nonsense LP/PVs, and structural variants in a multivariable analysis. However, in this previous research, no difference in clinicopathological features or in OS was observed.¹⁷

Patients with LP/PVs located in exon 10 of *BRCA2* exhibited a higher rate of early-onset diagnosis (≤ 30 years of age) than patients with variants in the other exons and these patients also showed better DFS compared with those with variants in most of the other exons. Additionally, patients with LP/PVs located in *BRCA1* exon 2 and exon 10 had worse OS compared with patients with variants in most of the other exons. In this case as well, poorer OS is associated with a higher rate of distant relapses. Studies investigating the relationship between LP/PV location and clinicopathological characteristics or prognosis in breast cancer remain limited. Contrary to the data described here, Rebbeck et al.¹⁴ found that LP/PVs in *BRCA1* exon 10 (formerly exon 11) and *BRCA2* exon 11 were associated with earlier ages at breast and ovarian cancer diagnosis. On the other hand, Bayraktar et al.⁵² did not find any difference in age at breast cancer diagnosis and phenotypic features of breast cancer by ethnicity or variant location. Bayraktar et al.⁵² also found no significant differences in OS in relation to variant location (exons 2, 11, and 20 among *BRCA1* carriers and 10, 11, and 25 among *BRCA2* carriers). Nevertheless, when the OS was analyzed according to three combined exon groups, the 10-year OS estimates were significantly better for *BRCA2* carriers with LP/PVs in exon 11 and with LP/PVs in exons 1-10, compared with LP/PVs in exons 12-25. Further research is needed in order to confirm and better understand the role of variant position in the prognosis of these patients.

Several limitations of this study must be considered when interpreting these results. Firstly, this retrospective study lacked a control cohort of healthy *BRCA* carriers. The study population specifically included patients diagnosed with breast cancer at a young age, excluding patients with diagnosis of ovarian cancer or other malignancies with no history of invasive breast cancer; therefore, these results may not be generalizable to the entire population of *BRCA* carriers. Additionally, family clustering was not collected in this study; therefore, although being relatively unlikely due to the special population included, this cohort might include relatives with the same variant who could also share other clinicopathological features and prognostic factors. Moreover, survival outcomes might have also been influenced by factors that were not included in our investigation (i.e. lifestyle, environmental factors, or

comorbidities), particularly given the regional variation observed in the distribution of *BRCA* variants. Data on treatments at relapse or in the metastatic setting were also not available. However, it is important to note that patients were enrolled between 2000 and 2020, a period during which Cyclin-dependent kinase 4/6 inhibitors, PARP inhibitors, immune checkpoint inhibitors, and antibody–drug conjugates were not yet standard of care in the treatment of patients with early-stage disease. Finally, no correction for multiple testing was applied with subsequent potential increased risk of false-positive findings.

In conclusion, this analysis enhances our understanding of the influence of the specific types of *BRCA1/BRCA2* LP/PVs on the characteristics and outcomes of breast cancer in young women. Specifically, this study identified young *BRCA* carriers affected by breast cancer who may have a better prognosis (*BRCA1* and *BRCA2* missense LP/PVs) and a worse prognosis (*BRCA1* protein-truncating variants, *BRCA1* exons 2 and 10). Notably, the predicted functional consequence, rather than the variant type *per se*, appears to hold greater biological relevance with respect to its impact on gene function and subsequent phenotypic outcomes. A deeper understanding of these variant-specific features will drive future research and support the development of tailored clinical strategies based on individual *BRCA1* or *BRCA2* LP/PV profiles.

ACKNOWLEDGEMENTS

We wish to thank Heather Thorne, Eveline Niedermayr, all the kConFab research nurses and staff, the heads and staff of the Family Cancer Clinics, and the Clinical Follow Up Study (which has received funding from the NHMRC, the National Breast Cancer Foundation, Cancer Australia, and the National Institute of Health (USA)) for their contributions to this resource, and the many families who contribute to kConFab. kConFab is supported by a grant from the National Breast Cancer Foundation, and previously by the National Health and Medical Research Council (NHMRC), the Queensland Cancer Fund, the Cancer Councils of New South Wales, Victoria, Tasmania and South Australia, and the Cancer Foundation of Western Australia. Part of the graphical abstract was designed by pch.vector/ Freepik.

FUNDING

This work was supported by the Italian Association for Cancer Research (AIRC) [grant number MFAG 2020 ID 24698]. ML received support from the European Society for Medical Oncology (ESMO) for a translational research fellowship at Institut Jules Bordet in Brussels, Belgium, at the time this study was initiated. KAP is an Australian National Health and Medical Research Council leadership fellow. Data collection for most Australian participants was through the kConFab Follow-Up Study with support from Cancer Australia and the National Breast Cancer Foundation [grant number PdCCRS 1100868], Cancer Australia [grant number 809195], the Australian National Breast

Cancer Foundation [grant number IF 17], the Australian National Health and Medical Research Council [grant numbers 454508, 288704, 145684], the United States National Institutes of Health [grant number 1R01CA159868], the Queensland Cancer Fund, the Cancer Councils of New South Wales, Victoria, Tasmania, and South Australia, and the Cancer Foundation of Western Australia (no grant number). FJC was supported by the United States National Institutes of Health grant [grant number R35CA253187], a Specialized Program of Research Excellence in breast cancer [grant number P50 CA116201], the Breast Cancer Research Foundation, and the BRCA Research and Cure Alliance (cureBRCA) (no grant number). The study supporters had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; or decision to submit the manuscript for publication.

DISCLOSURE

AT reports advisory role and/or honoraria from Eli/Lilly, Novartis, Pfizer, MSD, AstraZeneca, Gilead, Daiichi Sankyo, Menarini Stemline. EB reports speaker fee from Eli Lilly, AstraZeneca, and funding to the institution from Gilead Science. EA reports advisory role and/or honoraria from Eli Lilly, Sandoz, AstraZeneca, Bayer, Abscint, Gilead, Novartis; research grant to her institution from Gilead BeLux; meeting/travel grants from Novartis, Roche, Eli Lilly, Daiichi Sankyo, AstraZeneca, Abscint, Menarini, Gilead (all outside the submitted work). KP received honoraria for consultations/lectures/training/clinical trials and payment of conference fees from AstraZeneca, Gilead, Roche, Novartis, Eli Lilly, Pfizer, MSD, Swixx, all outside the submitted work. H.C.F.M has research funding paid to Cleveland Clinic from AstraZeneca, Daiichi Sankyo, Sermonix Pharmaceuticals, Seattle Genetics, and Pfizer. WC reports honoraria from AstraZeneca, Pfizer, Merck, Eisai. LDM reports grants or contracts from Eli Lilly, Novartis, Roche, Daiichi Sankyo, Seagen, AstraZeneca, Gilead, Pierre Fabre, Pfizer, Menarini Stemline, MSD, Olema, Exact sciences; Jazz Pharmaceuticals; Astellas. Consulting fees Eli Lilly, Novartis, Roche, Daiichi Sankyo, Seagen, AstraZeneca, Gilead, Pierre Fabre, Pfizer, Menarini Stemline, MSD, Exact sciences, GSK, Ipsen, Eisai. Support for attending meetings and/or travel Roche, Gilead, Menarini Stemline, AstraZeneca, Daiichi Sankyo. Participation on Advisory Board Eli Lilly, Novartis, Roche, Daiichi Sankyo, Seagen, AstraZeneca, Gilead, Pierre Fabre, Pfizer, Menarini Stemline, MSD, Exact sciences, GSK, Ipsen, Eisai, Olema, Signature biosciences. FJC reports an advisory role for AstraZeneca and research support from GRAIL. FP reports honoraria for advisory boards, activities as a speaker, travel grants, research grants from Amgen, AstraZeneca, Daiichi Sankyo, Celgene, Eisai, Eli Lilly, Exact Sciences, Gilead, Ipsen, Italfarmaco, Menarini, MSD, Novartis, Pierre Fabre, Pfizer, Roche, Seagen, Takeda, Viatrix; research funding from AstraZeneca, Eisai, Roche. PAM received consulting fees from Gilead, GSK, MSD, Novartis, Pfizer; speaker honoraria from AstraZeneca, Pfizer; travel/meeting grants from Gilead,

GSK, Pfizer, Roche. AM reports role as speaker (no personal fees) from Roche, Seagen; consultant (no compensation) from AstraZeneca, Roche, Veracyte; research funding (to institution) from AstraZeneca, Veracyte, Novartis, MSD. CG reports travel grants from Ipsen, Gilead and Pfizer; research funding (to institution) from Gilead all outside the submitted work; honoraria from Novartis. RY reports consulting role with Pfizer, consulting role, invited speaker with Novartis, consulting role with Rhenium, invited speaker with Medison, consulting role, invited speaker with MSD, AstraZeneca, and Eli Lilly, consulting role with Gilead, invited speaker, consulting role with Stemline. RSB reports advisory/consulting/speaker fees from Roche, AstraZeneca, Novartis, Lilly, Daiichi Sankyo, Pfizer, Eisai, GlaxoSmithKline, Reveal Genomics, and Gilead; travel expenses from Pfizer, AstraZeneca, Gilead, Novartis, and Roche. LM reports speaker/consultancy fee from Novartis, Daiichi Sankyo, AstraZeneca, Lilly, Pfizer, Roche; support for attending medical conferences from Novartis, Roche, Pfizer, Lilly. AK served as an invited lecturer for AstraZeneca, Pfizer, and Roche. CS reports speaker bureau from AstraZeneca, Daiichi Sankyo; travel/accommodation/expenses from Pfizer, AstraZeneca, Novartis, Lilly. AS reports consulting or advisory role for Eli Lilly, Pfizer, Novartis, Roche, Gilead, MSD, AstraZeneca, Progenetics, Rhenium, Stemline; travel, accommodations, expenses from Neopharm, Celgene, Medison, Roche, MSD, Gilead, Pfizer; speakers bureau for Gilead, Roche, Pfizer, Novartis, Eli Lilly; grant support from Novartis, Roche, Gilead. ML reports advisory role for Roche, Lilly, Novartis, AstraZeneca, Pfizer, Seagen, Gilead, MSD, Exact Sciences, Pierre Fabre, Menarini, Nordic Pharma; speaker honoraria from Roche, Lilly, Novartis, Pfizer, Sandoz, Libbs, Daiichi Sankyo, Takeda, Menarini, AstraZeneca; travel grants from Gilead, Daiichi Sankyo, Roche; research funding (to institution) from Gilead all outside the submitted work. YA reports consulting fees from AstraZeneca and Pfizer all outside of submitted work. NH reports honoraria for lectures and/or consulting: AstraZeneca, Daiichi-Sankyo, Exact Sciences, Gilead, IQVIA, Lilly, Menarini-Stemline, MSD, Novartis, Pierre-Fabre, Pfizer, Roche, Sandoz, Viatrix, Zuelligpharma and minority ownership of West German Study Group (WSG) - all outside the subject area. MKL reports equity and professional services and activities from Biologica. ET reports speaker fee and meeting/travel grants from MDS, SOPHiA Genetics and Devyser. RBM reports honoraria from AstraZeneca, Roche, Pfizer; support for attending meetings from AstraZeneca, Pfizer, Gilead. ZK declares honoraria for educational resources and advisory board, AstraZeneca. AF reports honoraria for advisory boards, invited speaker and travel grant from AstraZeneca and MSD. HW reports his institution received financial compensation on his behalf for advisory boards, lecture fees and/or consultancy fees from Agendia, AstraZeneca, Augustine Therapeutics NV, Boehringer Ingelheim int., Daiichi-Sankyo, Eli Lilly Benelux N.V., Gilead Sciences, Immuteq Limited, MediMix bv, Menarini, Novartis, NV Hict, NV Roche SA, Pfizer SA, PSI Cro AG, Seagen, Stemline Therapeutics, Syneos Health. He received

travel support from Gilead Sciences. All other authors have declared no conflicts of interest.

DATA SHARING

Deidentified individual patient data, data dictionary, and statistical analysis plan will be available for 5 years after publication upon reasonable request to ML (matteo.lambertini@unige.it), after proper revision of the data transfer agreement of each participating center and if ultimately allowed by local ethics committees.

REFERENCES

- Roy R, Chun J, Powell SN. BRCA1 and BRCA2: different roles in a common pathway of genome protection. *Nat Rev Cancer*. 2011;12(1):68-78.
- Fu X, Tan W, Song Q, Pei H, Li J. BRCA1 and breast cancer: molecular mechanisms and therapeutic strategies. *Front Cell Dev Biol*. 2022;10:813457.
- Shahid T, Soroka J, Kong E, et al. Structure and mechanism of action of the BRCA2 breast cancer tumor suppressor. *Nat Struct Mol Biol*. 2014;21(11):962-968.
- Shailani A, Kaur RP, Munshi A. A comprehensive analysis of BRCA2 gene: focus on mechanistic aspects of its functions, spectrum of deleterious mutations, and therapeutic strategies targeting BRCA2-deficient tumors. *Med Oncol*. 2018;35(3):18.
- Maxwell KN, Domchek SM, Nathanson KL, Robson ME. Population frequency of germline BRCA1/2 mutations. *J Clin Oncol*. 2016;34(34):4183-4185.
- Lejeune F. Nonsense-mediated mRNA decay at the crossroads of many cellular pathways. *BMB Rep*. 2017;50(4):175-185.
- Petrosino M, Novak L, Pasquo A, et al. Analysis and Interpretation of the impact of missense variants in cancer. *Int J Mol Sci*. 2021;22(11):5416.
- Scotti MM, Swanson MS. RNA mis-splicing in disease. *Nat Rev Genet*. 2016;17(1):19-32.
- Póczy T, Grolmusz VK, Papp J, Butz H, Patócs A, Bozsik A. Germline structural variations in cancer predisposition genes. *Front Genet*. 2021;12:634217.
- Kuchenbaecker KB, Hopper JL, Barnes DR, et al. Risks of breast, ovarian, and contralateral breast cancer for BRCA1 and BRCA2 mutation carriers. *J Am Med Assoc*. 2017;317(23):2402-2416.
- Deng M, Chen H-H, Zhu X, et al. Prevalence and clinical outcomes of germline mutations in BRCA1/2 and PALB2 genes in 2769 unselected breast cancer patients in China. *Int J Cancer*. 2019;145(6):1517-1528.
- Zattarin E, Tagliatela I, Lobefaro R, et al. Breast cancers arising in subjects with germline BRCA1 or BRCA2 mutations: different biological and clinical entities with potentially diverse therapeutic opportunities. *Crit Rev Oncol Hematol*. 2023;190:104109.
- Lambertini M, Blondeaux E, Tomasello LM, et al. Clinical behavior of breast cancer in young BRCA carriers and prediagnostic awareness of germline BRCA status. *J Clin Oncol*. 2025;43(14):1706-1719.
- Rebeck TR, Mitra N, Wan F, et al. Association of type and location of BRCA1 and BRCA2 mutations with risk of breast and ovarian cancer. *J Am Med Assoc*. 2015;313(13):1347-1361.
- Gayther SA, Mangion J, Russell P, et al. Variation of risks of breast and ovarian cancer associated with different germline mutations of the BRCA2 gene. *Nat Genet*. 1997;15(1):103-105.
- Li H, Engel C, de la Hoya M, et al. Risks of breast and ovarian cancer for women harboring pathogenic missense variants in BRCA1 and BRCA2 compared with those harboring protein truncating variants. *Genet Med*. 2022;24(1):119-129.
- Corso G, Girardi A, Calvillo M, et al. Prognostic impact of germline BRCA1/2 pathogenic variants in breast cancer. *Breast Cancer Res Treat*. 2023;197(1):103-112.
- Pal T, Mundt E, Richardson ME, et al. Reduced penetrance BRCA1 and BRCA2 pathogenic variants in clinical germline genetic testing. *NPJ Precis Oncol*. 2024;8(1):247.

19. Labidi-Galy SI, Rodrigues M, Sandoval JL, et al. Association of location of BRCA1 and BRCA2 mutations with benefit from olaparib and bevacizumab maintenance in high-grade ovarian cancer: phase III PAOLA-1/ENGOT-ov25 trial subgroup exploratory analysis. *Ann Oncol*. 2023;34(2):152-162.
20. Škof E, Stegel V, Dragoš VŠ, et al. Exploring the impact of BRCA1 and BRCA2 mutation type and location on olaparib maintenance therapy in platinum-sensitive relapsed ovarian cancer patients: a single center report. *Gynecol Oncol*. 2024;190:104-112.
21. Lambertini M, Blondeaux E, Agostinetto E, et al. Pregnancy after breast cancer in young BRCA carriers: an international hospital-based cohort study. *J Am Med Assoc*. 2024;331(1):49-59.
22. Blondeaux E, Sonnenblick A, Agostinetto E, et al. Association between risk-reducing surgeries and survival in young BRCA carriers with breast cancer: an international cohort study. *Lancet Oncol*. 2025;26(6):759-770.
23. von Elm E, Altman DG, Egger M, et al. The strengthening the reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. *Lancet*. 2007;370(9596):1453-1457.
24. den Dunnen JT, Dalgleish R, Maglott DR, et al. HGVS recommendations for the description of sequence variants: 2016 update. *Hum Mutat*. 2016;37(6):564-569.
25. Hart RK, Fokkema IFAC, DiStefano M, et al. HGVS nomenclature 2024: improvements to community engagement, usability, and computability. *Genome Med*. 2024;16(1):149.
26. Kopanos C, Tsiolkas V, Kouris A, et al. VarSome: the human genomic variant search engine. *Bioinformatics*. 2019;35(11):1978-1980.
27. Rehman HL, Berg JS, Brooks LD, et al. ClinGen—the clinical genome resource. *N Engl J Med*. 2015;372(23):2235-2242.
28. Richards S, Aziz N, Bale S, et al. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med*. 2015;17(5):405-424.
29. Morales J, Pujar S, Loveland JE, et al. A joint NCBI and EMBL-EBI transcript set for clinical genomics and research. *Nature*. 2022;604(7905):310-315.
30. McDevitt T, Durkie M, Arnold N, et al. Correction: EMQN best practice guidelines for genetic testing in hereditary breast and ovarian cancer. *Eur J Hum Genet*. 2024. <https://doi.org/10.1038/s41431-024-01670-3>.
31. HGVS (Human Genome Variation Society). Numbering recommendations. 2025. Available at <https://hgvs-nomenclature.org/stable/background/numbering/>. Accessed September 1, 2025.
32. Clark SL, Rodriguez AM, Snyder RR, Hankins GDV, Boehning D. Structure-function of the tumor suppressor BRCA1. *Comput Struct Biotechnol J*. 2012;1(1):e201204005.
33. Le HP, Heyer W-D, Liu J. Guardians of the genome: BRCA2 and its partners. *Genes (Basel)*. 2021;12(8):1229.
34. Tolane SM, Garrett-Mayer E, White J, et al. Updated standardized definitions for efficacy end points (STEEP) in adjuvant breast cancer clinical trials: STEEP version 2.0. *J Clin Oncol*. 2021;39(24):2720-2731.
35. Robson M, Im S-A, Senkus E, et al. Olaparib for metastatic breast cancer in patients with a germline BRCA mutation. *N Engl J Med*. 2017;377(6):523-533.
36. Karami F, Mehdipour P. A comprehensive focus on global spectrum of BRCA1 and BRCA2 mutations in breast cancer. *Biomed Res Int*. 2013;2013:928562.
37. Rebbeck TR, Friebel TM, Friedman E, et al. Mutational spectrum in a worldwide study of 29,700 families with BRCA1 or BRCA2 mutations. *Hum Mutat*. 2018;39(5):593-620.
38. Hamel N, Feng B-J, Foretova L, et al. On the origin and diffusion of BRCA1 c.5266dupC (5382insC) in European populations. *Eur J Hum Genet*. 2011;19(3):300-306.
39. Velez C, Palamara PF, Guevara-Aguirre J, et al. The impact of Converso Jews on the genomes of modern Latin Americans. *Hum Genet*. 2012;131(2):251-263.
40. Weitzel JN, Lagos V, Blazer KR, et al. Prevalence of BRCA mutations and founder effect in high-risk Hispanic families. *Cancer Epidemiol Biomarkers Prev*. 2005;14(7):1666-1671.
41. Kadalmani K, Deepa S, Bagavathi S, Anishetty S, Thangaraj K, Gajalakshmi P. Independent origin of 185delAG BRCA1 mutation in an Indian family. *Neoplasma*. 2007;54(1):51-56.
42. Rashid MU, Zaidi A, Torres D, et al. Prevalence of BRCA1 and BRCA2 mutations in Pakistani breast and ovarian cancer patients. *Int J Cancer*. 2006;119(12):2832-2839.
43. Qin Z, Li J, Tam B, et al. Ethnic-specificity, evolution origin and deleteriousness of Asian BRCA variation revealed by over 7500 BRCA variants derived from Asian population. *Int J Cancer*. 2023;152(6):1159-1173.
44. Kaufman B, Laitman Y, Gronwald J, Lubinski J, Friedman E. Haplotype of the C61G BRCA1 mutation in Polish and Jewish individuals. *Genet Test Mol Biomarkers*. 2009;13(4):465-469.
45. Im KM, Kirchhoff T, Wang X, et al. Haplotype structure in Ashkenazi Jewish BRCA1 and BRCA2 mutation carriers. *Hum Genet*. 2011;130(5):685-699.
46. Thorlacius S, Olafsdottir G, Tryggvadottir L, et al. A single BRCA2 mutation in male and female breast cancer families from Iceland with varied cancer phenotypes. *Nat Genet*. 1996;13(1):117-119.
47. Infante M, Durán M, Acedo A, et al. The highly prevalent BRCA2 mutation c.2808_2811del (3036delACAA) is located in a mutational hot-spot and has multiple origins. *Carcinogenesis*. 2013;34(11):2505-2511.
48. Heramb C, Wangenstein T, Grindedal EM, et al. BRCA1 and BRCA2 mutation spectrum - an update on mutation distribution in a large cancer genetics clinic in Norway. *Hered Cancer Clin Pract*. 2018;16:3.
49. Herzog JS, Chavarri-Guerra Y, Castillo D, et al. Genetic epidemiology of BRCA1- and BRCA2-associated cancer across Latin America. *NPJ Breast Cancer*. 2021;7(1):107.
50. Hakkaart C, Pearson JF, Marquart L, et al. Copy number variants as modifiers of breast cancer risk for BRCA1/BRCA2 pathogenic variant carriers. *Commun Biol*. 2022;5(1):1061.
51. Villarreal-Garza C, Ferrigno AS, Aranda-Gutierrez A, et al. Influence of germline BRCA genotype on the survival of patients with triple-negative breast cancer. *Cancer Res Commun*. 2021;1(3):140-147.
52. Bayraktar S, Jackson M, Gutierrez-Barrera AM, et al. Genotype-phenotype correlations by ethnicity and mutation location in BRCA mutation carriers. *Breast J*. 2015;21(3):260-267.