

# World Journal of *Orthopedics*

*World J Orthop* 2023 January 18; 14(1): 1-41



**EDITORIAL**

- 1 Osteoarthritis, an old wine in a new bottle!  
*Muthu S*

**MINIREVIEWS**

- 6 Update on surgical procedures for carpal tunnel syndrome: What is the current evidence and practice? What are the future research directions?  
*Pace V, Marzano F, Placella G*
- 13 Polydactyly: Clinical and molecular manifestations  
*Kyriazis Z, Kollia P, Grivea I, Stefanou N, Sotiriou S, Dailiana ZH*

**META-ANALYSIS**

- 23 Comparative effectiveness of adipose-derived mesenchymal stromal cells in the management of knee osteoarthritis: A meta-analysis  
*Muthu S, Patil SC, Jeyaraman N, Jeyaraman M, Gangadaran P, Rajendran RL, Oh EJ, Khanna M, Chung HY, Ahn BC*

**ABOUT COVER**

Editorial Board Member of *World Journal of Orthopedics*, Amr Abdel-Mordy Kandeel, MD, Assistant Professor, Department of Orthopedics and Traumatology, Faculty of Medicine, Menoufia University, Shbien El-Kom 002-048, Egypt. amr\_kandeel81@yahoo.com

**AIMS AND SCOPE**

The primary aim of *World Journal of Orthopedics (WJO, World J Orthop)* is to provide scholars and readers from various fields of orthopedics with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

*WJO* mainly publishes articles reporting research results and findings obtained in the field of orthopedics and covering a wide range of topics including arthroscopy, bone trauma, bone tumors, hand and foot surgery, joint surgery, orthopedic trauma, osteoarthropathy, osteoporosis, pediatric orthopedics, spinal diseases, spine surgery, and sports medicine.

**INDEXING/ABSTRACTING**

*WJO* is now abstracted and indexed in PubMed, PubMed Central, Emerging Sources Citation Index (Web of Science), Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 edition of Journal Citation Reports® cites the 2021 Journal Citation Indicator (JCI) for *WJO* as 0.62. The *WJO*'s CiteScore for 2021 is 2.4 and Scopus CiteScore rank 2021: Orthopedics and Sports Medicine is 139/284.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: *Ying-Yi Yuan*, Production Department Director: *Xiang Li*, Editorial Office Director: *Jin-Lei Wang*.

**NAME OF JOURNAL**

*World Journal of Orthopedics*

**ISSN**

ISSN 2218-5836 (online)

**LAUNCH DATE**

November 18, 2010

**FREQUENCY**

Monthly

**EDITORS-IN-CHIEF**

Massimiliano Leigheb

**EDITORIAL BOARD MEMBERS**

<http://www.wjgnet.com/2218-5836/editorialboard.htm>

**PUBLICATION DATE**

January 18, 2023

**COPYRIGHT**

© 2023 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>

## Update on surgical procedures for carpal tunnel syndrome: What is the current evidence and practice? What are the future research directions?

Valerio Pace, Fabrizio Marzano, Giacomo Placella

**Specialty type:** Orthopedics

**Provenance and peer review:**

Invited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's scientific quality classification**

Grade A (Excellent): 0

Grade B (Very good): 0

Grade C (Good): C, C

Grade D (Fair): 0

Grade E (Poor): 0

**P-Reviewer:** Anandan H, India;  
Tawonsawatruk T, Thailand

**Received:** September 20, 2022

**Peer-review started:** September 20, 2022

**First decision:** November 25, 2022

**Revised:** December 3, 2022

**Accepted:** December 21, 2022

**Article in press:** December 21, 2022

**Published online:** January 18, 2023



**Valerio Pace**, Department of Trauma & Orthopaedics, AOSP Terni - University of Perugia, Terni 05100, Italy

**Fabrizio Marzano**, Department of Trauma & Orthopaedics, University of Perugia, Perugia 06100, Italy

**Giacomo Placella**, Department of Trauma and Orthopaedics, IRCSS San Raffaele Hospital, Milan 20132, Italy

**Corresponding author:** Valerio Pace, MBBS, Senior Postdoctoral Fellow, Surgeon, Department of Trauma & Orthopaedics, AOSP Terni - University of Perugia, Via Joannuccio, Terni 05100, Italy. [valeripace@doctors.org.uk](mailto:valeripace@doctors.org.uk)

### Abstract

Carpal tunnel syndrome (CTS) is a multifactorial compression neuropathy. It is reported to be very common and rising globally. CTS's treatment varies from conservative measures to surgical treatments. Surgery has shown to be an effective method for more severe cases. However few unclear aspects and room for further research and improvements still remains. We performed a narrative literature review on the most up to date progress and innovation in terms of surgical treatments for CTS. The simple algorithm of leaving the choice of the surgical method to surgeons' preference and experience (together with consideration of patients' related factors) seem to be the best available option, which is supported by the most recent metanalysis and systematic reviews. We suggest that surgeons (unless in presence of precise indications towards endoscopic release) should tend to perform a minimally invasive open approach release, favoring the advantage of a better neurovascular structures visualization (and a consequent higher chance to perform a complete release with long term relief of symptoms) instead of favoring an early reduction (in the first postoperative days) of immobilization and pain. Research towards a universally accepted standardization should be aimed for by the researchers, who have failed to date to sufficiently limit bias and limitations.

**Key Words:** Carpal tunnel; Carpal tunnel release; Transverse ligament; Endoscopic release; Open release

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** After reviewing the most up to date literature, it could be said that evidence of superiority of one technique over the others is lacking from a high level of evidence point of view. Specific advantages and disadvantages of surgical methods can however be taken into account when choosing among treatments. The simple algorithm of leaving the choice of the surgical method to surgeons' preference and experience (together with consideration of patients' related factors) seem to be the best available option, which is supported by the most recent metanalysis and systematic reviews.

**Citation:** Pace V, Marzano F, Placella G. Update on surgical procedures for carpal tunnel syndrome: What is the current evidence and practice? What are the future research directions? *World J Orthop* 2023; 14(1): 6-12

**URL:** <https://www.wjgnet.com/2218-5836/full/v14/i1/6.htm>

**DOI:** <https://dx.doi.org/10.5312/wjo.v14.i1.6>

## INTRODUCTION

Carpal tunnel syndrome (CTS) is a multifactorial neuropathy caused by compression of the median nerve at the carpal tunnel. Its symptoms include pain, tingling or numbness affecting mainly the thumb, index and middle finger (sometimes the ring finger is also involved), with this sensation that could travel up the arm; hand weakness (mainly the thumb's pinching muscles).

The incidence of CTS in the general population is thought to be about 3%-4%; although it could reach 8% in the working population. Figures vary quite significantly among continents and countries, but unequivocally the trends suggest the CTS incidence and subsequent surgical management are rising globally[1-3].

The majority of cases seem to be idiopathic. Other causes such as fractures, infections and systemic diseases should also be taken into account. In fact the primary diseases should be properly assessed and treated, together with CTS in order to achieve complete resolution of symptoms[1-4].

The diagnosis is made with accurate history taking and clinical examination, supported by electromyography (EMG) and nerve conduction studies. Ultrasound has been used to aid diagnosis. Most people with mild to moderate symptoms are initially treated non-operatively. However surgery is thought to provide more effective and durable symptom relief, especially for the most severe cases[4].

Various surgical methods have been proposed and studied, with contradictory results. In fact little agreement on the best surgical procedure to treat CTS has been reached. Most of studies are characterized by significant limitations, which do not allow the clinicians to achieve a consensus on the matter [4,5].

Our aim is to present an update on surgical procedures for carpal tunnel syndrome, highlighting what the current evidence is, with advantages and limitations of the studied surgical methods. The final goal is to provide the most up to date scientific information in order to help the clinicians to maintain good practice with decisions based on the highest possible level of evidence.

## DIAGNOSIS

Accurate physical examination is warranted (after full history taking). It should include Tinel sign and Phalen sign, which are commonly utilized to reveal median nerve compression at the carpus. Durkan's compression test could also be used, but it is not useful to discriminate between symptomatic patients with and without EMG disturbances. The closed fist test is specific in these situations. Two-point discrimination is considered positive if greater than 4 mm. EMG shows that the median nerve transmission rate decreases and the latency period extends beyond normal values. Ultrasounds are sometimes used to aid diagnosis[4-7].

## SURGICAL MANAGEMENT

Surgical management is recommended after failure of conservative measures (splinting, physiotherapy, manual therapy, steroid injections, platelet-rich plasma injections, Kinesio taping, neurodynamic techniques, gabapentin, therapeutic ultrasound, and extracorporeal shockwave therapy) for mild and moderate cases of CTS, or for the most severe cases (numbness in the hand, atrophy of the hand muscles, restricted hand function). No high-grade clinical evidence currently supports specific surgical

indications. Surgery should be aimed at reduce the compression on the median nerve at the carpus, at the level of the transverse ligament. The cause of CTS should also be clearly identified, as the primary disease should be treated and resolved as well. However most of cases are thought to be idiopathic. For those cases, a carpal tunnel release is indicated[4,6-8].

## SURGICAL PROCEDURES

The first reported surgically treated cases of CTS we could found in the literature are the ones presented by Herbert Galloway in 1924. Since then, surgical methods have become more and more common. Many surgical options have been studied and proposed, but in the end the main open uncertainty is about the choice between open surgery and endoscopic assisted surgery. Ultrasound guided methods have also been developed[6-9].

### **Open carpal tunnel release**

Surgeons could use a traditional incision or a mini-incision. The landmarks for the traditional incision are the radial border of the hypothenar muscle, where the incision should be started and extended proximally till the distal wrist crease. 5 cm is the most common length of the incision, but it can be further extended to better visualize the structures. The mini-incision techniques include a transverse incision of about 2 cm on the ulnar side of the wrist stripes or a longitudinal incision starting from the mid-palm and ending at the most proximal portion of the palm. The main issues when using the two latter options are increased complexity, scar pain and increased chance of incomplete transverse ligament release. Other options such as double mini-incisions or other slightly different longitudinal incision options have been presented with promising results. Many studies have proved that all mini-incision technique are safe and provide good results[6,10,11].

After skin incision, the surgeon should go through the palmar fat and fascia, trough which the flexor retinaculum should be visualized. When adequately exposed, it should be completely split longitudinally, with decompression and visualization of the median nerve. Wound closure (skin stitches) and a wound dressing are the last surgical steps. Some authors have presented a method of open release through a small incision using a set of specially designed instruments, retaining advantages of observing the pathology under direct vision and avoiding complications of hazardous injuries to important structures. The instruments consist of a thin metal guide with a groove in the center to accommodate an angled knife holder. The procedure has been performed since 1997 with no complications[10,11].

Open carpal tunnel release (OCTR) has been reported to be a safe procedure overall. Only few cases of wound infections are reported. Scar formation on the palm could also be a complication, especially for traditional size incisions. The palmar nerve branch of the median nerve could also be damaged inadvertently during surgical exposure, as it arises at the distal part of the forearm palmarly and it divides into a medial and lateral branch, passing superficial to the flexor retinaculum of the hand. Another nerve that might be injured is the recurrent motor nerve branch of the median nerve, which normally supplies the thenar muscles. His injury might significantly affect the thumb function[9,11,12].

### **Endoscopic carpal tunnel release**

Single-portal and two-portal endoscopic carpal tunnel release (ECTR) have been reported. The first was introduced in 1986 by Okutsu *et al*[13] who started using the aid of endoscopy to perform carpal ligament release. This technique was used and modified by several other authors in the years, such as Agee and Linvatec[14].

Agee used an entry portal between the palmaris longus and the flexor carpi ulnaris at the level of the proximal wrist striatum. The palmaris longus tendon is then exposed after soft tissue dissection. The endoscope is then placed at the level of the ulnar side of the transverse ligament through the portal, and the ligament is cut from distal to proximal. Not good visualization of the neurovascular structures is the main issue of this approach[14-17].

Linvatec introduced a similar single-portal technique. The main characteristic and difference compared to the previous described technique is that the transverse ligament is cut from proximal to distal[14-17].

Among the two-portal ECTR, the Chow technique is the most widely used method. The first portal is made similarly to Agee method's entry portal, whilst the second portal is made in the palm surface (0.5-0.75 ncm in length) on the bisect line of the angle formed from the distal border of the fully abducted thumb and the third web space and approximately 1 cm proximal to the junction of these lines[18,19].

The surgeon should than push the sleeve till it enters into carpal tunnel from the entry portal and exits through the exit portal with the flexor tendon sliding. Cutting tools are then inserted into the sleeve and surgeons can cut the transverse ligament bilaterally under endoscopic monitoring[18,19].

It is thought that two-portal techniques allow a better visualization of the neurovascular structures and therefore they carry the lowest risk of complications and the higher chance to completely cut the transverse ligament. On the other hand, given the quite distal incision of the Chow technique, we must

say that the risk of injury to the arcus volaris superficialis is higher than in other endoscopic techniques, together with a higher risk of excessive palmar scar formation[15,16,18,19].

### **Comparison among surgical procedures**

A huge amount of work has been carried out and lots of papers published on the matter. Several reviews and metanalysis are also available. From the most up to date evidence it seems that no significant differences exist among the different surgical methods. However it must be said that relevant limitations often bias the results[12,15-17].

Results related to differences among mini-incision and traditional incision surgery are various and contradictory. There are studies reporting good outcomes and lower complication rates with mini-incision techniques. Undoubtedly a smaller approach results in less invasive surgery and this could contribute to better esthetic results both on the short and long run, with potential better patients' satisfaction. However symptoms resolution on the long run is the main objective of the performed surgery, and it is unclear whether open surgery could allow better results due to better visualization of the structures and reduced revision surgery rate. Similar results among mini and conventional approach have also been reported[19-24].

Despite what one could hypothesize regarding potential benefits of endoscopic procedures *vs* open surgery in terms of complication rate, operative time and outcomes and patients' satisfaction, overall data do not clearly highlight such advantages. Moreover most of papers present similar results. More evidence is present with regards to postoperative hand pain and recovery time, in favor of endoscopy procedures, allowing a quicker return to work and his consequent reduction of costs and resources. In fact open surgery (particularly the traditional approach) may prolong the immobilization time and augment postoperative pain and the risk for hypertrophic or hypersensitive scar formation. However the better visualization of the neurovascular structures allowed by the open procedures makes the latter safer from this point of view. However it must be said that most noted nerve injuries were transient, and patients still achieved full recovery after surgery. It should be taken as a worrying sign that often the choice between open and endoscopic surgery is left to surgeons' preference and experience, together with patients' related factors. It seems that a lack of universally accepted evidence on one of the most common syndromes and related surgical management still exist[15,16,20-23].

Among the endoscopic techniques, the majority of the studies have focused their attention on the two-portal technique, whilst fewer studies have reported the results of the Agee's technique. The latter is claimed to be used for his potentials of reducing the higher complication rates of the Chow's technique reported by some authors. Better results in terms of recovery time and return to work has been reported in favor of the single-portal techniques. However the utilization of just one portal could cause a not perfect visualization of the structures (including the transverse ligament), and this could lead to incomplete ligament section and the consequent recurrence of symptoms and the need of revision surgery[22-27].

Intuitively one could relate the endoscopic techniques to mini-incision techniques, as they are based on the same objectives (a smaller approach able to provide good or even better results). In fact surgeons utilizing these techniques aim to a better appearance of the scar (and less complications related to scarring processes) and a quicker recovery, with better patients' satisfaction and acceptance of the procedure. However insufficient evidence with regards of comparison between the two techniques is still present[26-29].

A mention to ultrasound-guided percutaneous carpal tunnel release is needed. Several authors have performed carpal tunnel release with such modality, reporting good results and claiming that it could be an effective treatment for CTS. However the overall level of evidence has been reported by review and metanalysis to be very low, with several studies with at least moderate risk of bias. A low complication rate and fast recovery have often been reported after such procedures[29-32].

Another special mention should be given to epineurotomy and flexor tenosynovectomy. They are often considered as a useful adjunct to the basic surgical procedure, but their indications are still controversial. It seems that tenosynovectomy is only recommended for patients with rheumatic disease (or inflammatory risk factors), or patients undergoing chronic hemodialysis. The intraoperative finding of excessive or abnormal synovial tissues makes a tenosynovectomy also indicated[33-37] **Table 1.**

---

## **CONCLUSION**

There has been controversy regarding the superiority of ECTR over OCTR in the last decades. Many original articles have been published on this issue; moreover, several meta-analyses have compared ECTR with OCTR as treatment options for CTS, but relevant bias and limitations have commonly been reported. Therefore a universal consensus has not been achieved yet, even if CTS is a very common pathology and his surgery is routinely and widely performed[26,27].

Over the years the wrist anatomy knowledge has improved and various surgical instruments and methods have been studied and presented. Sufficient effectiveness has always been reported, but a high quality evidence is lacking.

**Table 1 Advantages of surgical options (risk of hypertrophic or hypersensitive scar formation should be considered secondary, despite temporary discomfort for the patients)**

Advantages of open release	Advantages of endoscopic release
Good outcomes	Good outcomes
Low complication rates	Low complication rate (slightly better than the open release)
Mini-incision: best approach and better patients' satisfaction	Better esthetic results (much smaller scars)
Reduced revision surgery rate	Reduced risk for hypertrophic or hypersensitive scar formation
Better visualization of structures	Reduced postoperative hand pain
Safest approach for neurovascular structures	Reduced recovery time
Possibility to use a mini-incision approach	Quicker return to work
Possibility to perform epineurotomy and flexor tenosynovectomy	Reduced costs and resources

Many studies determined that ECTR was superior to OCTR in terms of higher satisfaction rates, improved key pinch strengths, earlier recovery times, and fewer scar-related complications. This suggests that patients with CTS can be effectively managed with ECTR; however, the possibility of transient nerve injury should be considered. However most of studies are characterized by significant limitations, which do not allow the clinicians to achieve a consensus on the matter. Clear and high level of evidence advantages of one technique over the others have not been provided yet, and sufficient results seem to be provided with any of the studied methods[4,5,26-29,38].

We believe that the lack of high level of evidence regarding the surgical techniques should be taken as a worrying sign, especially because clear evidence is most of the time provided for the most common diseases and related management options. In fact it seems that often the choice between open and endoscopic surgery is left to surgeons' preference and experience, together with patients' related factors. To date, this simple algorithm has shown to be able to provide the best results. Surgeons should refrain from attempting potentially less invasive procedures if not familiar with the technique, as the risks of arming would outweigh the potential better results.

Relevant authors tend to strongly recommend their proposed technique, commonly providing evidence of excellent results and elements in favor of their method over the other ones. However when it comes to systematic reviews and metaanalysis, data suggest that relevant bias and limitations do not allow standardization and do not provide sufficient evidence of the superiority of a technique over the others.

We suggest that surgeons (unless in presence of precise indications towards endoscopic release) should tend to perform a minimally invasive open approach release, favoring the advantage of a better neurovascular structures visualization (and a consequent higher chance to perform a complete release with long term relief of symptoms) instead of favoring an early reduction (in the first postoperative days) of immobilization and pain. Moreover, in view of higher chances to obtain long term symptoms relief, the risk of hypertrophic or hypersensitive scar formation should be considered secondary, despite temporary discomfort for the patients [Table 1](#).

We believe there is room for further research evidence, possibly high level of evidence works (level 1 or 2) which should separately study specific aspects and provide detailed and clear advantages and disadvantages of every single treatment option. Cohorts should be sufficiently big and bias reduced to the minimum. The problem of the current heterogeneity should also be overcome.

CTS is becoming more and more common, as is its surgical management. This constitutes a significant economic burden for societies. All surgical techniques have provided satisfactory results and have been proven to be effective options. After reviewing the most up to date literature, it could be said that evidence of superiority of one technique over the others is lacking from a high level of evidence point of view. Specific advantages and disadvantages of surgical methods can however be taken into account when choosing among treatments. The simple algorithm of leaving the choice of the surgical method to surgeons' preference and experience (together with consideration of patients' related factors) seem to be the best available option, which is supported by the most recent metaanalysis and systematic reviews. Research towards a universally accepted standardization should be aimed for by the authors, who have failed to date to sufficiently limit bias and limitations.

## FOOTNOTES

**Author contributions:** Pace V, Marzano F, and Placella G designed the article, performed research and literature review, analyzed data and wrote the review.



**Conflict-of-interest statement:** All the authors declare that they have no conflict of interest.

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country/Territory of origin:** United Kingdom

**ORCID number:** Valerio Pace [0000-0002-4499-9157](https://orcid.org/0000-0002-4499-9157).

**S-Editor:** Liu JH

**L-Editor:** A

**P-Editor:** Liu JH

## REFERENCES

- 1 **Atroshi I**, Gummesson C, Johnsson R, Ornstein E, Ranstam J, Rosén I. Prevalence of carpal tunnel syndrome in a general population. *JAMA* 1999; **282**: 153-158 [PMID: [10411196](https://pubmed.ncbi.nlm.nih.gov/10411196/) DOI: [10.1001/jama.282.2.153](https://doi.org/10.1001/jama.282.2.153)]
- 2 **Centers for Disease Control and Prevention.** Morbidity and Mortality Weekly Report, 2011. December 23, 2011; 60: 1697–1728. Available from: <https://www.cdc.gov/mmwr/index2011.html>
- 3 **Dale AM**, Harris-Adamson C, Rempel D, Gerr F, Hegmann K, Silverstein B, Burt S, Garg A, Kapellusch J, Merlino L, Thiese MS, Eisen EA, Evanoff B. Prevalence and incidence of carpal tunnel syndrome in US working populations: pooled analysis of six prospective studies. *Scand J Work Environ Health* 2013; **39**: 495-505 [PMID: [23423472](https://pubmed.ncbi.nlm.nih.gov/23423472/) DOI: [10.5271/sjweh.3351](https://doi.org/10.5271/sjweh.3351)]
- 4 **Karjalainen T**, Raatikainen S, Jaatinen K, Lusa V. Update on Efficacy of Conservative Treatments for Carpal Tunnel Syndrome. *J Clin Med* 2022; **11** [PMID: [35207222](https://pubmed.ncbi.nlm.nih.gov/35207222/) DOI: [10.3390/jcm11040950](https://doi.org/10.3390/jcm11040950)]
- 5 **MacDonald E**, Rea PM. A Systematic Review of Randomised Control Trials Evaluating the Efficacy and Safety of Open and Endoscopic Carpal Tunnel Release. *Adv Exp Med Biol* 2022; **1356**: 141-172 [PMID: [35146621](https://pubmed.ncbi.nlm.nih.gov/35146621/) DOI: [10.1007/978-3-030-87779-8\\_7](https://doi.org/10.1007/978-3-030-87779-8_7)]
- 6 **Meyers A**, Annunziata MJ, Rampazzo A, Bassiri Gharb B. A Systematic Review of the Outcomes of Carpal Ligament Release in Severe Carpal Tunnel Syndrome. *J Hand Surg Am* 2022 [PMID: [35058091](https://pubmed.ncbi.nlm.nih.gov/35058091/) DOI: [10.1016/j.jhsa.2021.11.015](https://doi.org/10.1016/j.jhsa.2021.11.015)]
- 7 **De Smet L**, Steenwerckx A, Van den Bogaert G, Cnudde P, Fabry G. Value of clinical provocative tests in carpal tunnel syndrome. *Acta Orthop Belg* 1995; **61**: 177-182 [PMID: [8525813](https://pubmed.ncbi.nlm.nih.gov/8525813/)]
- 8 **Piazzini DB**, Aprile I, Ferrara PE, Bertolini C, Tonali P, Maggi L, Rabini A, Piantelli S, Padua L. A systematic review of conservative treatment of carpal tunnel syndrome. *Clin Rehabil* 2007; **21**: 299-314 [PMID: [17613571](https://pubmed.ncbi.nlm.nih.gov/17613571/) DOI: [10.1177/0269215507077294](https://doi.org/10.1177/0269215507077294)]
- 9 **Wu PT**, Chern TC, Wu TT, Shao CJ, Wu KC, Kuo LC, Jou IM. Safe Zones for Percutaneous Carpal Tunnel Release. *Hand Clin* 2022; **38**: 83-90 [PMID: [34802612](https://pubmed.ncbi.nlm.nih.gov/34802612/) DOI: [10.1016/j.hcl.2021.08.008](https://doi.org/10.1016/j.hcl.2021.08.008)]
- 10 **Cellocco P**, Rossi C, El Boustany S, Di Tanna GL, Costanzo G. Minimally invasive carpal tunnel release. *Orthop Clin North Am* 2009; **40**: 441-448, vii [PMID: [19773048](https://pubmed.ncbi.nlm.nih.gov/19773048/) DOI: [10.1016/j.ocl.2009.06.002](https://doi.org/10.1016/j.ocl.2009.06.002)]
- 11 **Kim PT**, Lee HJ, Kim TG, Jeon IH. Current approaches for carpal tunnel syndrome. *Clin Orthop Surg* 2014; **6**: 253-257 [PMID: [25177448](https://pubmed.ncbi.nlm.nih.gov/25177448/) DOI: [10.4055/cios.2014.6.3.253](https://doi.org/10.4055/cios.2014.6.3.253)]
- 12 **Pripotnev S**, Mackinnon SE. Revision of Carpal Tunnel Surgery. *J Clin Med* 2022; **11** [PMID: [35268477](https://pubmed.ncbi.nlm.nih.gov/35268477/) DOI: [10.3390/jcm11051386](https://doi.org/10.3390/jcm11051386)]
- 13 **Okutsu I**, Ninomiya S, Takatori Y, Ugawa Y. Endoscopic management of carpal tunnel syndrome. *Arthroscopy* 1989; **5**: 11-18 [PMID: [2706046](https://pubmed.ncbi.nlm.nih.gov/2706046/) DOI: [10.1016/0749-8063\(89\)90084-4](https://doi.org/10.1016/0749-8063(89)90084-4)]
- 14 **Hansen TB.** [Endoscopic decompression of carpal tunnel syndrome with the "Agee one-portal system"]. *Ugeskr Laeger* 1997; **159**: 5672-5674 [PMID: [9340875](https://pubmed.ncbi.nlm.nih.gov/9340875/)]
- 15 **Orhurhu V**, Orman S, Peck J, Urits I, Orhurhu MS, Jones MR, Manchikanti L, Kaye AD, Odonkor C, Hirji S, Cornett EM, Imani F, Varrassi G, Viswanath O. Carpal Tunnel Release Surgery- A Systematic Review of Open and Endoscopic Approaches. *Anesth Pain Med* 2020; **10**: e112291 [PMID: [34150584](https://pubmed.ncbi.nlm.nih.gov/34150584/) DOI: [10.5812/aapm.112291](https://doi.org/10.5812/aapm.112291)]
- 16 **Karamanos E**, Jillian BQ, Person D. Endoscopic Carpal Tunnel Release: Indications, Technique, and Outcomes. *Orthop Clin North Am* 2020; **51**: 361-368 [PMID: [32498954](https://pubmed.ncbi.nlm.nih.gov/32498954/) DOI: [10.1016/j.ocl.2020.02.001](https://doi.org/10.1016/j.ocl.2020.02.001)]
- 17 **Tulipan JE**, Ilyas AM. Carpal Tunnel Syndrome Surgery: What You Should Know. *Plast Reconstr Surg Glob Open* 2020; **8**: e2692 [PMID: [32537349](https://pubmed.ncbi.nlm.nih.gov/32537349/) DOI: [10.1097/GOX.0000000000002692](https://doi.org/10.1097/GOX.0000000000002692)]
- 18 **Chow JC.** Endoscopic carpal tunnel release. Two-portal technique. *Hand Clin* 1994; **10**: 637-646 [PMID: [7868631](https://pubmed.ncbi.nlm.nih.gov/7868631/)]
- 19 **Chow JC**, Hantes ME. Endoscopic carpal tunnel release: thirteen years' experience with the Chow technique. *J Hand Surg Am* 2002; **27**: 1011-1018 [PMID: [12457351](https://pubmed.ncbi.nlm.nih.gov/12457351/) DOI: [10.1053/jhsu.2002.35884](https://doi.org/10.1053/jhsu.2002.35884)]
- 20 **Zuo D**, Zhou Z, Wang H, Liao Y, Zheng L, Hua Y, Cai Z. Endoscopic versus open carpal tunnel release for idiopathic carpal tunnel syndrome: a meta-analysis of randomized controlled trials. *J Orthop Surg Res* 2015; **10**: 12 [PMID: [25627324](https://pubmed.ncbi.nlm.nih.gov/25627324/) DOI: [10.1186/s13018-014-0148-6](https://doi.org/10.1186/s13018-014-0148-6)]
- 21 **de Roo SF**, Sprangers PN, Walbeehm ET, van der Heijden B. Systematic review and meta-analysis of surgical options for recurrent or persistent carpal tunnel syndrome: simple decompression versus coverage of the median nerve. *J Hand Surg*

- Eur Vol* 2021; **46**: 749-753 [PMID: 33775163 DOI: 10.1177/17531934211001715]
- 22 **Vasiliadis HS**, Nikolakopoulou A, Shrier I, Lunn MP, Brassington R, Scholten RJ, Salanti G. Endoscopic and Open Release Similarly Safe for the Treatment of Carpal Tunnel Syndrome. A Systematic Review and Meta-Analysis. *PLoS One* 2015; **10**: e0143683 [PMID: 26674211 DOI: 10.1371/journal.pone.0143683]
  - 23 **Aslani HR**, Alizadeh K, Eajazi A, Karimi A, Karimi MH, Zaferani Z, Hosseini Khameneh SM. Comparison of carpal tunnel release with three different techniques. *Clin Neurol Neurosurg* 2012; **114**: 965-968 [PMID: 22421246 DOI: 10.1016/j.clineuro.2012.02.017]
  - 24 **Bai J**, Kong L, Zhao H, Yu K, Zhang B, Zhang J, Tian D. Carpal tunnel release with a new mini-incision approach versus a conventional approach, a retrospective cohort study. *Int J Surg* 2018; **52**: 105-109 [PMID: 29471152 DOI: 10.1016/j.ijssu.2018.02.033]
  - 25 **Palmer DH**, Paulson JC, Lane-Larsen CL, Peulen VK, Olson JD. Endoscopic carpal tunnel release: a comparison of two techniques with open release. *Arthroscopy* 1993; **9**: 498-508 [PMID: 8280321 DOI: 10.1016/s0749-8063(05)80396-2]
  - 26 **Li Y**, Luo W, Wu G, Cui S, Zhang Z, Gu X. Open versus endoscopic carpal tunnel release: a systematic review and meta-analysis of randomized controlled trials. *BMC Musculoskelet Disord* 2020; **21**: 272 [PMID: 32340621 DOI: 10.1186/s12891-020-03306-1]
  - 27 **Müller LP**, Rudig L, Degreif J, Rommens PM. Endoscopic carpal tunnel release: results with special consideration to possible complications. *Knee Surg Sports Traumatol Arthrosc* 2000; **8**: 166-172 [PMID: 10883429 DOI: 10.1007/s001670050209]
  - 28 **Kang HJ**, Koh IH, Lee TJ, Choi YR. Endoscopic carpal tunnel release is preferred over mini-open despite similar outcome: a randomized trial. *Clin Orthop Relat Res* 2013; **471**: 1548-1554 [PMID: 23100191 DOI: 10.1007/s11999-012-2666-z]
  - 29 **Sanati KA**, Mansouri M, Macdonald D, Ghafghazi S, Macdonald E, Yadegarfar G. Surgical techniques and return to work following carpal tunnel release: a systematic review and meta-analysis. *J Occup Rehabil* 2011; **21**: 474-481 [PMID: 21528400 DOI: 10.1007/s10926-011-9310-8]
  - 30 **Chou RC**, Robinson DM, Homer S. Ultrasound-guided percutaneous carpal tunnel release: A systematic review. *PM R* 2022 [PMID: 35254722 DOI: 10.1002/pmjr.12801]
  - 31 **David I**. Sonography-Guided Carpal Tunnel Release. *Hand Clin* 2022; **38**: 75-82 [PMID: 34802611 DOI: 10.1016/j.hcl.2021.08.007]
  - 32 **Moungondo F**, Feipel V. Percutaneous Sonographically Guided Release of Carpal Tunnel and Trigger Finger: Biomechanics, Clinical Results, Technical Developments. *Hand Clin* 2022; **38**: 91-100 [PMID: 34802613 DOI: 10.1016/j.hcl.2021.08.010]
  - 33 **Lanzetti RM**, Astone A, Pace V, D'Abbondanza L, Braghiroli L, Lupariello D, Altissimi M, Vadalà A, Spoliti M, Topa D, Perugia D, Caraffa A. Neurolysis versus anterior transposition of the ulnar nerve in cubital tunnel syndrome: a 12 years single secondary specialist centre experience. *Musculoskelet Surg* 2021; **105**: 69-74 [PMID: 32036564 DOI: 10.1007/s12306-020-00647-x]
  - 34 **Pace V**, Lanzetti RM, Venditto T, Park C, Kim WJ, Rinonapoli G, Caraffa A. Dorsally displaced distal radius fractures: introduction of Pacetti's line as radiological measurement to predict dorsal fracture displacement. *Acta Biomed* 2021; **92**: e2021200 [PMID: 34212906 DOI: 10.23750/abm.v92i3.11392]
  - 35 **Valerio Pace**, Pasquale Sessa, Matteo Guzzini, Marco Spoliti, Alessandro Carcangiu, Criseo N, Alessandro Giai Via, Luigi Meccariello, Auro Caraffa, Riccardo Maria Lanzetti. Clinical, functional and radiological outcomes of the use of fixed angle volar locking plates in corrective distal radius osteotomy for fracture malunion. *Acta Biomed* 2021; **92**: e2021180 [PMID: 34212911 DOI: 10.23750/abm.v92i3.11265]
  - 36 **Ting J**, Weiland AJ. Role of ancillary procedures in surgical management of carpal tunnel syndrome: epineurotomy, internal neurolysis, tenosynovectomy, and tendon transfers. *Hand Clin* 2002; **18**: 315-323 [PMID: 12371034 DOI: 10.1016/s0749-0712(02)00027-6]
  - 37 **Chiang J**, An VVG, Graham D, Lawson R, Sivakumar B. Flexor Synovectomy as an Adjunct to Carpal Tunnel Release in Primary Carpal Tunnel Syndrome: A Meta-Analysis. *J Hand Surg Asian Pac Vol* 2021; **26**: 497-501 [PMID: 34789113 DOI: 10.1142/S2424835521500454]
  - 38 **Iwase Y**, Ikai T, Hara A, Mori K, Kusonose K. Carpal tunnel release through a small incision using a special knife guide. *Tech Hand Up Extrem Surg* 2002; **6**: 193-195 [PMID: 16520600 DOI: 10.1097/00130911-200212000-00006]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

**Telephone:** +1-925-3991568

**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)

**Help Desk:** <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

